


# CMSC 754 - Computational Geometry

## Lecture II: Delaunay Triangulations (Properties)

Last lecture - Voronoi Diagrams  
This - The dual structure - Delaunay Triangulations



### Delaunay Triangulation:


Given a set  $P = \{p_1, \dots, p_n\}$  of sites in  $\mathbb{R}^2$ , the Delaunay Triangulation is the cell complex whose vertices are sites & there is an edge  $\overline{p_i p_j}$  iff  $V(p_i) \cap V(p_j)$  share a common edge. Called  $DT(P)$

### Properties:

**Triangulation:** If general position (no four sites cocircular), the internal faces are all triangles

**Hull:** The boundary of the external face is the boundary of  $\text{conv}(P)$


**Circumcircle:** The circumcircle of any triangle is empty (no sites in its interior)



**Empty Circle:** Sites  $p_i$  &  $p_j$  are adjacent in  $\text{DT}(P)$  iff there is an empty circle through  $p_i$  &  $p_j$ .


**Closest Pair:** The **closest pair** of sites are **Delannay neighbors**

- Consider the circle with diameter  $\overline{p_i p_j}$ .
- No site  $p_k$  can lie within
- Apply empty circle prop.



## Combinatorial Complexity:

By applying Euler's formula, there are at most  $2n$  triangles and at most  $3n$  edges




[In  $\mathbb{R}^d$ , size is  $\mathcal{O}(n^{\lfloor d/2 \rfloor})$ ]

## Euclidean Minimum Spanning Tree: (EMST)

Euclidean graph: Complete graph on vertex set  $P = \{p_1, \dots, p_n\}$ , where edge weight is Euclidean distance ( $w(p_i, p_j) = \|p_i - p_j\|$ )


EMST( $P$ ) = MST of Euclidean graph  
(lowest weight tree spanning  $P$ )

Thm: EMST( $P$ )  $\subseteq$  DT( $P$ )



EMST( $P$ )

$\subseteq$



DT( $P$ )

Proof: (Contradiction)

- Suppose some edge  $\overline{ab} \in \text{EMST}(P)$   
but not in  $\text{DT}(P)$


- Empty circle  $\Rightarrow$  circle with diameter  $\overline{ab}$  contains site  $c$

-  $\|ac\| < \|ab\|$

$\|bc\| < \|ab\|$

- Can remove  $\overline{ab}$  from EMST + replace  
with either  $\overline{ac}$  or  $\overline{bc}$  to produce  
a spanning tree of lower weight

- Contradiction!



□

Minimum Weight Triangulation: No!

$\text{MWT}(P)$  = triangulation of  $P$  whose  
sum of edge lengths is minimum

Generally  $\text{MWT}(P) \neq \text{DT}(P)$

**Notation:** Given graph  $G=(V,E)$  and vertices  $u,v \in V$ , let  $d_G(u,v)$  = shortest path distance in  $G$  from  $u$  to  $v$ .

### Spanner Properties:

Given a graph  $G$  and  $t \geq 1$ , a  $t$ -spanner is a subgraph  $G'$  of  $G$  on same vertex set s.t.  $\forall u,v \in V$ ,

$$d_{G'}(u,v) \leq t \cdot d_G(u,v)$$


(Path lengths don't stretch too much)

**Theorem (Keil + Gutwin, '92)** Given a set  $P$  of sites in the plane,  $DT(P)$  is a  $\frac{4\pi\sqrt{3}}{9} \approx 2.418$  spanner of the Euclidean graph. That is,  $\forall p,q \in P$

$$d_{DT(P)}(p,q) \leq \frac{4\pi\sqrt{3}}{9} \cdot \|p-q\|$$


Avoids skinny triangles:

Let  $P$  be a set of sites in the plane. Among all possible triangulations of  $P$ ,  $DT(P)$  maximizes the size of the smallest angle.



other triangulation

smallest angle




$DT(P)$

**Thm:** If all angles of all triangulations are ordered small to large,  $DT(P)$  is the largest lexicographically compared to all triangulations of  $P$ .

(See full lecture notes)

# Relationship to polytopes in $\mathbb{R}^{d+1}$



Delaunay triangulation in  $\mathbb{R}^d$  is the projection of a lower convex hull in  $\mathbb{R}^{d+1}$


Voronoi diagram in  $\mathbb{R}^d$  is the projection of a lower envelope of hyperplanes in  $\mathbb{R}^{d+1}$

→ We'll prove the first only:  $\mathbb{R}^2 \rightarrow \mathbb{R}^3$

Consider the paraboloid:

$$z = f(x, y) = x^2 + y^2$$

Given  $p = (p_x, p_y)$ , define  $p^\uparrow$  to be  $(p_x, p_y, p_x^2 + p_y^2)$



### Lemma:

Three pts  $p, q, r \in \mathbb{R}^2$   
have an empty  
circumcircle w.r.t.  $P$

$\Leftrightarrow$

Three pts  $p^\uparrow, q^\uparrow, r^\uparrow$   
lie on plane  $h$   
with all pts of  $P^\uparrow$   
above

- Let  $c = (c_x, c_y)$  be center of circumcircle through  $p, q, r$  + let  $r$  be its radius
- The plane tangent to paraboloid at  $c^\uparrow$  is:

$$z = 2c_x \cdot x + 2c_y \cdot y - (c_x^2 + c_y^2)$$

- Shift this plane up by distance  $r^2$ :

$$h: z = 2c_x \cdot x + 2c_y \cdot y - (c_x^2 + c_y^2) + r^2$$



- All 3 lifted pts lie on this plane:

$$P_x \text{ on circle: } (p_x - c_x)^2 + (p_y - c_y)^2 = r^2$$

$$\Leftrightarrow (p_x^2 - 2p_x c_x + c_x^2) + (p_y^2 - 2p_y c_y + c_y^2) = r^2$$

$$\Leftrightarrow p_x^2 + p_y^2 = 2c_x \cdot p_x + 2c_y \cdot p_y - (c_x^2 + c_y^2) + r^2$$

$$\Leftrightarrow p_z^\uparrow = 2c_x \cdot p_x^\uparrow + 2c_y \cdot p_y^\uparrow - (c_x^2 + c_y^2) + r^2$$

$\Leftrightarrow p^\uparrow$  lies on plane  $h$