


CMSC 754 - Computational Geometry

Lecture 8 - Trapezoidal Maps


Trapezoidal Maps:

- Given a set $S = \{s_1, \dots, s_n\}$ of line segments in \mathbb{R}^2 , which we assume do not intersect (except at their endpoints)
- General position: No duplicate x-coords + no vertical segments
- Enclose in large bounding rectangle
- Shoot a bullet path vertically above + below each endpoint until it hits something

S



$T(S)$




- This subdivides the rectangle into trapezoids (degenerating possibly to triangles)

Lemma: If $|S| = n$, $T(S)$ has $\leq 6n + 4$ vertices and $\leq 3n + 1$ trapezoids


Proof:

- Each segment contributes 6 vertices to $T(S)$




Plus 4 for the bounding rectangle $\Rightarrow 6n + 4$ vertices

- Charge each trapezoid to vertex on its left side. Each segment is charged 3 times



□

Obs: Each trapezoid owes its existence to ≤ 4 segments



Construction:


- Plane sweep - $O(n \log n)$ [exercise]
- Randomized incremental - $O(n \log n)$ [this lect]

Incremental Construction:

Random order


- Add segments one-by-one
- Update the map after each insertion
- Two types of updates:
 - Endpt of new segment
 - shoot bullet paths up + down
 - Crash through a vertical wall
 - trim the wall back

Endpoint:




Find the trapezoid containing this endpoint, and add vertical segments to top + bottom

Crash through wall:



Determine whether the shooting vertex is above or below, and trim away the excess

These updates implicitly generate new trapezoids + destroy old ones



Running time: to insert segment $s_i, i=1, \dots, n$

- Find trapezoid containing s_i 's left endpoint } $O(\log n)$
(next lect.)


- Trace segment through trapezoids + update } Depends!
 $O(1) \dots O(n)$

Lemma: For $1 \leq i \leq n$, let k_i denote the number of new trapezoids created by insertion of i^{th} segment. Ignoring the time to locate the left endpoint, the insert time is $O(k_i)$

(Note: k_i is a random variable, depending on insertion order $O(1) \dots O(n)$)

Proof: Let w_i denote num. of walls hit.

$$k_i = 4 + w_i$$



$$\text{Insert time} \sim O(2 + 2 + w_i) = O(k_i)$$

Bullets
for left
end pt

Bullets
for right

Trim walls
hit

□

Overall run time: (Ignoring endpt location)

Worst-case: Adding segment i can create $O(i)$ new trapezoids

$$\Rightarrow W(n) = \sum_{i=1}^n i = O(n^2)$$

Expected-case: We will show that if segs are inserted in random order, $E(k_i) = O(1)$

Wow - This does not depend on i !

$$\Rightarrow \text{Exp. time} = \sum_{i=1}^n E(k_i) \leq n \cdot O(1) = O(n)$$

(ignoring left
endpt location)

Lemma: Assuming segments are inserted in random order, $E(k_i)$ (the expected number of new trapezoids with i^{th} insert) is $O(1)$.

\mathcal{T}_i does not depend on insert order

Proof: (Backwards analysis)


- Let $\mathcal{T}_i =$ trapezoidal map after $S_i = \{s_1, s_2, \dots, s_i\}$

- Each seg. is equally likely to be last


$$\text{Prob}(s_i \text{ is last inserted}) = 1/i$$

- Given any trapezoid $\Delta \in \mathcal{T}_i$ and any segment $s \in \{s_1, \dots, s_n\}$ we say Δ depends on s if Δ would have been created if s was inserted last.

$$\delta(\Delta, s) = \begin{cases} 1 & \text{if } \Delta \text{ depends on } s \\ 0 & \text{ow.} \end{cases}$$



Trapezoids that depend on s



Segments on which Δ depends

Note: $\delta(\Delta, s)$ does not depend on insertion order


$$\bar{E}(k_i) = \sum_{s \in S_i} \text{Prob}(s \text{ inserted last}) \cdot \left(\begin{array}{l} \text{Num. of traps} \\ \text{that depend on} \\ s \end{array} \right)$$

$$= \sum_{s \in S_i} \left(\frac{1}{i} \right) \sum_{\Delta \in \mathcal{T}_i} \delta(\Delta, s)$$

(Reverse sum. order)

$$= \frac{1}{i} \sum_{s \in S_i} \sum_{\Delta \in \mathcal{T}_i} \delta(\Delta, s)$$

$$= \frac{1}{i} \sum_{\Delta \in \mathcal{T}_i} \sum_{s \in S_i} \delta(\Delta, s)$$



$$\leq \frac{1}{i} \sum_{\Delta \in \mathcal{T}_i} 4$$

Δ depends on at most 4 segs

$$= \frac{4}{i} \cdot (\text{No. of trapezoids in } \mathcal{T}_i)$$

$$\leq \frac{4}{i} (3i + 1) \quad [\text{By earlier lemma}]$$

$$= 12 + \frac{4}{i} = \mathcal{O}(1)$$

□

Summary:

- Shown that if segs. inserted in random order, total no. of updates $\mathcal{O}(n)$
- Next: How to locate left endpoints.