

2022, January 15th

Ramsey theory

by Federica Baccini

a.y. 2021/2022

What is Ramsey theory?

«According to a 3500-year-old cuneiform text, an ancient Sumerian scholar once looked to the stars in the heavens and saw a lion, a bull and a scorpion. A modern astronomer would be more likely to describe a constellation as a temporary collection of stars, which we earthlings observe from the edge of an ordinary galaxy. Yet most stargazers would agree that the night sky appears to be filled with constellations in the shape of straight lines, rectangles and pentagons. Could it be that such geometric patterns arise from unknown forces in the cosmos?

Mathematics provides a much more plausible explanation.»

(1990, Graham R. L., Spencer J.H.)

What is Ramsey theory?

- Frank Plumpton Ramsey (Cambridge, 1903-Londra, 1930) proved that patterns like those we see in the stars are always present in any large enough structure.
- Ramsey theory is an entire branch of mathematics whose purpose is to find ordered substructure in arbitrary structures.
- First contribution to Ramsey theory given by Ramsey in 1928; other contributions given by the Hungarian mathematicians Paul Erdős (Budapest, 1913-Warsaw, 1996) and George Szekeres (Budapest, 1911-Adelaide, 2005).

Ramsey Theory and graphs: a historical example



- William Gasarch, a computer scientist, studied the work of a scholar of Pre-Christian England, named Sir Woodsor Kneading, who observed a strange phenomenon.
- 577 b. C. : 5 lords in Wessex, all with armies. No war.
- A sixth lord settled and war broke out.
- 552 b. C. : 5 lords in what is now Sout Wales. No war.
- A sixth lord settled and war broke out.
- Phenomenon observed 42 times by Kneading.


Why do six lords always mean war?

One of the following holds:

1. There are 3, 4 or 5 allies, and they attack the other kingdoms.
2. There are 3 or more of them who are pairwise enemies, so a «pairwise war» breaks out.
3. (Rare case) All the 6 kingdoms form an alliance and there is peace.

Conjecture: whenever there are 6 lords, not all in alliance, there must either be 3, 4 or 5 lords allied one with each other, or 3, 4, 5 or 6 lords who are pairwise enemies.

Formalizing the problem in terms of graphs



- Complete graph (K_6), whose edges are all the possible «relationships» between the 6 lords.
- Select one lord (one vertex).
- Suppose there are at least 3 lords **allied** with the selected one. Colour with **red** the edges linking these allies. So the selected vertex has at least 3 incident red edges.
- **Case 1: for some $i, j \in \{1, 2, 3\}$ A_i, A_j are allies.** Then we have (at least) 3 lords who are pairwise allied.
- **Case 2: all A_i, A_j are enemies.** Colour the edges between **enemies** with **blue**. Then $A_1A_2A_3$ are pairwise enemies.

Formalizing the problem in terms of graphs

Definition 1. *A k -edge colouring of a graph G is a labeling $f : E(G) \rightarrow [k]$*

The previous conjecture proves in fact the following result.

Theorem 1. *For every 2-colouring of K_6 , there is a monochromatic (with respect to edge colourings) subgraph of at least 3 vertices.*

Central question

how big a structure has to be, in order to contain *ordered substructures*?

Edge colourings

Given an integer $r \geq 0$, I ask if there exists an integer n such that K_n contains a monochromatic K_r , for any 2-edge colouring of K_n .

Cliques and independent sets

Given two integers $k, l \geq 0$, I ask if there exists an integer n such that every graph G on at least n vertices contains either K_k or \bar{K}_l .



Ramsey's answer

- Ramsey gave positive answer to this question, proving the following theorem, which here is stated in terms of cliques and independent sets.

Theorem 2. *Given any integers $k, l \geq 0$ there exists a smallest integer, denoted as $R(k, l)$, such that every graph on (at least) $R(k, l)$ vertices contains either a clique of k vertices or an independent set of l vertices.*

- The number $R(k, l)$ is the so called *Ramsey number*. If $k = l$, $R(k, k)$ is a *diagonal Ramsey number*.

Properties of Ramsey numbers

Proposition 1. *The following properties for Ramsey numbers hold:*

1. $R(1, l) = R(k, 1) = 1$, for every $k, l \geq 0$;

2. $R(2, l) = l, R(k, 2) = k$ for every $k, l \geq 0$;

3. $R(k, l) = R(l, k)$ for every $k, l \geq 0$;

4. for every $k, l \geq 2$

$$R(k, l) \leq R(k, l - 1) + R(k - 1, l);$$

strict inequality holds if $R(k, l - 1)$ and $R(k - 1, l)$ are both even;

5. for every $k, l \geq 2$

$$R(k, l) \leq \binom{k+l-2}{k-1}.$$

Proof. (1), (2), (3) can be easily verified.

(4) Let G be a graph on $R(k, l - 1) + R(k - 1, l)$ vertices. Let $v \in V(G)$.

The number of vertices adjacent to v , plus the number of vertices nonadjacent to v is $R(k, l - 1) + R(k - 1, l) - 1$.

Case 1 There exists a set of vertices S of at least $R(k, l - 1)$ elements, to which v is nonadjacent;

Case 2 there exists a set of vertices T of at least $R(k - 1, l)$ elements, to which v is adjacent.

If **case 1** holds, $G[S]$ contains either a k -clique or an $l - 1$ -independent set.

Then $G[S \cup v]$ contains either a k -clique or an l -independent set.

If **case 2** holds, $G[T]$ contains either a $k - 1$ -clique or an l -independent set.

Then $G[T \cup v]$ contains either a k -clique or an l -independent set.

In any case I can find either a k -clique or an l -independent set.

Then $R(k, l) \leq R(k, l - 1) + R(k - 1, l)$.

Now suppose $R(k, l - 1)$ and $R(k - 1, l)$ are both even.

Consider a graph G on $R(k, l - 1) + R(k - 1, l) - 1$ vertices.

The number of vertices is odd, so there exists a vertex v of even degree.

Observe that v cannot be adjacent to $R(k - 1, l) - 1$ vertices, or to $R(k, l - 1) - 1$ vertices, hence **case 1** or **case 2** must hold.

Repeating the argument used before, I can find a k -clique or an l -independent set in G , thus proving the claim.

Case 1 There exists a set of vertices S of at least $R(k, l - 1)$ elements, to which v is nonadjacent;

Case 2 there exists a set of vertices T of at least $R(k - 1, l)$ elements, to which v is adjacent.

(5) We prove the claim by induction on $k + l$.

Observe it holds if $k + l \leq 5$.

Suppose the claim is true for every k, l such that $5 \leq k + l < m + n$.

Let's show it holds for $m + n$.

Apply (4) to $R(m, n)$, then apply the inductive hypothesis and compute.

$$R(m, n) \leq R(m, n - 1) + R(m - 1, n) \leq \binom{m+n-3}{m-1} + \binom{m+n-3}{m-2} = \binom{m+n-2}{m-1}$$

□

Corollary 1. For all $k, l \geq 0$, $R(k, l) \leq 2^{k+l-2}$.

Ramsey numbers are in general difficult to calculate. Only few of them have been found, with the help of the previous properties and of some special graphs, called *Ramsey graphs*.

k	3	3	3	3	3	3	3	4	4
ℓ	3	4	5	6	7	8	9	4	5
$r(k, \ell)$	6	9	14	18	23	28	36	18	25

Ramsey numbers known to date (Bondy J.A., Murty U.S.R, 2008)


Definition 2. A (k, l) -Ramsey graph is a graph on $R(k, l) - 1$ vertices, containing no k -cliques and no l -independent sets.

Example: proof of $R(3,3)$ with the help of the (3,3)-Ramsey graph


- The graph contains no 3-cliques and no 3-independent sets.

So $R(3,3) \geq 6$.


- By *property (4)* we have $R(3,3) \leq R(2,3) + R(3,2) = 3 + 3 = 6$.
- Conclusion: $R(3,3) = 6$.




(3,3)-Ramsey graph. (Bondy J.A., Murty U.S.R., 2008)



A (3,4)-Ramsey graph



A (3,5)-Ramsey graph



A (4,4)-Ramsey graph

Ramsey's result for diagonal Ramsey numbers

Theorem 3 (Ramsey, 1930). *For every integer $r \geq 0$ there exists an integer $n \geq 0$ such that every graph on at least n vertices contains either K_r or \bar{K}_r .*



Proof. Suppose $r \geq 1$, otherwise the assertion is trivial.

Define $n = 2^{2r-3}$ and consider a graph G with $|V(G)| \geq n$.

We construct a sequence of subsets of $V(G)$, say $V_1, V_2, \dots, V_{2r-2}$, and choose $v_1 \in V_1, \dots, v_{2r-2} \in V_{2r-2}$ satisfying the following properties:

- (i) $|V_i| = 2^{2r-2-i}$, for $1 \leq i \leq 2r - 2$;
- (ii) $V_i \subseteq V_{i-1} \setminus \{v_{i-1}\}$, for $2 \leq i \leq 2r - 2$;
- (iii) v_{i-1} is adjacent either to all the vertices of V_i , or to no vertex of V_i , for $2 \leq i \leq 2r - 2$.

Construction of the sequence with the required properties.
(Diestel, 1997)



- (i) $|V_i| = 2^{2r-2-i}$, for $1 \leq i \leq 2r - 2$;
- (ii) $V_i \subseteq V_{i-1} \setminus \{v_{i-1}\}$, for $2 \leq i \leq 2r - 2$;
- (iii) v_{i-1} is adjacent either to all the vertices of V_i , or to no vertex of V_i , for $2 \leq i \leq 2r - 2$.

Such a sequence can be constructed inductively.

Start by choosing $V_1 \subseteq V(G)$, $|V_1| = n$. Pick v_1 arbitrarily.

Suppose to have constructed V_1, \dots, V_{i-1} for some i . Let's show I can define V_i .

Choose $v_{i-1} \in V_{i-1}$;

by (i), $|V_{i-1} - \{v_{i-1}\}| = 2^{2r-2-(i-1)} - 1 = 2^{2r-i-1} - 1$.

$|V_{i-1} - \{v_{i-1}\}|$ is an odd number, then:

either v_{i-1} is adjacent to more than a half of the vertices in V_{i-1} ,

or v_{i-1} is adjacent to less than a half of the vertices in V_{i-1} .

In both cases I can select $V_i \subseteq V_{i-1} - \{v_{i-1}\}$, satisfying (i), (ii), (iii).

We construct a sequence of subsets of $V(G)$, say $V_1, V_2, \dots, V_{2r-2}$, and choose $v_1 \in V_1, \dots, v_{2r-2} \in V_{2r-2}$ satisfying the following properties:

(i) $|V_i| = 2^{2r-2-i}$, for $1 \leq i \leq 2r - 2$;

(ii) $V_i \subseteq V_{i-1} \setminus \{v_{i-1}\}$, for $2 \leq i \leq 2r - 2$;

(iii) v_{i-1} is adjacent either to all the vertices of V_i , or to no vertex of V_i , for $2 \leq i \leq 2r - 2$.

(iii) v_{i-1} is adjacent either to all the vertices of V_i , or to no vertex of V_i , for $2 \leq i \leq 2r - 2$.

Pick any v_i . Observe that $2r - 2$ is even.

Among v_1, \dots, v_{2r-3} there is a set of $r - 1$ vertices with same behaviour in the sense of (iii).

Together with v_{2r-2} they induce either a K_r or a \bar{K}_r .

To understand this, observe that for every i , $v_i, v_{i+1}, \dots, v_{2r-2} \in V_i$.

□

Observe that the theorem gives an upper bound for diagonal Ramsey numbers, i. e.

$$R(r, r) \leq 2^{2r-3}$$

A lower bound for Ramsey numbers

Theorem 4 (Erdős, 1947). *For all $k \geq 2$*

$$R(k, k) \geq 2^{\frac{k}{2}}.$$

Proof. The case $k = 2$ is trivial.

If $k = 3$, $6 = R(3, 3) \geq 2^{\frac{3}{2}}$.

Suppose $k \geq 4$.

We will show with a probabilistic method that there exists a simple graph on $2^{\frac{k}{2}}$ vertices with no k -cliques and no k -independent sets.

Claim: \mathbb{P} (a simple graph on n vertices contains at least a K_k or a \bar{K}_k) < 1 .

Let $n \leq 2^{\frac{k}{2}}$.

In a simple graph we have that \mathbb{P} (a selected set of k elements forms a K_k or a \bar{K}_k) $= \frac{2}{2^{\binom{k}{2}}}$.

Hence \mathbb{P} (a simple graph on n vertices contains at least a K_k or a \bar{K}_k) $\leq \binom{n}{k} \frac{2}{2^{\binom{k}{2}}}$.

Now, $\binom{n}{k} \frac{2}{2^{\binom{k}{2}}} = \binom{n}{k} 2^{1-\binom{k}{2}} = 2^{1-\binom{k}{2}} \frac{n!}{k!(n-k)!} < \frac{2n^k}{k!2^{\binom{k}{2}}}$.

Computing $\binom{k}{2}$, and remembering that $n \leq 2^{\frac{k}{2}}$ we obtain

$\frac{2n^k}{k!2^{\binom{k}{2}}} \leq \frac{2^{\frac{k}{2}+1}}{k!} < 1$ (for $k! > 2^k$ for $k \geq 4$). This proves our **claim**.

Then, there must exist a graph on at most $2^{\frac{k}{2}}$ vertices with no k -cliques and no k -independent sets.

So $R(k, k) > 2^{\frac{k}{2}}$.

□

Conclusions

«Yet workers have only begun to explore the implications of the theory. It suggests that much of the essential structure of mathematics consists of extremely large numbers and sets, objects so huge that they are difficult to express, much less understand.

As we learn to handle these large numbers, we may find mathematical relations that help engineers to build large communications networks or help scientists to recognize patterns in large-scale physical systems.»

(1990, Graham R. L., Spencer J. H.)

Bibliography

Barton, Lane. *Ramsey Theory*. 13 May, 2016.

<https://www.Whitman.edu/Documents/Academics/Mathematics/2016/Barton.pdf>.

Bondy, Adrian, Murty M. Ram. *Graph Theory*, volume 244 of *Graduate Texts in Mathematics*. Springer-Verlag, London, 2008.

Diestel, Reinhard. *Graph Theory*, volume 173 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1997.

Gasarch, William. "Ramsey Theory and the History of Pre-Christian England: An Example

of Interdisciplinary Research." *Journal of Ramsey Theory* (2015): Web. 16 Feb. 2016. <http://www.cs.umd.edu/gasarch/BLOGPAPERS/ramseykings.pdf>.

Graham, Ronald L., and Joel H. Spencer. "Ramsey Theory." *Scientific American*. N.p., July 1990. Web. 8 Feb. 2016.