

START


RECORDING

Sets & Quantifiers

CMSC250


What is a set?

- A set is a collection of **distinct** objects.
- We use the notation $x \in S$ to say that S contains x.
- We'd like to know if $x \in S$ fast!
- Unless explicitly specified otherwise, **sets are unordered**.



What is a set?

- A set is a collection of **distinct** objects.
- We use the notation $x \in S$ to say that S contains x .
- We'd like to know if $x \in S$ fast!
- Unless explicitly specified otherwise, **sets are unordered**.
- Given the last two requirements, what's the **best possible data structure to implement a set in memory?**



Doubly Linked List


Binary Tree

Stack

Something else
(what?)

What is a set?

- A set is a collection of **distinct** objects.
- We use the notation $x \in S$ to say that S contains x.
- We'd like to know if $x \in S$ fast!
- Unless explicitly specified otherwise, **sets are unordered**.
- Given the last two requirements, what's the **best possible data structure to implement a set in memory?**



Doubly Linked List

Binary Tree

Stack

Something else
(what?)

Hash table!

Elementary number sets

- \mathbb{N} : the **natural** numbers
 - $\mathbb{N} = \{0, 1, 2, 3, \dots\}$. In our class, $0 \in \mathbb{N}$!
- \mathbb{Z} : the **integers**
 - $\mathbb{Z} = \{\dots - 3, -2, -1, 0, 1, 2, 3, \dots\}$
- \mathbb{Q} : the **rationals**
 - $\mathbb{Q} = \left\{ \frac{a}{b}, (a \in \mathbb{Z}) \wedge (b \in \mathbb{Z}) \wedge (b \neq 0) \right\}$
 - **Any** number that can be written as a **ratio of integers**!
- \mathbb{R} : the **reals**
 - This will typically be our “upper limit” in 250.
 - That is, we don’t usually care about \mathbb{C} , the set of **complex** numbers

Fill those in!

	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
0	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
-1	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$1/2$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$-1/2$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$0.333333\dots$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$0.333333\dots/0.11111111\dots$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
π	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i , such that $i^2 = -1$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Fill those in!

	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
0	■	■	■	■
-1	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$1/2$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$-1/2$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$0.333333\dots$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$0.333333\dots/0.11111111\dots$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
π	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i , such that $i^2 = -1$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Fill those in!

	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
0	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
-1	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
$1/2$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$-1/2$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$0.333333\dots$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$0.333333\dots/0.11111111\dots$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
π	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i , such that $i^2 = -1$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Fill those in!

	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
0	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
-1	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
$1/2$	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>
$-1/2$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$0.333333\dots$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
$0.333333\dots/0.11111111\dots$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
π	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
i , such that $i^2 = -1$	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Fill those in!

	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
0	■	■	■	■
-1	□	■	■	■
1/2	□	□	■	■
-1/2	□	□	■	■
0.333333...	□	□	□	□
0.333333.../0.11111111...	□	□	□	□
π	□	□	□	□
i , such that $i^2 = -1$	□	□	□	□

Fill those in!

	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
0	■	■	■	■
-1	□	■	■	■
$1/2$	□	□	■	■
$-1/2$	□	□	■	■
$0.333333\dots$	□	□	■	■
$0.333333\dots/0.11111111\dots$	□	□	□	□
π	□	□	□	□
i , such that $i^2 = -1$	□	□	□	□

Fill those in!

	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
0	■	■	■	■
-1	□	■	■	■
$1/2$	□	□	■	■
$-1/2$	□	□	■	■
$0.333333\dots$	□	□	■	■
$0.333333\dots/0.11111111\dots$	■	■	■	■
π	□	□	□	□
i , such that $i^2 = -1$	□	□	□	□

Fill those in!




	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
0	■	■	■	■
-1	□	■	■	■
$1/2$	□	□	■	■
$-1/2$	□	□	■	■
$0.333333\dots$	□	□	■	■
$0.333333\dots/0.11111111\dots$	■	■	■	■
π	□	□	□	■
i , such that $i^2 = -1$	□	□	□	□

Fill those in!




	\mathbb{N}	\mathbb{Z}	\mathbb{Q}	\mathbb{R}
0	■	■	■	■
-1	□	■	■	■
$1/2$	□	□	■	■
$-1/2$	□	□	■	■
$0.333333\dots$	□	□	■	■
$0.333333\dots/0.11111111\dots$	■	■	■	■
π	□	□	□	■
i , such that $i^2 = -1$	□	□	□	□

Not even
a real
number!

Venn Diagrams



Venn Diagrams



- U is the *Universal Domain*: a set that we imagine holds every *conceivable* element.
- When talking about sets of numbers, U is usually \mathbb{R} , the reals.

“There exists” (\exists)

- The symbol \exists (*LaTeX: `\exists`*) is read “There exists”.
- Examples:
 - $(\exists x \in \mathbb{R}) [8x = 1]$

“There exists” (\exists)

- The symbol \exists (*LaTeX: `\exists`*) is read “There exists”.
- Examples:
 - $(\exists x \in \mathbb{R}) [8x = 1]$ True

“There exists” (\exists)

- The symbol \exists (*LaTeX: \exists*) is read “There exists”.
- Examples:
 - $(\exists x \in \mathbb{R}) [8x = 1]$ True
 - $(\exists n \in \mathbb{Z}) [n^2 = -1]$

“There exists” (\exists)

- The symbol \exists (*LaTeX: \exists*) is read “There exists”.
- Examples:
 - $(\exists x \in \mathbb{R}) [8x = 1]$ True
 - $(\exists n \in \mathbb{Z}) [n^2 = -1]$ False

“There exists” (\exists)

- The symbol \exists (*LaTeX: \exists*) is read “There exists”.
- Examples:
 - $(\exists x \in \mathbb{R}) [8x = 1]$ True
 - $(\exists n \in \mathbb{Z}) [n^2 = -1]$ False
- Is there a domain D where $(\exists n \in D) [n^2 = -1]$ is true?

Yes

No

Something
else

“There exists” (\exists)

- The symbol \exists (*LaTeX: \exists*) is read “There exists”.
- Examples:
 - $(\exists x \in \mathbb{R}) [8x = 1]$ True
 - $(\exists n \in \mathbb{Z}) [n^2 = -1]$ False
- Is there a domain D where $(\exists n \in D) [n^2 = -1]$ is true?

The
complex
numbers \mathbb{C}

Yes

No

Something
else

“For all” (\forall)

- The symbol \forall (*LaTeX: `\forall`*) is read “for all”.
- Examples:
 - $(\forall x \in \mathbb{N}) [((x > 2) \wedge (x \text{ is prime})) \Rightarrow (x \text{ is odd})]$

“For all” (\forall)

- The symbol \forall (*LaTeX: `\forall`*) is read “for all”.
- Examples:
 - $(\forall x \in \mathbb{N}) [((x > 2) \wedge (x \text{ is prime})) \Rightarrow (x \text{ is odd})]$
True

“For all” (\forall)

- The symbol \forall is read “for all”.
- Examples:
 - $(\forall x \in \mathbb{N}) [((x > 2) \wedge (x \text{ is prime})) \Rightarrow (x \text{ is odd})]$
True
 - $(\forall n \in \mathbb{Z}) [n^2 \geq 0]$

“For all” (\forall)

- The symbol \forall is read “for all”.
- Examples:
 - $(\forall x \in \mathbb{N}) [((x > 2) \wedge (x \text{ is prime})) \Rightarrow (x \text{ is odd})]$
True
 - $(\forall n \in \mathbb{Z}) [n^2 \geq 0]$
True

“For all” (\forall)

- Let D be the set of all students in this class who are over 8 feet tall.
- $(\forall x \in D)[x \text{ has perfect attendance so far!}]$


True

False

Something
else

“For all” (\forall)

- Let D be the set of all students in this class who are over 8 feet tall.
- $(\forall x \in D)[x \text{ has perfect attendance so far!}]$



- If disagree, need to find $x \in D$ who missed a class
- Called **vacuously true!**

Nesting quantifiers

- $(\exists x \in \mathbb{N})(\exists y \in \mathbb{N})[x + 2y = 3x + y = 4]$

Nesting quantifiers

- $(\exists x \in \mathbb{N})(\exists y \in \mathbb{N})[x + 2y = 3x + y = 4]$ **False**

Nesting quantifiers

- $(\exists x \in \mathbb{N})(\exists y \in \mathbb{N})[x + 2y = 3x + y = 4]$ **False**
- $(\exists x \in \mathbb{Q})(\exists y \in \mathbb{Q})[x + 2y = 3x + y = 4]$

Nesting quantifiers

- $(\exists x \in \mathbb{N})(\exists y \in \mathbb{N})[x + 2y = 3x + y = 4]$ **False**
- $(\exists x \in \mathbb{Q})(\exists y \in \mathbb{Q})[x + 2y = 3x + y = 4]$

$$\text{True, } x = \frac{4}{5}, y = \frac{8}{5}$$

Nesting quantifiers

- $(\exists x \in \mathbb{N})(\exists y \in \mathbb{N})[x + 2y = 3x + y = 4]$ **False**
- $(\exists x \in \mathbb{Q})(\exists y \in \mathbb{Q})[x + 2y = 3x + y = 4]$

$$\text{True, } x = \frac{4}{5}, y = \frac{8}{5}$$

- Common abbreviation: $(\exists x, y \in D)[\dots]$
- Generally: $(\exists x_1, x_2, \dots, x_n \in D)[\dots]$

Alternating nested quantifiers

- Notice the differences between the following:
 - $(\forall x \in \mathbb{N})(\exists y \in \mathbb{N})[x < y]$
 - $(\exists x \in \mathbb{N})(\forall y \in \mathbb{N})[x < y]$

Alternating nested quantifiers

- Notice the differences between the following:
 - $(\forall x \in \mathbb{N})(\exists y \in \mathbb{N})[x < y]$ True (\mathbb{N} unbounded from above)
 - $(\exists x \in \mathbb{N})(\forall y \in \mathbb{N})[x < y]$

Alternating nested quantifiers

- Notice the differences between the following:
 - $(\forall x \in \mathbb{N})(\exists y \in \mathbb{N})[x < y]$ True (\mathbb{N} unbounded from above)
 - $(\exists x \in \mathbb{N})(\forall y \in \mathbb{N})[x < y]$ False (\mathbb{N} bounded from below)

Alternating nested quantifiers

- Notice the differences between the following:
 - $(\forall x \in \mathbb{N})(\exists y \in \mathbb{N})[x < y]$ True (\mathbb{N} unbounded from above)
 - $(\exists x \in \mathbb{N})(\forall y \in \mathbb{N})[x < y]$ False (\mathbb{N} bounded from below)
- ***WHEN QUANTIFIERS ARE DIFFERENT, THEIR ORDER MATTERS!!!!!!***

Fill this in!

Statement	True	False
$(\exists n \in \mathbb{N})[n + n = 0]$	<input type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{N})[n + n = 1]$	<input type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{Z})[n + n = 1]$	<input type="radio"/>	<input type="radio"/>
$(\exists x, y \in \mathbb{Z})[x + y = 1]$	<input type="radio"/>	<input type="radio"/>
$(\exists x \in \mathbb{R})[x(x + 1) = -1]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[((x < y < z) \Rightarrow (x^2 < y^2 < z^2))]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[((x < y < z) \Rightarrow (x^3 < y^3 < z^3))]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y \in \mathbb{N})(\exists z \in \mathbb{Q})[(x < y) \Rightarrow (x < z < y)]$	<input type="radio"/>	<input type="radio"/>

Fill this in!

Statement	True	False
$(\exists n \in \mathbb{N})[n + n = 0]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{N})[n + n = 1]$	<input type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{Z})[n + n = 1]$	<input type="radio"/>	<input type="radio"/>
$(\exists x, y \in \mathbb{Z})[x + y = 1]$	<input type="radio"/>	<input type="radio"/>
$(\exists x \in \mathbb{R})[x(x + 1) = -1]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[((x < y < z) \Rightarrow (x^2 < y^2 < z^2))]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[((x < y < z) \Rightarrow (x^3 < y^3 < z^3))]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y \in \mathbb{N})(\exists z \in \mathbb{Q})[(x < y) \Rightarrow (x < z < y)]$	<input type="radio"/>	<input type="radio"/>

$n = 0$

Fill this in!

Statement	True	False
$(\exists n \in \mathbb{N})[n + n = 0]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{N})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists n \in \mathbb{Z})[n + n = 1]$	<input type="radio"/>	<input type="radio"/>
$(\exists x, y \in \mathbb{Z})[x + y = 1]$	<input type="radio"/>	<input type="radio"/>
$(\exists x \in \mathbb{R})[x(x + 1) = -1]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^2 < y^2 < z^2)]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^3 < y^3 < z^3)]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y \in \mathbb{N})(\exists z \in \mathbb{Q})[(x < y) \Rightarrow (x < z < y)]$	<input type="radio"/>	<input type="radio"/>

$$n = 0$$

$$2n = 1 \Rightarrow n = \frac{1}{2} \notin \mathbb{N}$$

Fill this in!

Statement	True	False
$(\exists n \in \mathbb{N})[n + n = 0]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{N})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists n \in \mathbb{Z})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists x, y \in \mathbb{Z})[x + y = 1]$	<input type="radio"/>	<input type="radio"/>
$(\exists x \in \mathbb{R})[x(x + 1) = -1]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^2 < y^2 < z^2)]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^3 < y^3 < z^3)]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y \in \mathbb{N})(\exists z \in \mathbb{Q})[(x < y) \Rightarrow (x < z < y)]$	<input type="radio"/>	<input type="radio"/>

$n = 0$

$2n = 1 \Rightarrow n = \frac{1}{2} \notin \mathbb{N}$

Similarly, $\frac{1}{2} \notin \mathbb{Z}$

Fill this in!

Statement	True	False
$(\exists n \in \mathbb{N})[n + n = 0]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{N})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists n \in \mathbb{Z})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists x, y \in \mathbb{Z})[x + y = 1]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists x \in \mathbb{R})[x(x + 1) = -1]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^2 < y^2 < z^2)]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^3 < y^3 < z^3)]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y \in \mathbb{N})(\exists z \in \mathbb{Q})[(x < y) \Rightarrow (x < z < y)]$	<input type="radio"/>	<input type="radio"/>

$n = 0$

$2n = 1 \Rightarrow n = \frac{1}{2} \notin \mathbb{N}$

Similarly, $\frac{1}{2} \notin \mathbb{Z}$

$x = 0, y = 1$ or
 $x = -1, y = 2, \text{ or...}$

Fill this in!

Statement	True	False
$(\exists n \in \mathbb{N})[n + n = 0]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{N})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists n \in \mathbb{Z})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists x, y \in \mathbb{Z})[x + y = 1]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists x \in \mathbb{R})[x(x + 1) = -1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^2 < y^2 < z^2)]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^3 < y^3 < z^3)]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y \in \mathbb{N})(\exists z \in \mathbb{Q})[(x < y) \Rightarrow (x < z < y)]$	<input type="radio"/>	<input type="radio"/>

$n = 0$

$2n = 1 \Rightarrow n = \frac{1}{2} \notin \mathbb{N}$

Similarly, $\frac{1}{2} \notin \mathbb{Z}$

$x = 0, y = 1$ or
 $x = -1, y = 2, \text{ or...}$

$x^2 + x + 1 = 0$ has no
real solutions

Fill this in!

Statement	True	False
$(\exists n \in \mathbb{N})[n + n = 0]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{N})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists n \in \mathbb{Z})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists x, y \in \mathbb{Z})[x + y = 1]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists x \in \mathbb{R})[x(x + 1) = -1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^2 < y^2 < z^2)]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^3 < y^3 < z^3)]$	<input type="radio"/>	<input type="radio"/>
$(\forall x, y \in \mathbb{N})(\exists z \in \mathbb{Q})[(x < y) \Rightarrow (x < z < y)]$	<input type="radio"/>	<input type="radio"/>

$n = 0$

$2n = 1 \Rightarrow n = \frac{1}{2} \notin \mathbb{N}$

Similarly, $\frac{1}{2} \notin \mathbb{Z}$

$x = 0, y = 1$ or
 $x = -1, y = 2$, or...

$x^2 + x + 1 = 0$ has no
real solutions

Think of graph of $f(x) = x^2$

Fill this in!

Statement	True	False
$(\exists n \in \mathbb{N})[n + n = 0]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists n \in \mathbb{N})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists n \in \mathbb{Z})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\exists x, y \in \mathbb{Z})[x + y = 1]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\exists x \in \mathbb{R})[x(x + 1) = -1]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^2 < y^2 < z^2)]$	<input type="radio"/>	<input checked="" type="radio"/>
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^3 < y^3 < z^3)]$	<input checked="" type="radio"/>	<input type="radio"/>
$(\forall x, y \in \mathbb{N})(\exists z \in \mathbb{Q})[(x < y) \Rightarrow (x < z < y)]$	<input type="radio"/>	<input type="radio"/>

$n = 0$

$2n = 1 \Rightarrow n = \frac{1}{2} \notin \mathbb{N}$

Similarly, $\frac{1}{2} \notin \mathbb{Z}$

$x = 0, y = 1$ or
 $x = -1, y = 2$, or...

$x^2 + x + 1 = 0$ has no
real solutions

Think of graph of $f(x) = x^2$

Think of graph of $f(x) = x^3$

Fill this in!

Statement	True	False	
$(\exists n \in \mathbb{N})[n + n = 0]$	<input checked="" type="radio"/>	<input type="radio"/>	$n = 0$
$(\exists n \in \mathbb{N})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>	$2n = 1 \Rightarrow n = \frac{1}{2} \notin \mathbb{N}$
$(\exists n \in \mathbb{Z})[n + n = 1]$	<input type="radio"/>	<input checked="" type="radio"/>	Similarly, $\frac{1}{2} \notin \mathbb{Z}$
$(\exists x, y \in \mathbb{Z})[x + y = 1]$	<input checked="" type="radio"/>	<input type="radio"/>	$x = 0, y = 1$ or $x = -1, y = 2$, or...
$(\exists x \in \mathbb{R})[x(x + 1) = -1]$	<input type="radio"/>	<input checked="" type="radio"/>	$x^2 + x + 1 = 0$ has no real solutions
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^2 < y^2 < z^2)]$	<input type="radio"/>	<input checked="" type="radio"/>	Think of graph of $f(x) = x^2$
$(\forall x, y, z \in \mathbb{R})[(x < y < z) \Rightarrow (x^3 < y^3 < z^3)]$	<input checked="" type="radio"/>	<input type="radio"/>	Think of graph of $f(x) = x^3$
$(\forall x, y \in \mathbb{N})(\exists z \in \mathbb{Q})[(x < y) \Rightarrow (x < z < y)]$	<input checked="" type="radio"/>	<input type="radio"/>	E.g: arithmetic mean

Finding domains

- Give infinite sets D such that $(\forall x \in D)(\exists y \in D)[x < y]$
 1. Is **true**

Finding domains

- Give infinite sets D such that $(\forall x \in D)(\exists y \in D)[x < y]$
 1. Is true ($D = \mathbb{N}$, select $y = x + 1$)

Finding domains

- Give infinite sets D such that $(\forall x \in D)(\exists y \in D)[x < y]$
 1. Is true ($D = \mathbb{N}$, select $y = x + 1$)
 2. Is false

Finding domains

- Give infinite sets D such that $(\forall x \in D)(\exists y \in D)[x < y]$
 1. Is true ($D = \mathbb{N}$, select $y = x + 1$)
 2. Is false ($D = \mathbb{Z}^{\leq 0}$, counter-example is 0)

Finding domains

- Give infinite sets D such that $(\forall x \in D)(\exists y \in D)[x < y]$
 1. Is true ($D = \mathbb{N}$, select $y = x + 1$)
 2. Is false ($D = \mathbb{Z}^{\leq 0}$, counter-example is 0)
- Do the same thing for

$$(\forall x \in D)[x \leq 1] \wedge (\forall x \in D)(\exists y \in D)[x < y]$$

Finding domains

- Give infinite sets D such that $(\forall x \in D)(\exists y \in D)[x < y]$
 1. Is true ($D = \mathbb{N}$, select $y = x + 1$)
 2. Is false ($D = \mathbb{Z}^{\leq 0}$, counter-example is 0)
- Do the same thing for

$$(\forall x \in D)[x \leq 1] \wedge (\forall x \in D)(\exists y \in D)[x < y]$$

1. True for $D = (-\infty, 1)$

Finding domains

- Give infinite sets D such that $(\forall x \in D)(\exists y \in D)[x < y]$
 1. Is true ($D = \mathbb{N}$, select $y = x + 1$)
 2. Is false ($D = \mathbb{Z}^{\leq 0}$, counter-example is 0)
- Do the same thing for

$$(\forall x \in D)[x \leq 1] \wedge (\forall x \in D)(\exists y \in D)[x < y]$$

1. True for $D = (-\infty, 1)$
2. False for $D = (-\infty, 1]$ (!)


Subset

- We say that A is a subset of B ($A \subseteq B$) iff

$$(\forall x \in A)[x \in B]$$




$$(\forall x \in U)[(x \in A) \Rightarrow (x \in B)]$$




Superset and proper subset/superset

- We say that B is a **superset** of A ($B \supseteq A$) iff $A \subseteq B$.
- We say that A is a **proper subset** of B ($A \subset B$) iff $(A \subseteq B) \wedge (A \neq B) \wedge (A \neq \emptyset)$.
- We say that B is a **proper superset** of A ($B \supset A$) iff $A \subset B$



Union


$$A \cup B = \{(x \in A) \vee (x \in B)\}$$



Union


$$A \cup B = \{(x \in A) \vee (x \in B)\}$$

Connection between union
and logical disjunction!




Intersection

$$A \cap B = \{(x \in A) \wedge (x \in B)\}$$



Absolute complement


$$A^c = \{x \notin A\} = \{(x \in U) \wedge (\sim(x \in A))\}$$



Absolute complement

$$A^c = \{(x \notin A)\} = \{(x \in U) \wedge (\sim(x \in A))\}$$

Connection between
absolute complement and
logical negation!




Absolute complement

$$A^c = \{x \notin A\} = \{(x \in U) \wedge (\sim(x \in A))\}$$


Some use A' . They are Wrong, we are right.

Connection between
absolute complement and
logical negation!




Relative Complement

$$A - B = \{(x \in A) \wedge (x \notin B)\}$$




Relative Complement

$$A - B = \{(x \in A) \wedge (x \notin B)\}$$



Some use $A \setminus B$. They are wrong, we are right!



Careful about membership and subset!

- Be careful to distinguish between **members** of a set and **subsets** of a set...

True

False

Careful about membership and subset!

- Be careful to distinguish between **members** of a set and **subsets** of a set...

True

False

1. $1 \in \{-2, 0, 1, 3\}$

Careful about membership and subset!

- Be careful to distinguish between **members** of a set and **subsets** of a set...

True

False

1. $1 \in \{-2, 0, 1, 3\}$ T
2. $1 \in \{-2, 0, \{1\}, 3\}$

Careful about membership and subset!

- Be careful to distinguish between **members** of a set and **subsets** of a set...

True

False

1. $1 \in \{-2, 0, 1, 3\}$ T
2. $1 \in \{-2, 0, \{1\}, 3\}$ F
3. $1 \subseteq \{-2, 0, \{1\}, 3\}$

Careful about membership and subset!

- Be careful to distinguish between **members** of a set and **subsets** of a set...

True

False

1. $1 \in \{-2, 0, 1, 3\}$ T
2. $1 \in \{-2, 0, \{1\}, 3\}$ F
3. $1 \subseteq \{-2, 0, \{1\}, 3\}$ F, in fact, not even mathematically correct syntax
4. $\{1\} \subseteq \{-2, 0, \{1\}, 3\}$

Careful about membership and subset!

- Be careful to distinguish between **members** of a set and **subsets** of a set...

True

False

1. $1 \in \{-2, 0, 1, 3\}$ T
2. $1 \in \{-2, 0, \{1\}, 3\}$ F
3. $1 \subseteq \{-2, 0, \{1\}, 3\}$ F, in fact, not even mathematically correct syntax
4. $\{1\} \subseteq \{-2, 0, \{1\}, 3\}$ F
5. $\{1\} \in \{-2, 0, \{1\}, 3\}$

Careful about membership and subset!

- Be careful to distinguish between **members** of a set and **subsets** of a set...

True

False

1. $1 \in \{-2, 0, 1, 3\}$ T
2. $1 \in \{-2, 0, \{1\}, 3\}$ F
3. $1 \subseteq \{-2, 0, \{1\}, 3\}$ F, in fact, not even mathematically correct syntax
4. $\{1\} \subseteq \{-2, 0, \{1\}, 3\}$ F
5. $\{1\} \in \{-2, 0, \{1\}, 3\}$ T
6. $\{1\} \subseteq \{-2, 0, 1, 3\}$

Careful about membership and subset!

- Be careful to distinguish between **members** of a set and **subsets** of a set...

True

False

1. $1 \in \{-2, 0, 1, 3\}$ T
2. $1 \in \{-2, 0, \{1\}, 3\}$ F
3. $1 \subseteq \{-2, 0, \{1\}, 3\}$ F, in fact, not even mathematically correct syntax
4. $\{1\} \subseteq \{-2, 0, \{1\}, 3\}$ F
5. $\{1\} \in \{-2, 0, \{1\}, 3\}$ T
6. $\{1\} \subseteq \{-2, 0, 1, 3\}$ T

The empty set (\emptyset , $\{ \}$)

- The empty set, denoted either \emptyset or $\{ \}$, is the **unique** set with **no elements**.
 - Uniqueness can be proven, through a proof by contradiction!

The empty set (\emptyset , $\{ \}$)

- The empty set, denoted either \emptyset or $\{ \}$, is the **unique** set with **no elements**.
 - Uniqueness can be proven, through a proof by contradiction!

True

False

The empty set ($\emptyset, \{ \}$)

- The empty set, denoted either \emptyset or $\{ \}$, is the **unique** set with **no elements**.
 - Uniqueness can be proven, through a proof by contradiction!

True

False

1. $\emptyset \subseteq \mathbb{N}$

The empty set ($\emptyset, \{ \}$)

- The empty set, denoted either \emptyset or $\{ \}$, is the **unique** set with **no elements**.
 - Uniqueness can be proven, through a proof by contradiction!

True

False

1. $\emptyset \subseteq \mathbb{N}$ **T**
2. $\emptyset \subseteq A$ for **any set** A

The empty set ($\emptyset, \{ \}$)

- The empty set, denoted either \emptyset or $\{ \}$, is the **unique** set with **no elements**.
 - Uniqueness can be proven, through a proof by contradiction!

True

False

1. $\emptyset \subseteq \mathbb{N}$ T
2. $\emptyset \subseteq A$ for **any set** A T
3. $\emptyset \subset A$ for **any set** A

The empty set ($\emptyset, \{ \}$)

- The empty set, denoted either \emptyset or $\{ \}$, is the **unique** set with **no elements**.
 - Uniqueness can be proven, through a proof by contradiction!

True

False

1. $\emptyset \subseteq \mathbb{N}$ **T**
2. $\emptyset \subseteq A$ for **any set** A **T**
3. $\emptyset \subset A$ for **any set** A **F**
4. $\emptyset \subseteq \emptyset$

The empty set ($\emptyset, \{ \}$)

- The empty set, denoted either \emptyset or $\{ \}$, is the **unique** set with **no elements**.
 - Uniqueness can be proven, through a proof by contradiction!

True

False

1. $\emptyset \subseteq \mathbb{N}$ **T**
2. $\emptyset \subseteq A$ for **any set** A **T**
3. $\emptyset \subset A$ for **any set** A **F**
4. $\emptyset \subseteq \emptyset$ **T**

The powerset

- Given a set A , the powerset $\mathcal{P}(A)$ is **the set of all subsets of A .**
 - $\mathcal{P}(\{0, 1\}) = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$
 - $\mathcal{P}(\{0, 1, 2\}) = \{\emptyset, \{0\}, \{1\}, \{2\}, \{0, 1\}, \{1, 2\}, \{0, 2\}, \{0, 1, 2\}\}$
 - Evens, Odds, Primes, Squares
 - And lots more...

Facts about the powerset

- The following are **facts** about the powerset:
 - Since $\emptyset \subseteq A$ for all sets A , $\emptyset \in \mathcal{P}(A)$ for all sets A
 - Since $A \subseteq A$ for all sets A , $A \in \mathcal{P}(A)$ for all sets A

Powerset quizzing

- Let $A = \{1, 2, \dots, n\}$
- Then, $|P(A)|$

$$\approx n \cdot \log n$$

$$= n^2$$

$$= 2^n$$

$$= n!$$

Powerset quizzing

- Let $A = \{1, 2, \dots, n\}$
- Then, $|P(A)|$

$$\approx n \cdot \log n$$

$$= n^2$$

$$= 2^n$$

$$= n!$$

Powerset quizzing

- $P(\{1\}) =$

Powerset quizzing

- $P(\{1\}) = \{\emptyset, \{1\}\}$
- $P(P(\{1\})) =$

Powerset quizzing

- $P(\{1\}) = \{\emptyset, \{1\}\}$
- $P(P(\{1\})) = \{\emptyset, \{\emptyset\}, \{\{1\}\}, \{\emptyset, \{1\}\}\}$
- $P(\emptyset) =$

Powerset quizzing

- $P(\{1\}) = \{\emptyset, \{1\}\}$
- $P(P(\{1\})) = \{\emptyset, \{\emptyset\}, \{\{1\}\}, \{\emptyset, \{1\}\}\}$
- $P(\emptyset) = \{\emptyset\}$
- $P(\{\emptyset\}) =$

Powerset quizzing

- $P(\{1\}) = \{\emptyset, \{1\}\}$
- $P(P(\{1\})) = \{\emptyset, \{\emptyset\}, \{\{1\}\}, \{\emptyset, \{1\}\}\}$
- $P(\emptyset) = \{\emptyset\}$
- $P(\{\emptyset\}) = \{\emptyset, \{\emptyset\}\}$

STOP

RECORDING