

HW 11 CMSC 452
Morally Due TUES April 22 11:00AM
Dead-Cat Due THU April 24 11:00AM

1. (40 points) In class we did an example of taking a formula ϕ and producing a graph G and a number k such that

ϕ is satisfiable IFF G has an ind. set of size k .

The formula was in 3-CNF form which means it is of the form

$C_1 \wedge \cdots \wedge C_k$ where each C_i is the \vee of 3 literals.

The reduction produced k Δ s and then edges between the Δ s.

- (a) (10 points) Write psuedocode for a program that takes any formula in 3-CNF form and outputs a graph G and a number k .

Here is a HINT in that its the program with FILL IN THE BLANKS.

Begin with

- i. Input $C_1 \wedge \cdots \vee C_k$ where

$$C_1 = (L_{11} \vee L_{12} \vee L_{13})$$

$$C_2 = (L_{21} \vee L_{22} \vee L_{23})$$

$$\vdots \quad \quad \quad \vdots$$

$$C_k = (L_{k1} \vee L_{k2} \vee L_{k3})$$

- ii. V is YOU NEED TO DESCRIBE V .

- iii. E is YOU NEED TO DESCRIBE E .

- iv. Output (V, E) .

You do not need to prove or even mention that the program works in polynomial time. It will.

- (b) Describe the graph you get on input

$$(\neg x \vee y \vee z) \wedge (x \vee \neg y \vee z) \wedge (w \vee \neg x \vee y) \wedge (\neg w \vee x \vee \neg y).$$

- (c) (0 points) Think about. Come up with a 3-CNF formula that is NOT in 3-SAT. Apply your algorithm to it. What does the graph look like?

2. (30 points) In this problem all numbers are written in binary. Hence the number x takes $\lg_2(x)$ bits to represent and hence is of LENGTH $\lg_2(x)$.

In this problem all of the quantifiers range over $\{0, 1, 2, \dots\}$.

For $k \geq 2$ let

$$SQ_k = \{x : (\exists y_1, \dots, y_k)[x = y_1^2 + \dots + y_k^2]\}.$$

- (a) (6 points)

Show that, for all k , SQ_k is in NP.

(Just describe the witness y and the set B .)

- (b) (6 points) Look on the web to find out what is known about the following questions: (Here and for later questions, you don't have to look on the web if you are sure you know the answer.)

Is $SQ_2 \in P$? Is SQ_2 NP-Complete?

- (c) (6 points) Look on the web to find out what is known about the following questions:

Is $SQ_3 \in P$? Is SQ_3 NP-Complete?

- (d) (6 points) Look on the web to find out what is known about the following questions:

Is $SQ_4 \in P$? Is SQ_4 NP-Complete?

- (e) (6 points) Look on the web to find out what is known about the following questions:

Is $SQ_5 \in P$? Is SQ_5 NP-Complete?

3. (30 points)

A *Graph* is a $G = (V, E)$ where V is a set and E is a set of unordered pairs of elements of V . (Note that we do not allow self-loops and the edges are undirected.)

A graph $G = (V, E)$ is *k-colorable* if there is a function

$$f : V \rightarrow \{1, \dots, k\}$$

such that if $(x, y) \in E$ then $f(x) \neq f(y)$. (So two neighbors cannot have the same color.)

A graph is *Planar* if it can be drawn in the plane without crossing.

You can assume the following is true (it is!): $\{G : G \text{ is Planar}\} \in \text{P}$.

For all $k \geq \mathbb{N}$ let

$$A_k = \{G : G \text{ is Planar and } G \text{ is } k\text{-colorable}\}.$$

- (a) (6 points) Show that, for all k , the set A_k is in NP.
(Just describe the witness y and the set B .)
- (b) (6 points) Look on the web to find out what is known about the following questions: (Here and for later questions, you don't have to look on the web if you are sure you know the answer.)
Is $A_2 \in \text{P}$? Is A_2 NP-complete?
- (c) (6 points) Look on the web to find out what is known about the following questions:
Is $A_3 \in \text{P}$? Is A_3 NP-complete?
- (d) (6 points) Look on the web to find out what is known about the following questions:
Is $A_4 \in \text{P}$? Is A_4 NP-complete?
- (e) (6 points) Look on the web to find out what is known about the following questions: Is $A_5 \in \text{P}$? Is A_5 NP-complete?