

# BILL, RECORD LECTURE!!!!

BILL RECORD LECTURE!!!

# The Square Theorem

**Exposition by William Gasarch**

January 23, 2025

# The Square Theorem

**Definition** Let  $G \in \mathbb{N}$  and  $c \in \mathbb{N}$ . Let  $\text{COL}: [G] \times [G] \rightarrow [c]$ .

1. A **mono L** is 3 points

$$(x, y), (x + d, y), (x, y + d)$$

that are all the same color ( $d \geq 1$ ). (This should be called an *mono isosceles right triangle* but we just call it a *mono L*.)

2. A **mono Square** is 4 points

$$(x, y), (x + d, y), (x, y + d), (x + d, y + d)$$

that are all the same color ( $d \geq 1$ ). This is a square.

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

1. The proof of **The Square Theorem** gives enormous bounds on  $G$ ; however, the answer is known to be 15.

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

1. The proof of **The Square Theorem** gives enormous bounds on  $G$ ; however, the answer is known to be 15.
2. We will first prove *For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .*

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

1. The proof of **The Square Theorem** gives enormous bounds on  $G$ ; however, the answer is known to be 15.
2. We will first prove *For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .*
3. To prove **The Square Theorem** (about 2-coloring) we need to know that  $GG(c)$  exists for a very large  $c$ .

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

1. The proof of **The Square Theorem** gives enormous bounds on  $G$ ; however, the answer is known to be 15.
2. We will first prove *For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .*
3. To prove **The Square Theorem** (about 2-coloring) we need to know that  $GG(c)$  exists for a very large  $c$ .
4. More Colors: *For all  $c$  there exists  $G = G(c)$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [c]$  there exists a mono square. Proof needs a larger  $c'$  for  $GG(c')$ .*



# The $L$ Theorem for $c = 2$

**Theorem** For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .

# The $L$ Theorem for $c = 2$

**Theorem** For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .

**Proof** We prove this for  $c = 2$ . We will set  $H$  later. Let  $\text{COL}: [H] \times [H] \rightarrow [c]$ .

# The $L$ Theorem for $c = 2$

**Theorem** For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .

**Proof** We prove this for  $c = 2$ . We will set  $H$  later. Let  $\text{COL}: [H] \times [H] \rightarrow [c]$ .

Take the  $[H] \times [H]$  grid and tile it with  $3 \times 3$  tiles.

*View a 2-coloring of  $[H] \times [H]$  as a  $2^9$ -coloring of the tiles.*

# The $L$ Theorem for $c = 2$

**Theorem** For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .

**Proof** We prove this for  $c = 2$ . We will set  $H$  later. Let  $\text{COL}: [H] \times [H] \rightarrow [c]$ .

Take the  $[H] \times [H]$  grid and tile it with  $3 \times 3$  tiles.

*View a 2-coloring of  $[H] \times [H]$  as a  $2^9$ -coloring of the tiles.*

**This is very typical of VDW-Ramsey Theory: a 2-coloring of BLAH is viewed as a  $X$ -coloring of a different object where  $X$  is quite large.**

# Why This Size Tile?

Any 2-coloring of the  $3 \times 3$  tile will have two of the same color in the first column and hence an **almost  $L$**

# Why This Size Tile?

Any 2-coloring of the  $3 \times 3$  tile will have two of the same color in the first column and hence an **almost  $L$**

Goto White Board.

# Make $H$ Big Enough To Get Two Tiles Same Color

Take  $H = 3(2^9 + 1)$ .

# Make $H$ Big Enough To Get Two Tiles Same Color

Take  $H = 3(2^9 + 1)$ .

View  $[H] \times [H]$  grid of **points** as  $[2^9 + 1] \times [2^9 + 1]$  grid of **tiles**.



# Make $H$ Big Enough To Get Two Tiles Same Color

Take  $H = 3(2^9 + 1)$ .

View  $[H] \times [H]$  grid of **points** as  $[2^9 + 1] \times [2^9 + 1]$  grid of **tiles**.

Look at the first column of tiles. Two are the same color.

# Make $H$ Big Enough To Get Two Tiles Same Color

Take  $H = 3(2^9 + 1)$ .

View  $[H] \times [H]$  grid of **points** as  $[2^9 + 1] \times [2^9 + 1]$  grid of **tiles**.

Look at the first column of tiles. Two are the same color.

Go to White Board.

# The $L$ Theorem for $c = 3$

First take  $4 \times 4$ -tiles.

# The $L$ Theorem for $c = 3$

First take  $4 \times 4$ -tiles.

Any 3-coloring of the  $4 \times 4$  tile will have two of the same color in the first column and hence an **almost  $L$**

# The $L$ Theorem for $c = 3$

First take  $4 \times 4$ -tiles.

Any 3-coloring of the  $4 \times 4$  tile will have two of the same color in the first column and hence an **almost  $L$**

Goto White Board.

# Make $H$ Big Enough To Get Two Tiles Same Color

Take  $H = 4(3^{16} + 1)$ .

# Make $H$ Big Enough To Get Two Tiles Same Color

Take  $H = 4(3^{16} + 1)$ .

View  $[H] \times [H]$  grid of **points** as  $[3^{16} + 1] \times [3^{16} + 1]$  grid of **tiles**.

# Make $H$ Big Enough To Get Two Tiles Same Color

Take  $H = 4(3^{16} + 1)$ .

View  $[H] \times [H]$  grid of **points** as  $[3^{16} + 1] \times [3^{16} + 1]$  grid of **tiles**.

Look at the first column of tiles. Two are the same color.



# Make $H$ Big Enough To Get Two Tiles Same Color

Take  $H = 4(3^{16} + 1)$ .

View  $[H] \times [H]$  grid of **points** as  $[3^{16} + 1] \times [3^{16} + 1]$  grid of **tiles**.

Look at the first column of tiles. Two are the same color.

Go to White Board.

# Take Much Bigger Tiles

Take Tile so big that any 3-coloring of it has two different colored almost-L's converging to the same point.

# Take Much Bigger Tiles

Take Tile so big that any 3-coloring of it has two different colored almost-L's converging to the same point.

Go to White Board.

# Full $L$ Theorem

**Theorem** For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .

# Full $L$ Theorem

**Theorem** For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .

- ▶ We won't prove this but I am sure any of you could prove it given what we have done so far. Would be messy.

# Full $L$ Theorem

**Theorem** For all  $c$  there exists  $GG = GG(c)$  such that for all  $\text{COL}: [GG] \times [GG] \rightarrow [c]$  there exists a mono  $L$ .

- ▶ We won't prove this but I am sure any of you could prove it given what we have done so far. Would be messy.
- ▶ Easier to prove it from the Hales-Jewitt Theorem, which we won't be doing.

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

**Proof**  $G$  will be  $GG(2)GG(2^{GG(2)^2})$ .



# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

**Proof**  $G$  will be  $GG(2)GG(2^{GG(2)^2})$ .

Tile the  $[G] \times [G]$  plane with  $GG(2) \times GG(2)$  Tiles.

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

**Proof**  $G$  will be  $GG(2)GG(2^{GG(2)^2})$ .

Tile the  $[G] \times [G]$  plane with  $GG(2) \times GG(2)$  Tiles.

*View the 2-coloring of  $[G] \times [G]$  as a  $2^{GG(2)^2}$ -coloring of the tiles.*

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

**Proof**  $G$  will be  $GG(2)GG(2^{GG(2)^2})$ .

Tile the  $[G] \times [G]$  plane with  $GG(2) \times GG(2)$  Tiles.

*View the 2-coloring of  $[G] \times [G]$  as a  $2^{GG(2)^2}$ -coloring of the tiles.*

For any 2-coloring of  $[G] \times [G]$ :

- ▶ Every tile has a mono  $L$
- ▶ There is a mono  $L$  of tiles.

# The Square Theorem

**Theorem** There exists  $G$  such that for all  $\text{COL}: [G] \times [G] \rightarrow [2]$  there exists a mono square.

**Proof**  $G$  will be  $GG(2)GG(2^{GG(2)^2})$ .

Tile the  $[G] \times [G]$  plane with  $GG(2) \times GG(2)$  Tiles.

*View the 2-coloring of  $[G] \times [G]$  as a  $2^{GG(2)^2}$ -coloring of the tiles.*

For any 2-coloring of  $[G] \times [G]$ :

- ▶ Every tile has a mono  $L$
- ▶ There is a mono  $L$  of tiles.

Go to Whiteboard for rest of proof.