

BILL, RECORD LECTURE!!!!

BILL RECORD LECTURE!!!

Probabilistic Method

Proof of Turan's Theorem

Exposition by William Gasarch

Turan's Theorem

Theorem If $G = (V, E)$ is a graph, $|V| = n$, and $|E| = e$, then G has an ind set of size at least

$$\frac{n}{\frac{2e}{n} + 1}.$$

Turan's Theorem

Theorem If $G = (V, E)$ is a graph, $|V| = n$, and $|E| = e$, then G has an ind set of size at least

$$\frac{n}{\frac{2e}{n} + 1}.$$

Turan proved this in 1941 with a complicated proof.

Turan's Theorem

Theorem If $G = (V, E)$ is a graph, $|V| = n$, and $|E| = e$, then G has an ind set of size at least

$$\frac{n}{\frac{2e}{n} + 1}.$$

Turan proved this in 1941 with a complicated proof. We proof this more easily using Probability, but first need a lemma.

Turan's Theorem

Theorem If $G = (V, E)$ is a graph, $|V| = n$, and $|E| = e$, then G has an ind set of size at least

$$\frac{n}{\frac{2e}{n} + 1}.$$

Turan proved this in 1941 with a complicated proof. We proof this

more easily using Probability, but first need a lemma. The proof

we give is due to Ravi Boppana and appears in the Alon-Spencer book on *The Probabilistic Method*

Lemma

Lemma If $G = (V, E)$ is a graph. Then

$$\sum_{v \in V} \deg(v) = 2e.$$

Lemma

Lemma If $G = (V, E)$ is a graph. Then

$$\sum_{v \in V} \deg(v) = 2e.$$

Proof: Try to count the edges by summing the degrees at each vertex. This counts every edge TWICE.

Proof of Turan's Theorem

Theorem If $G = (V, E)$ is a graph, $|V| = n$, and $|E| = e$, then G has an ind set of size

$$\geq \frac{n}{\frac{2e}{n} + 1}.$$

Proof of Turan's Theorem

Theorem If $G = (V, E)$ is a graph, $|V| = n$, and $|E| = e$, then G has an ind set of size

$$\geq \frac{n}{\frac{2e}{n} + 1}.$$

Proof: Take the graph and RANDOMLY permute the vertices.


Proof of Turan's Theorem

Theorem If $G = (V, E)$ is a graph, $|V| = n$, and $|E| = e$, then G has an ind set of size

$$\geq \frac{n}{\frac{2e}{n} + 1}.$$

Proof: Take the graph and RANDOMLY permute the vertices.

Example:




Proof of Turan's Theorem

Theorem If $G = (V, E)$ is a graph, $|V| = n$, and $|E| = e$, then G has an ind set of size

$$\geq \frac{n}{\frac{2e}{n} + 1}.$$

Proof: Take the graph and RANDOMLY permute the vertices.

Example:



The set of vertices that have NO edges coming out on the right form an Ind Set. Call this set I .

How Big is I ?

How big is I

How Big is I ?

How big is I

WRONG QUESTION!

How Big is I ?


How big is I

WRONG QUESTION!

What is the EXPECTED VALUE of the size of I .
(NOTE- we permuted the vertices RANDOMLY)


What is Prob $v \in I$

Let $v \in V$. What is prob that $v \in I$



What is Prob $v \in I$


Let $v \in V$. What is prob that $v \in I$



v has degree d_v . How many ways can v and its vertices be laid out: $(d_v + 1)!$. In how many of them is v on the right? $d_v!$.

What is Prob $v \in I$

Let $v \in V$. What is prob that $v \in I$




v has degree d_v . How many ways can v and its vertices be laid out: $(d_v + 1)!$. In how many of them is v on the right? $d_v!$.

$$\Pr(v \in I) = \frac{d_v!}{(d_v + 1)!} = \frac{1}{d_v + 1}.$$

What is Prob $v \in I$

Let $v \in V$. What is prob that $v \in I$




v has degree d_v . How many ways can v and its vertices be laid out: $(d_v + 1)!$. In how many of them is v on the right? $d_v!$.

$$\Pr(v \in I) = \frac{d_v!}{(d_v + 1)!} = \frac{1}{d_v + 1}.$$

Hence

What is Prob $v \in I$

Let $v \in V$. What is prob that $v \in I$



v has degree d_v . How many ways can v and its vertices be laid out: $(d_v + 1)!$. In how many of them is v on the right? $d_v!$.

$$\Pr(v \in I) = \frac{d_v!}{(d_v + 1)!} = \frac{1}{d_v + 1}.$$

Hence

$$E(|I|) = \sum_{v \in V} \frac{1}{d_v + 1}.$$

How Big is this Sum?

Need to find lower bound on

$$\sum_{v \in V} \frac{1}{d_v + 1}.$$

Rephrase

NEW PROBLEM:

Minimize

$$\sum_{v \in V} \frac{1}{x_v + 1}$$

relative to the constraint:

$$\sum_{v \in V} x_v = 2e.$$

KNOWN: This sum is minimized when all of the x_v are $\frac{2e}{|V|} = \frac{2e}{n}$.
So the min the sum can be is

$$\sum_{v \in V} \frac{1}{\frac{2e}{n} + 1} = \frac{n}{\frac{2e}{n} + 1}.$$

Recap and Done

$$E(|I|) = \sum_{v \in V} \frac{1}{d_v + 1} \text{ and } \sum_{v \in V} d_v = 2e.$$

Recap and Done

$$E(|I|) = \sum_{v \in V} \frac{1}{d_v + 1} \text{ and } \sum_{v \in V} d_v = 2e.$$

To lower bound $E(|I|)$ we solve a continuous problem: minimize $\sum_{v \in V} \frac{1}{x_v + 1}$ with constraint $\sum_{v \in V} x_v = 2e$.

Recap and Done

$$E(|I|) = \sum_{v \in V} \frac{1}{d_v + 1} \text{ and } \sum_{v \in V} d_v = 2e.$$

To lower bound $E(|I|)$ we solve a continuous problem: minimize $\sum_{v \in V} \frac{1}{x_v + 1}$ with constraint $\sum_{v \in V} x_v = 2e$.

The min occurs when $(\forall v)[x_v = \frac{2e}{n}]$. Hence

Recap and Done

$$E(|I|) = \sum_{v \in V} \frac{1}{d_v + 1} \text{ and } \sum_{v \in V} d_v = 2e.$$

To lower bound $E(|I|)$ we solve a continuous problem: minimize $\sum_{v \in V} \frac{1}{x_v + 1}$ with constraint $\sum_{v \in V} x_v = 2e$.

The min occurs when $(\forall v)[x_v = \frac{2e}{n}]$. Hence

$$E(I) \geq \sum_{v \in V} \frac{1}{x_v + 1} \geq \sum_{v \in V} \frac{1}{\frac{2e}{n} + 1} = \frac{n}{\frac{2e}{n} + 1}.$$