

The Muffin Problem

William Gasarch - University of MD

Erik Metz - University of MD

Jacob Prinz-University of MD

Daniel Smolyak- University of MD

How it Began


A Recreational Math Conference (Gathering for Gardner) May 2016

I found a pamphlet:

The Julia Robinson Mathematics Festival: A Sample of Mathematical Puzzles Compiled by Nancy Blachman

which had this problem, proposed by Alan Frank:






How can you divide and distribute 5 muffins to 3 students so that every student gets $\frac{5}{3}$ where nobody gets a tiny sliver?



5 Muffins, 3 Students, Proc by Picture

Person	Color	What they Get
Alice	RED	$1 + \frac{2}{3} = \frac{5}{3}$
Bob	BLUE	$1 + \frac{2}{3} = \frac{5}{3}$
Carol	GREEN	$1 + \frac{1}{3} + \frac{1}{3} = \frac{5}{3}$

Smallest Piece: $\frac{1}{3}$



Can We Do Better?

The smallest piece in the above solution is $\frac{1}{3}$.

Is there a procedure with a larger smallest piece?

Work on it with your neighbor


You have 7 minutes

5 Muffins, 3 People—Proc by Picture

YES WE CAN!

Person	Color	What they Get
Alice	RED	$\frac{6}{12} + \frac{7}{12} + \frac{7}{12}$
Bob	BLUE	$\frac{6}{12} + \frac{7}{12} + \frac{7}{12}$
Carol	GREEN	$\frac{5}{12} + \frac{5}{12} + \frac{5}{12} + \frac{5}{12}$

Smallest Piece: $\frac{5}{12}$



Can We Do Better?

The smallest piece in the above solution is $\frac{5}{12}$.

Is there a procedure with a larger smallest piece?

Work on it with your neighbor

You have 4 minutes

5 Muffins, 3 People—Can't Do Better Than $\frac{5}{12}$

NO WE CAN'T!

There is a procedure for 5 muffins, 3 students where each student gets $\frac{5}{3}$ muffins, smallest piece N . We want $N \leq \frac{5}{12}$.

Case 0: Some muffin is uncut. Cut it $(\frac{1}{2}, \frac{1}{2})$ and give both $\frac{1}{2}$ -sized pieces to whoever got the uncut muffin. (Note $\frac{1}{2} > \frac{5}{12}$.) Reduces to other cases. (**Henceforth:** All muffins cut into ≥ 2 pieces.)

5 Muffins, 3 People—Can't Do Better Than $\frac{5}{12}$

NO WE CAN'T!

There is a procedure for 5 muffins, 3 students where each student gets $\frac{5}{3}$ muffins, smallest piece N . We want $N \leq \frac{5}{12}$.

Case 0: Some muffin is uncut. Cut it $(\frac{1}{2}, \frac{1}{2})$ and give both $\frac{1}{2}$ -sized pieces to whoever got the uncut muffin. (Note $\frac{1}{2} > \frac{5}{12}$.) Reduces to other cases. (**Henceforth:** All muffins cut into ≥ 2 pieces.)

Case 1: Some muffin is cut into ≥ 3 pieces. Then $N \leq \frac{1}{3} < \frac{5}{12}$. (**Henceforth:** All muffins cut into 2 pieces.)

5 Muffins, 3 People—Can't Do Better Than $\frac{5}{12}$

NO WE CAN'T!

There is a procedure for 5 muffins, 3 students where each student gets $\frac{5}{3}$ muffins, smallest piece N . We want $N \leq \frac{5}{12}$.

Case 0: Some muffin is uncut. Cut it $(\frac{1}{2}, \frac{1}{2})$ and give both $\frac{1}{2}$ -sized pieces to whoever got the uncut muffin. (Note $\frac{1}{2} > \frac{5}{12}$.) Reduces to other cases. (**Henceforth:** All muffins cut into ≥ 2 pieces.)

Case 1: Some muffin is cut into ≥ 3 pieces. Then $N \leq \frac{1}{3} < \frac{5}{12}$. (**Henceforth:** All muffins cut into 2 pieces.)

Case 2: All muffins are cut into 2 pieces. 10 pieces, 3 students: **Someone** gets ≥ 4 pieces. He has some piece

$$\leq \frac{5}{3} \times \frac{1}{4} = \frac{5}{12} \quad \text{Great to see } \frac{5}{12}$$

What Else Was in the Pamphlet?

The pamphlet also had asked about

1. 4 muffins, 7 students.
2. 12 muffins, 11 students.
3. a few others

What Else Was in the Pamphlet?

The pamphlet also had asked about

1. 4 muffins, 7 students.
2. 12 muffins, 11 students.
3. a few others

This seemed like a nice exercise and it was.

What Else Was in the Pamphlet?

The pamphlet also had asked about

1. 4 muffins, 7 students.
2. 12 muffins, 11 students.
3. a few others

This seemed like a nice exercise and it was.

There can't be much more to this.

If there is not much more to this then how come

[https://www.amazon.com/
Mathematical-Muffin-Morsels-Problem-Mathematics/dp/
9811215170](https://www.amazon.com/Mathematical-Muffin-Morsels-Problem-Mathematics/dp/9811215170)

If there is not much more to this then how come

[https://www.amazon.com/
Mathematical-Muffin-Morsels-Problem-Mathematics/dp/
9811215170](https://www.amazon.com/Mathematical-Muffin-Morsels-Problem-Mathematics/dp/9811215170)

The following happened:

If there is not much more to this then how come

[https://www.amazon.com/
Mathematical-Muffin-Morsels-Problem-Mathematics/dp/
9811215170](https://www.amazon.com/Mathematical-Muffin-Morsels-Problem-Mathematics/dp/9811215170)

The following happened:

- Find a technique that solves many problems.

If there is not much more to this then how come

[https://www.amazon.com/
Mathematical-Muffin-Morsels-Problem-Mathematics/dp/
9811215170](https://www.amazon.com/Mathematical-Muffin-Morsels-Problem-Mathematics/dp/9811215170)

The following happened:

- ▶ Find a technique that solves many problems.
- ▶ Find least problem the set of techniques does not work for.

If there is not much more to this then how come

`https://www.amazon.com/
Mathematical-Muffin-Morsels-Problem-Mathematics/dp/
9811215170`

The following happened:

- ▶ Find a technique that solves many problems.
- ▶ Find least problem the set of techniques does not work for.
- ▶ Find a new technique **which is interesting**.

If there is not much more to this then how come

[https://www.amazon.com/
Mathematical-Muffin-Morsels-Problem-Mathematics/dp/
9811215170](https://www.amazon.com/Mathematical-Muffin-Morsels-Problem-Mathematics/dp/9811215170)

The following happened:

- ▶ Find a technique that solves many problems.
- ▶ Find least problem the set of techniques does not work for.
- ▶ Find a new technique **which is interesting**.
- ▶ Lather, Rinse, Repeat.

General Problem

$f(m, s)$ be the smallest piece in the best procedure (best in that the smallest piece is maximized) to divide m muffins among s students so that everyone gets $\frac{m}{s}$.

General Problem

$f(m, s)$ be the smallest piece in the best procedure (best in that the smallest piece is maximized) to divide m muffins among s students so that everyone gets $\frac{m}{s}$.

We just showed showed $f(5, 3) = \frac{5}{12}$.

General Problem

$f(m, s)$ be the smallest piece in the best procedure (best in that the smallest piece is maximized) to divide m muffins among s students so that everyone gets $\frac{m}{s}$.

We just showed showed $f(5, 3) = \frac{5}{12}$.

We showed $f(m, s)$ exists, rational, computable, via a Mixed Int Program.

Amazing Results!/Amazing Theorems!

1. $f(43, 33) = \frac{91}{264}$.
2. $f(52, 11) = \frac{83}{176}$.
3. $f(35, 13) = \frac{64}{143}$.

Amazing Results!/Amazing Theorems!

1. $f(43, 33) = \frac{91}{264}$.
2. $f(52, 11) = \frac{83}{176}$.
3. $f(35, 13) = \frac{64}{143}$.

All done by hand, no use of a computer

Amazing Results!/Amazing Theorems!

1. $f(43, 33) = \frac{91}{264}$.
2. $f(52, 11) = \frac{83}{176}$.
3. $f(35, 13) = \frac{64}{143}$.

All done by hand, no use of a computer
by Co-author Erik Metz is a muffin savant !

Amazing Results!/Amazing Theorems!

1. $f(43, 33) = \frac{91}{264}$.
2. $f(52, 11) = \frac{83}{176}$.
3. $f(35, 13) = \frac{64}{143}$.

All done by hand, no use of a computer
by Co-author Erik Metz is a **muffin savant** !

Have **General Theorems** from which **upper bounds** follow.

Amazing Results!/Amazing Theorems!

1. $f(43, 33) = \frac{91}{264}$.
2. $f(52, 11) = \frac{83}{176}$.
3. $f(35, 13) = \frac{64}{143}$.

All done by hand, no use of a computer
by Co-author Erik Metz is a **muffin savant** !

Have **General Theorems** from which **upper bounds** follow.
Have **General Procedures** from which **lower bounds** follow.

7 Muffins, 3 Students

Work on $f(7, 3)$ in groups.

7 Muffins, 3 Students.

Get upper and lower bounds that match!

You have 4 minutes

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\geq \lceil \frac{14}{3} \rceil = 5$ pieces.

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\geq \lceil \frac{14}{3} \rceil = 5$ pieces.
5. That student must get a piece $\leq \frac{7}{3} \times \frac{1}{5} = \frac{7}{15}$.

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\geq \lceil \frac{14}{3} \rceil = 5$ pieces.
5. That student must get a piece $\leq \frac{7}{3} \times \frac{1}{5} = \frac{7}{15}$.
6. Great! We know $f(7, 3) \leq \frac{7}{15}$.

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\geq \lceil \frac{14}{3} \rceil = 5$ pieces.
5. That student must get a piece $\leq \frac{7}{3} \times \frac{1}{5} = \frac{7}{15}$.
6. Great! We know $f(7, 3) \leq \frac{7}{15}$.
7. Can we show a protocol that gives $f(7, 3) \geq \frac{7}{15}$.

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\geq \lceil \frac{14}{3} \rceil = 5$ pieces.
5. That student must get a piece $\leq \frac{7}{3} \times \frac{1}{5} = \frac{7}{15}$.
6. Great! We know $f(7, 3) \leq \frac{7}{15}$.
7. Can we show a protocol that gives $f(7, 3) \geq \frac{7}{15}$.
8. We tried. We failed. Darn :-)

7 Muffins, 3 Students: How to think about it

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\geq \lceil \frac{14}{3} \rceil = 5$ pieces.
5. That student must get a piece $\leq \frac{7}{3} \times \frac{1}{5} = \frac{7}{15}$.
6. Great! We know $f(7, 3) \leq \frac{7}{15}$.
7. Can we show a protocol that gives $f(7, 3) \geq \frac{7}{15}$.
8. We tried. We failed. Darn :-)

Now what?

7 Muffins, 3 Students: How to think about it again

We first look at LIMITS on what we can expect.

7 Muffins, 3 Students: How to think about it again

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.

7 Muffins, 3 Students: How to think about it again

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.

7 Muffins, 3 Students: How to think about it again

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.

7 Muffins, 3 Students: How to think about it again

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\leq \lfloor \frac{14}{3} \rfloor = 4$ pieces.

7 Muffins, 3 Students: How to think about it again

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\leq \lfloor \frac{14}{3} \rfloor = 4$ pieces.
5. That student must get a piece $\geq \frac{7}{3} \times \frac{1}{4} = \frac{7}{12}$.

7 Muffins, 3 Students: How to think about it again

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\leq \lfloor \frac{14}{3} \rfloor = 4$ pieces.
5. That student must get a piece $\geq \frac{7}{3} \times \frac{1}{4} = \frac{7}{12}$.
6. That piece came from a muffin. Other piece is $\leq 1 - \frac{7}{12} = \frac{5}{12}$.

7 Muffins, 3 Students: How to think about it again

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\leq \lfloor \frac{14}{3} \rfloor = 4$ pieces.
5. That student must get a piece $\geq \frac{7}{3} \times \frac{1}{4} = \frac{7}{12}$.
6. That piece came from a muffin. Other piece is $\leq 1 - \frac{7}{12} = \frac{5}{12}$.
7. Great! We know $f(7, 3) \leq \frac{5}{12}$.

7 Muffins, 3 Students: How to think about it again

We first look at LIMITS on what we can expect.

1. If a muffin is uncut, can cut it in two.
2. If a muffin is cut in ≥ 3 pieces then some piece $\leq \frac{1}{3}$. Unlikely.
3. 7 muffins, each one cut in two 2 pieces, so 14 pieces total.
4. 3 students, so some student gets $\leq \lfloor \frac{14}{3} \rfloor = 4$ pieces.
5. That student must get a piece $\geq \frac{7}{3} \times \frac{1}{4} = \frac{7}{12}$.
6. That piece came from a muffin. Other piece is $\leq 1 - \frac{7}{12} = \frac{5}{12}$.
7. Great! We know $f(7, 3) \leq \frac{5}{12}$.
8. Can we show a protocol that gives $f(7, 3) \geq \frac{5}{12}$?

7 Muffins, 3 Students: How to think about protocol

Want $f(7, 3) \geq \frac{5}{12}$.

7 Muffins, 3 Students: How to think about protocol

Want $f(7, 3) \geq \frac{5}{12}$.

Will be cutting some muffins $(\frac{5}{12}, \frac{7}{12})$.

7 Muffins, 3 Students: How to think about protocol

Want $f(7, 3) \geq \frac{5}{12}$.

Will be cutting some muffins $(\frac{5}{12}, \frac{7}{12})$.

Can also cut some muffins $(\frac{6}{12}, \frac{6}{12})$.

With this, You have 4 minutes to find a protocol

7 Muffins, 3 Students: How to think about protocol

Need to know what combos of $\frac{5}{12}$, $\frac{6}{12}$, $\frac{7}{12}$ add to $\frac{7}{3} = \frac{28}{12}$.

7 Muffins, 3 Students: How to think about protocol

Need to know what combos of $\frac{5}{12}, \frac{6}{12}, \frac{7}{12}$ add to $\frac{7}{3} = \frac{28}{12}$.

Need to know what combos of 5, 6, 7 add to 28.

$$7 + 7 + 7 + 7 = 28$$

$$5 + 5 + 6 + 6 + 6 = 28$$

7 Muffins, 3 Students: How to think about protocol

Need to know what combos of $\frac{5}{12}, \frac{6}{12}, \frac{7}{12}$ add to $\frac{7}{3} = \frac{28}{12}$.

Need to know what combos of 5, 6, 7 add to 28.

$$7 + 7 + 7 + 7 = 28$$

$$5 + 5 + 6 + 6 + 6 = 28$$

1. Cut 4 muffins $(\frac{5}{12}, \frac{7}{12})$.

7 Muffins, 3 Students: How to think about protocol

Need to know what combos of $\frac{5}{12}, \frac{6}{12}, \frac{7}{12}$ add to $\frac{7}{3} = \frac{28}{12}$.

Need to know what combos of 5, 6, 7 add to 28.

$$7 + 7 + 7 + 7 = 28$$

$$5 + 5 + 6 + 6 + 6 = 28$$

1. Cut 4 muffins $(\frac{5}{12}, \frac{7}{12})$.
2. Cut 3 muffins $(\frac{6}{12}, \frac{6}{12})$.

7 Muffins, 3 Students: How to think about protocol

Need to know what combos of $\frac{5}{12}, \frac{6}{12}, \frac{7}{12}$ add to $\frac{7}{3} = \frac{28}{12}$.

Need to know what combos of 5, 6, 7 add to 28.

$$7 + 7 + 7 + 7 = 28$$

$$5 + 5 + 6 + 6 + 6 = 28$$

1. Cut 4 muffins $(\frac{5}{12}, \frac{7}{12})$.
2. Cut 3 muffins $(\frac{6}{12}, \frac{6}{12})$.
3. Give 1 student 4 pieces of size $\frac{7}{12}$.

7 Muffins, 3 Students: How to think about protocol

Need to know what combos of $\frac{5}{12}, \frac{6}{12}, \frac{7}{12}$ add to $\frac{7}{3} = \frac{28}{12}$.

Need to know what combos of 5, 6, 7 add to 28.

$$7 + 7 + 7 + 7 = 28$$

$$5 + 5 + 6 + 6 + 6 = 28$$

1. Cut 4 muffins $(\frac{5}{12}, \frac{7}{12})$.
2. Cut 3 muffins $(\frac{6}{12}, \frac{6}{12})$.
3. Give 1 student 4 pieces of size $\frac{7}{12}$.
4. Give 2 students 2 pieces of size $\frac{5}{12}$ and 3 pieces of size $\frac{6}{12}$.

Conventions

We know and use the following:

1. Known: $f(m, s) = \frac{m}{s} f(s, m)$. Hence we assume $m > s$.
2. If s divides m then $f(m, s) = 1$ so assume s does not divide m .
3. All muffins are cut in ≥ 2 pcs. Replace uncut muff with $2 \frac{1}{2}$'s
4. If assuming $f(m, s) > \alpha > \frac{1}{3}$, assume all muffin in ≤ 2 pcs.
5. $f(m, s) > \alpha > \frac{1}{3}$, so exactly 2 pcs, is common case.

We do not know this but still use: $f(m, s)$ only depends on $\frac{m}{s}$.

All of our techniques that hold for (m, s) hold for (Am, As) .

For particular numbers, we only look at m, s rel prime.

FC Thm Generalizes $f(5, 3) \leq \frac{5}{12}$

$$f(m, s) \leq \text{FC}(m, s) = \max \left\{ \frac{1}{3}, \min \left\{ \frac{m}{s \lceil 2m/s \rceil}, 1 - \frac{m}{s \lfloor 2m/s \rfloor} \right\} \right\}.$$

Case 0: Some muffin is uncut. Cut it $(\frac{1}{2}, \frac{1}{2})$ and give both halves to whoever got the uncut muffin, so reduces to other cases.

Case 1: Some muffin is cut into ≥ 3 pieces. Some piece $\leq \frac{1}{3}$.

Case 2: Every muffin is cut into 2 pieces, so $2m$ pieces.

Someone gets $\geq \lceil \frac{2m}{s} \rceil$ pieces. $\exists$ piece $\leq \frac{m}{s} \times \frac{1}{\lceil 2m/s \rceil} = \frac{m}{s \lceil 2m/s \rceil}$.

Someone gets $\leq \lfloor \frac{2m}{s} \rfloor$ pieces. $\exists$ piece $\geq \frac{m}{s} \frac{1}{\lfloor 2m/s \rfloor} = \frac{m}{s \lfloor 2m/s \rfloor}$.

The other piece from that muffin is of size $\leq 1 - \frac{m}{s \lfloor 2m/s \rfloor}$.

THREE Students

CLEVERNESS, COMP PROGS for the procedure.

FC Theorem for optimality.

$$f(1, 3) = \frac{1}{3}$$

$$f(3k, 3) = 1.$$

$$f(3k + 1, 3) = \frac{3k-1}{6k}, k \geq 1.$$

$$f(3k + 2, 3) = \frac{3k+2}{6k+6}.$$

Note: A Mod 3 Pattern.

Theorem: For all $m \geq 3$, $f(m, 3) = \text{FC}(m, 3)$.

FOUR Students

CLEVERNESS, COMP PROGS for procedures.

FC Theorem for optimality.

$$f(4k, 4) = 1 \text{ (easy)}$$

$$f(1, 4) = \frac{1}{4} \text{ (easy)}$$

$$f(4k + 1, 4) = \frac{4k-1}{8k}, k \geq 1.$$

$$f(4k + 2, 4) = \frac{1}{2}.$$

$$f(4k + 3, 4) = \frac{4k+1}{8k+4}.$$

Note: A Mod 4 Pattern.

Theorem: For all $m \geq 4$, $f(m, 4) = \text{FC}(m, 4)$.

FC-Conjecture: For all m, s with $m \geq s$, $f(m, s) = \text{FC}(m, s)$.

FIVE Students

CLEVERNESS, COMP PROGS for procedures.

FC Theorem for optimality.

For $k \geq 1$, $f(5k, 5) = 1$.

For $k = 1$ and $k \geq 3$, $f(5k + 1, 5) = \frac{5k+1}{10k+5}$. $f(11, 5)$?

For $k \geq 2$, $f(5k + 2, 5) = \frac{5k-2}{10k}$. $f(7, 5) = \text{FC}(7, 5) = \frac{1}{3}$

For $k \geq 1$, $f(5k + 3, 5) = \frac{5k+3}{10k+10}$

For $k \geq 1$, $f(5k + 4, 5) = \frac{5k+1}{10k+5}$

Note: A Mod 5 Pattern.

Theorem: For all $m \geq 5$ **except $m=11$** , $f(m, 5) = \text{FC}(m, 5)$.

What About FIVE students, ELEVEN muffins?

$$f(11, 5) \leq \max \left\{ \frac{1}{3}, \min \left\{ \frac{11}{5 \lceil 22/5 \rceil}, 1 - \frac{11}{5 \lfloor 22/5 \rfloor} \right\} \right\} = \frac{11}{25}.$$

What About FIVE students, ELEVEN muffins?

$$f(11, 5) \leq \max \left\{ \frac{1}{3}, \min \left\{ \frac{11}{5 \lceil 22/5 \rceil}, 1 - \frac{11}{5 \lfloor 22/5 \rfloor} \right\} \right\} = \frac{11}{25}.$$

We tried to find a protocol to divide 11 muffins for 5 people, each gets $\frac{11}{5}$, and smallest piece is size $\frac{11}{25} = 0.44$.

What About FIVE students, ELEVEN muffins?

$$f(11, 5) \leq \max \left\{ \frac{1}{3}, \min \left\{ \frac{11}{5 \lceil 22/5 \rceil}, 1 - \frac{11}{5 \lfloor 22/5 \rfloor} \right\} \right\} = \frac{11}{25}.$$

We tried to find a protocol to divide 11 muffins for 5 people, each gets $\frac{11}{5}$, and smallest piece is size $\frac{11}{25} = 0.44$.

We found a protocol with smallest piece $\frac{13}{30} = 0.4333\dots$

1. Divide 1 muffin $(\frac{15}{30}, \frac{15}{30})$.
2. Divide 2 muffins $(\frac{14}{30}, \frac{16}{30})$.
3. Divide 8 muffins $(\frac{13}{30}, \frac{17}{30})$.
4. Give 2 students $[\frac{13}{30}, \frac{13}{30}, \frac{13}{30}, \frac{13}{30}, \frac{14}{30}]$
5. Give 1 students $[\frac{16}{30}, \frac{16}{30}, \frac{17}{30}, \frac{17}{30}]$
6. Give 2 students $[\frac{15}{30}, \frac{17}{30}, \frac{17}{30}, \frac{17}{30}]$

So Now What?

We have:

$$\frac{13}{30} \leq f(11, 5) \leq \frac{11}{25} \quad \text{Diff} = 0.006666\dots$$

So Now What?

We have:

$$\frac{13}{30} \leq f(11, 5) \leq \frac{11}{25} \quad \text{Diff} = 0.006666\dots$$

Options:

1. $f(11, 5) = \frac{11}{25}$. Need to find procedure.
2. $f(11, 5) = \frac{13}{30}$. Need to find new technique for upper bounds.
3. $f(11, 5)$ in between. Need to find both.
4. $f(11, 5)$ unknown to science!

Vote

So Now What?

We have:

$$\frac{13}{30} \leq f(11, 5) \leq \frac{11}{25} \quad \text{Diff} = 0.006666 \dots$$

Options:

1. $f(11, 5) = \frac{11}{25}$. Need to find procedure.
2. $f(11, 5) = \frac{13}{30}$. Need to find new technique for upper bounds.
3. $f(11, 5)$ in between. Need to find both.
4. $f(11, 5)$ unknown to science!

Vote WE SHOW: $f(11, 5) = \frac{13}{30}$. **Exciting** new technique!

Terminology: Buddy

Assume that in some protocol every muffin is cut into two pieces.

Let x be a piece from muffin M .

The *other piece* from muffin M is the *buddy* of x .

Note that the buddy of x is of size

$$1 - x.$$

$f(11, 5) = \frac{13}{30}$, Easy Case Based on Muffins

There is a procedure for 11 muffins, 5 students where each student gets $\frac{11}{5}$ muffins, smallest piece N . We want $N \leq \frac{13}{30}$.

Case 0: Some muffin is uncut. Cut it $(\frac{1}{2}, \frac{1}{2})$ and give both halves to whoever got the uncut muffin. Reduces to other cases.

$f(11, 5) = \frac{13}{30}$, Easy Case Based on Muffins

There is a procedure for 11 muffins, 5 students where each student gets $\frac{11}{5}$ muffins, smallest piece N . We want $N \leq \frac{13}{30}$.

Case 0: Some muffin is uncut. Cut it $(\frac{1}{2}, \frac{1}{2})$ and give both halves to whoever got the uncut muffin. Reduces to other cases.

Case 1: Some muffin is cut into ≥ 3 pieces. $N \leq \frac{1}{3} < \frac{13}{30}$.

$f(11, 5) = \frac{13}{30}$, Easy Case Based on Muffins

There is a procedure for 11 muffins, 5 students where each student gets $\frac{11}{5}$ muffins, smallest piece N . We want $N \leq \frac{13}{30}$.

Case 0: Some muffin is uncut. Cut it $(\frac{1}{2}, \frac{1}{2})$ and give both halves to whoever got the uncut muffin. Reduces to other cases.

Case 1: Some muffin is cut into ≥ 3 pieces. $N \leq \frac{1}{3} < \frac{13}{30}$.

Negation of Case 0 and Case 1: All muffins cut into 2 pieces.

$f(11, 5) = \frac{13}{30}$, Easy Case Based on Students

Case 2: Some student gets ≥ 6 pieces.

$$N \leq \frac{11}{5} \times \frac{1}{6} = \frac{11}{30} < \frac{13}{30}.$$

$f(11, 5) = \frac{13}{30}$, Easy Case Based on Students

Case 2: Some student gets ≥ 6 pieces.

$$N \leq \frac{11}{5} \times \frac{1}{6} = \frac{11}{30} < \frac{13}{30}.$$

Case 3: Some student gets ≤ 3 pieces.

One of the pieces is

$$\geq \frac{11}{5} \times \frac{1}{3} = \frac{11}{15}.$$

Look at the muffin it came from to find a piece that is

$$\leq 1 - \frac{11}{15} = \frac{4}{15} < \frac{13}{30}.$$

$f(11, 5) = \frac{13}{30}$, Easy Case Based on Students

Case 2: Some student gets ≥ 6 pieces.

$$N \leq \frac{11}{5} \times \frac{1}{6} = \frac{11}{30} < \frac{13}{30}.$$

Case 3: Some student gets ≤ 3 pieces.

One of the pieces is

$$\geq \frac{11}{5} \times \frac{1}{3} = \frac{11}{15}.$$

Look at the muffin it came from to find a piece that is

$$\leq 1 - \frac{11}{15} = \frac{4}{15} < \frac{13}{30}.$$

Negation of Cases 2 and 3: Every student gets 4 or 5 pieces.

$f(11, 5) = \frac{13}{30}$, Fun Cases

Case 4: Every muffin is cut in 2 pieces, every student gets 4 or 5 pieces. Number of pieces: 22. Note ≤ 11 pieces are $> \frac{1}{2}$.

- ▶ s_4 is number of students who get 4 pieces
- ▶ s_5 is number of students who get 5 pieces

$$4s_4 + 5s_5 = 22$$

$$s_4 + s_5 = 5$$

$s_4 = 3$: There are 3 students who have 4 shares.

$s_5 = 2$: There are 2 students who have 5 shares.

We call a share that goes to a person who gets 4 shares a **4-share**.

We call a share that goes to a person who gets 5 shares a **5-share**.

$f(11, 5) = \frac{13}{30}$, Fun Cases

Case 4.1: Some 4-share is $\leq \frac{1}{2}$.

Alice gets w, x, y, z and $w \leq \frac{1}{2}$.

Since $w + x + y + z = \frac{11}{5}$ and $w \leq \frac{1}{2}$

$$x + y + z \geq \frac{11}{5} - \frac{1}{2} = \frac{17}{10}$$

Let x be the largest of x, y, z

$$x \geq \frac{17}{10} \times \frac{1}{3} = \frac{17}{30}$$

Look at **buddy** of x .

$$B(x) \leq 1 - x = 1 - \frac{17}{30} = \frac{13}{30}$$

GREAT! This is where $\frac{13}{30}$ comes from!

$$f(11, 5) = \frac{13}{30}, \text{ Fun Cases}$$

Case 4.2: All 4-shares are $> \frac{1}{2}$. There are $4s_4 = 12$ 4-shares.
There are ≥ 12 pieces $> \frac{1}{2}$. Can't occur.

Other Techniques

Here are the list of the Techniques we came up with:

1. Floor-Ceiling
2. Half
3. INT
4. GAP
5. Easy buddy-match
6. Hard buddy-match
7. Train (only worked on 3 (m, s) 's).

Other Techniques

Here are the list of the Techniques we came up with:

1. Floor-Ceiling
2. Half
3. INT
4. GAP
5. Easy buddy-match
6. Hard buddy-match
7. Train (only worked on 3 (m, s) 's).

Time to say we are NOT going to find a finite set of techniques that covers all cases and take what we got and write a book.

Later Results by Other People

1. In Fall 2018 Scott Huddleston emailed me code for an algorithm that, on input m, s , found $f(m, s)$ REALLY FAST.
2. Jacob and Erik Understand WHAT his algorithm does and Jacob coded it up to make sure he understood it. Jacob's code is also REALLY FAST.
3. Neither Scott, Bill, Jacob, or Erik had a proof that Scott's algorithm was fast (poly in m, s).
4. Richard Chatwin independently came up with the same algorithm; however, he also has a proof that it works. Its on arXiv.
5. One corollary of the work: $f(m, s)$ only depends on m/s .

How Our Book Worked

How Our Book Worked

We all had tasks we were good at and did those:

How Our Book Worked

We all had tasks we were good at and did those:

1. Erik: A Math Genius (solves muffin problems)

How Our Book Worked

We all had tasks we were good at and did those:

1. Erik: A Math Genius (solves muffin problems)
2. Jacob and Daniel: Programmers (codes up techniques)

How Our Book Worked

We all had tasks we were good at and did those:

1. Erik: A Math Genius (solves muffin problems)
2. Jacob and Daniel: Programmers (codes up techniques)
3. Bill: The Mastermind (guides the work and writes it up)

How Our Book Worked

We all had tasks we were good at and did those:

1. Erik: A Math Genius (solves muffin problems)
2. Jacob and Daniel: Programmers (codes up techniques)
3. Bill: The Mastermind (guides the work and writes it up)

We also all could do some of the other tasks.

How it worked

We kept increasing s .

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.
Example: **Erik, our techniques do not work on (35,13).**
2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.
Example: **Erik, our techniques do not work on (35,13).**
2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.
Example: **Erik, our techniques do not work on (35,13).**
2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:
Floor Ceiling,

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:
Floor Ceiling, Half,

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:
Floor Ceiling, Half, Int,

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:
Floor Ceiling, Half, Int,
Midpoint,

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:

Floor Ceiling, Half, Int,

Midpoint, Gaps,

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:

Floor Ceiling, Half, Int,

Midpoint, Gaps,

Easy Buddy-Match,

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:

Floor Ceiling, Half, Int,

Midpoint, Gaps,

Easy Buddy-Match, Hard buddy-Match,

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:

Floor Ceiling, Half, Int,

Midpoint, Gaps,

Easy Buddy-Match, Hard buddy-Match, Train

How it worked

We kept increasing s .

1. Bill tells Erik the least case we can't do.

Example: **Erik, our techniques do not work on (35,13).**

2. Erik solves and sends Bill a 1-page sketch.
3. Bill fills in the details and obtains general technique.
4. Jacob & Daniel code up technique and find least case that can't be done. For example **We can't do (47,23).**
5. Sends to Bill who either does it or Goto Step 1.

This happened 7 times leading to techniques now called:

Floor Ceiling, Half, Int,

Midpoint, Gaps,

Easy Buddy-Match, Hard buddy-Match, Train

Also a chapter that sketched out Scott H's method.

I meet Alan Frank!

I emailed Alan Frank, the **creator** of the Muffin Problem and we planned to meet at the MIT combinatorics seminar where I was scheduled to give a talk.

I meet Alan Frank!

I emailed Alan Frank, the **creator** of the Muffin Problem and we planned to meet at the MIT combinatorics seminar where I was scheduled to give a talk.

- ▶ He was delighted that his innocent problem, that he viewed as recreational, has lead to so much math of interest.

I meet Alan Frank!

I emailed Alan Frank, the **creator** of the Muffin Problem and we planned to meet at the MIT combinatorics seminar where I was scheduled to give a talk.

- ▶ He was delighted that his innocent problem, that he viewed as recreational, has lead to so much math of interest.
- ▶ He brought to the seminar 11 muffins:
1 cut $(\frac{15}{30}, \frac{15}{30})$, 2 cut $(\frac{14}{30}, \frac{16}{30})$, 8 cut $(\frac{13}{30}, \frac{17}{30})$.

I meet Alan Frank!

I emailed Alan Frank, the **creator** of the Muffin Problem and we planned to meet at the MIT combinatorics seminar where I was scheduled to give a talk.

- ▶ He was delighted that his innocent problem, that he viewed as recreational, has lead to so much math of interest.
- ▶ He brought to the seminar 11 muffins:
1 cut $(\frac{15}{30}, \frac{15}{30})$, 2 cut $(\frac{14}{30}, \frac{16}{30})$, 8 cut $(\frac{13}{30}, \frac{17}{30})$.
The five us of took pieces so we each got $\frac{11}{5}$ muffins.

I meet Alan Frank!

I emailed Alan Frank, the **creator** of the Muffin Problem and we planned to meet at the MIT combinatorics seminar where I was scheduled to give a talk.

- ▶ He was delighted that his innocent problem, that he viewed as recreational, has lead to so much math of interest.
- ▶ He brought to the seminar 11 muffins:
1 cut $(\frac{15}{30}, \frac{15}{30})$, 2 cut $(\frac{14}{30}, \frac{16}{30})$, 8 cut $(\frac{13}{30}, \frac{17}{30})$.
The five us of took pieces so we each got $\frac{11}{5}$ muffins.
- ▶ He does a Bike-For-Food Charity. I asked him if I should give \$40.00 a year OR my Royalties. He chose the \$40.00.

I meet Alan Frank!

I emailed Alan Frank, the **creator** of the Muffin Problem and we planned to meet at the MIT combinatorics seminar where I was scheduled to give a talk.

- ▶ He was delighted that his innocent problem, that he viewed as recreational, has lead to so much math of interest.
- ▶ He brought to the seminar 11 muffins:
1 cut $(\frac{15}{30}, \frac{15}{30})$, 2 cut $(\frac{14}{30}, \frac{16}{30})$, 8 cut $(\frac{13}{30}, \frac{17}{30})$.
The five us of took pieces so we each got $\frac{11}{5}$ muffins.
- ▶ He does a Bike-For-Food Charity. I asked him if I should give \$40.00 a year OR my Royalties. He chose the \$40.00.
First Year Royalties: \$50.00

I meet Alan Frank!

I emailed Alan Frank, the **creator** of the Muffin Problem and we planned to meet at the MIT combinatorics seminar where I was scheduled to give a talk.

- ▶ He was delighted that his innocent problem, that he viewed as recreational, has lead to so much math of interest.
- ▶ He brought to the seminar 11 muffins:
1 cut $(\frac{15}{30}, \frac{15}{30})$, 2 cut $(\frac{14}{30}, \frac{16}{30})$, 8 cut $(\frac{13}{30}, \frac{17}{30})$.
The five us of took pieces so we each got $\frac{11}{5}$ muffins.
- ▶ He does a Bike-For-Food Charity. I asked him if I should give \$40.00 a year OR my Royalties. He chose the \$40.00.
First Year Royalties: \$50.00
Second Year Royalties: \$40.00

I meet Alan Frank!

I emailed Alan Frank, the **creator** of the Muffin Problem and we planned to meet at the MIT combinatorics seminar where I was scheduled to give a talk.

- ▶ He was delighted that his innocent problem, that he viewed as recreational, has lead to so much math of interest.
- ▶ He brought to the seminar 11 muffins:
1 cut $(\frac{15}{30}, \frac{15}{30})$, 2 cut $(\frac{14}{30}, \frac{16}{30})$, 8 cut $(\frac{13}{30}, \frac{17}{30})$.
The five us of took pieces so we each got $\frac{11}{5}$ muffins.
- ▶ He does a Bike-For-Food Charity. I asked him if I should give \$40.00 a year OR my Royalties. He chose the \$40.00.
First Year Royalties: \$50.00
Second Year Royalties: \$40.00
Third Year and beyond Royalties: $\leq$ \$20.00

Final Thoughts and Advice

Final Thoughts and Advice

1. I got this problem **from a pamphlet**. Look around you!
Inspiration can come from anywhere!

Final Thoughts and Advice

1. I got this problem **from a pamphlet**. Look around you!
Inspiration can come from anywhere!
2. When I could not solve a muffin problem I got help. **Do not be shy about asking for help.**

Final Thoughts and Advice

1. I got this problem **from a pamphlet**. Look around you! Inspiration can come from anywhere!
2. When I could not solve a muffin problem I got help. **Do not be shy about asking for help.**
3. The co-authors were a diverse collection of strengths as mentioned before. This worked because

Final Thoughts and Advice

1. I got this problem **from a pamphlet**. Look around you! Inspiration can come from anywhere!
2. When I could not solve a muffin problem I got help. **Do not be shy about asking for help.**
3. The co-authors were a diverse collection of strengths as mentioned before. This worked because
Team Work Makes the Dream Work!