

using n equal sized intervals $[x_{i-1}, x_i]$, $1 \leq i \leq n$. If $h = 1/n$, demonstrate numerically how the error $e_n \equiv |I(f) - Q(f)|$ behaves as a function of h . This should be done by applying the rule for a sequence of values of n that are doubling in size, say $n = 8, 16, \dots, 1024$ (so that the interval width h is divided by 2) and showing that the errors are reduced by an appropriate factor with each successive refinement.

To make this demonstration clear, you should print a table showing, for each n , the values of n , h , the computed estimates, the error e_n , and the ratio of errors e_{n-1}/e_n .

Problem 5. (20)

a. The normal distribution function used in statistics is defined to be

$$\mathcal{N}(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt.$$

Use an adaptive quadrature rule of your choice to plot $\mathcal{N}(x)$ for $x \in [-4, 4]$. To do this, you need to replace the lower limit $-\infty$ in the integral with a finite number. For the plot, use the value determined from your experiments in part (b) below.

b. Explore the accuracy of the computed answer as a function of the value chosen for the lower limit. That is, for several choices of the lower limit, identify the maximum error for a set of points in $[-5, 5]$. You can use as an accurate answer the values obtained using -50 as the lower limit and a tolerance of 10^{-14} for the adaptive quadrature.