

WAIT FOR INSTRUCTIONS BEFORE BEGINNING

HONOR PLEDGE: "I pledge on my honor that I have not given or received any unauthorized assistance on this examination."

Signature and UID: _____

Print name: _____

- *Write your answers with enough detail about your approach and concepts used, so that the grader will be able to understand it easily. You should PROVE the correctness of your algorithms either directly or by referring to a proof in the book.*
- *The sum of the grades is 110, but your grades would be out of 100 (thus you get 10 bonus points by solving all the problems).*
- *Select the best choice for the first 5 problems and mark it by **X** in the table below*

Problem	1	2	3	4	5	6	7	8
a								
b								
c								
d								
e								

DO NOT WRITE BELOW THIS LINE

Problem 1-8:	
Problem 9:	
Problem 10:	
Problem 11:	
Problem 12:	
TOTAL:	

Multiple-choice Problems (Answer ONLY in the TABLE in the FIRST PAGE and NOT HERE):

Problem 1. (5 points) Let $T(n) = 3T\left(\left\lfloor \frac{n}{2} \right\rfloor\right) + 5n$ and $T(0) = 7$. Then $T(n)$ is?


- a) $\Theta(n)$ b) $\Theta(n^{\log_2 3} \log n)$ c) $\Theta(n \log n)$
d) $\Theta(n^{1.5})$ e) $\Theta(n^{\log_2 3})$

Problem 2. (5 points) Let $f(n) = 3n^2 + \log n$, $g(n) = n \log n + n \log^2 n$, and $h(n) = 4n^2 - 5n$. Select the correct answer.

- a) $f(n) = O(g(n)), f(n) = O(h(n))$ b) $f(n) = \Omega(g(n)), f(n) = \Theta(h(n))$
c) $f(n) = O(g(n)), f(n) = \Theta(h(n))$ d) $f(n) = \Theta(g(n)), f(n) = \Omega(h(n))$
e) None

Problem 3. (5 points) How many ways can you walk from A to B such you do not pass through C, D, and E. In each move you are allowed to go one step up or one step to the right. (Hint: use dynamic programming technique)

- a) 8 b) 10 c) 12
d) 14 e) 18



Problem 4. (5 points) Problem A is polynomially reducible to problem B. Which of the following statements is correct?

- I. If problem A is solvable in a polynomial time then problem B is solvable in polynomial time.
II. If problem A is NP-complete then problem B is NP-complete.
III. If problem A is NP-hard then problem B is NP-hard
- a) I b) II c) III
d) I and II e) I, II, and III

Problem 5. (5 points) For dense graphs with $\Omega(n^2)$ weighted edges what is the best running time for Prim's algorithm?

a) $O(n \log n)$

b) $O(n^2)$


c) $O(n^2 \log n)$

d) $O(n^3)$

e) $O(n \log^2 n)$

Problem 6. (5 points) Consider the weighted graph below. Run Dijkstra's algorithm starting from vertex 1. Write the vertices in the order which they are marked.

- a) 1, 5, 4, 6, 2, 3
- b) 1, 5, 4, 2, 6, 3
- c) 1, 5, 6, 2, 4, 3
- d) 1, 5, 4, 2, 3, 6
- e) 1, 5, 6, 2, 3, 4




Problem 7. (5 points) Consider the weighted graph of Problem 6. Run Prim's algorithm starting from vertex 1. Write the vertices in the order which they are marked.

- a) 1, 5, 4, 6, 2, 3
- b) 1, 5, 2, 3, 6, 4
- c) 1, 5, 6, 3, 2, 4
- d) 1, 5, 6, 2, 3, 4
- e) 1, 5, 4, 2, 6, 3

Problem 8. (5 points) Consider tree T below. Assume tree T is the DFS tree of undirected graph G. Which of the following cannot be an edge of graph G?

- a) (1, 5)
- b) (2, 6)
- c) (1, 6)
- d) (1, 3)
- e) (4, 3)



Regular Problems (you should always *PROVE THE CORRECTNESS* of your solutions):

Problem 9. Given a sequence of (not necessary positive) integers $x_1, x_2, \dots, x_n$ find a subsequence $x_i, x_{i+1}, \dots, x_j$ (of consecutive elements) such that the sum of the numbers in it is **even and maximum** over all subsequences of consecutive elements with an even sum of elements. For example if the sequence is $5, -1, 4, -10, 3, 3, 3$, the maximum even consecutive subsequence is $5, -1, 4$. The running time of your algorithm should be in $O(n)$.

Problem 10. Let T be a minimum spanning tree of a graph G , and let L be the sorted list of the edge weights of T . Prove that for any other minimum spanning tree T' of G , the list L is also the sorted list of edge weights on T' .

Problem 11. Given two sets S_1 and S_2 of natural numbers, and a natural number a . Find whether there exists an element x from S_1 and an element y from S_2 such that $2x + 5y = a$. The algorithm should run in time $O(n)$, where n is the total number of elements in both sets.

Problem 12. Let $d[v]$ be the discovery time and $f[v]$ be the finishing time of vertex v in DFS. Prove that it is not possible to find two vertices v and u such that $d[u] < d[v] < f[u] < f[v]$.

Problem 13. Explain Dijkstra's algorithm in details and prove its correctness.