

# Announcements

- Homework #7 is due tomorrow
- Midterm #2 is in 9 days

# Unit 10


# Functions

(We will return to probability next week.)

# Function

- A **function** assigns members of one set (the **domain**) to members of another set (**co-domain**)
- The **range** is the subset of the co-domain that gets “hit”



$$f : A \rightarrow B$$



# Function

- A member of the domain can only be assigned to *one* member of the co-domain

Are these functions?



# Function


There are many ways to express functions:

Let  $A = \{1, 2, 3\}$      $B = \{4, 8, 12\}$

$f : A \rightarrow B$  such that for all  $a \in A$ ,  $f(a) = 4a$

$f : A \rightarrow B$  such that  $a \mapsto 4a$


$f = \{ \langle 1, 4 \rangle, \langle 2, 8 \rangle, \langle 3, 12 \rangle \}$




# Total vs. Partial

- A **total function** assigns *every* member of the domain to an element of the co-domain
- A **partial function** may not assign every member of the domain

**Total Function**




**Partial Function**




# Injective Function

A function is **one-to-one** or **injective** if every element of the range is associated with *exactly one* element from the domain.

**Not an Injection**




**Injection**




# Surjective Function

A function is **onto** or **surjective** if the range is equal to the entire co-domain.

**Not a Surjection**



**Surjection**






# Bijjective Function

A function is **bijjective** if it is both injective (one-to-one) and surjective (onto)

## Bijection



Sometimes we call this a “one to one correspondence”

# Let's vote... (No discussion please!)

Consider the function  $f(x) = \sin(x)$

Which is true?

- A. Both one-to-one and onto (bijection)
- B. Neither one-to-one nor onto
- C. One-to-one but not onto
- D. Onto but not one-to-one
- E. I don't know

Answer: E

Domain and Co-Domain matter!


# Inverse Image

Let  $y$  be an element of the co-domain of a function.

The **inverse image** of  $y$  is the subset of the domain that maps to  $y$ .

What is the inverse image of...

- A?
- C?
- B?



# Inverse Function

Let  $f$  be a function. The **inverse** of  $f$ , denoted  $f^{-1}$ , is a function that “reverses”  $f$ .

$$\text{e.g.: } f(7) = \text{“aardvark”} \iff f^{-1}(\text{“aardvark”}) = 7$$

Not all functions have inverses.

Suppose  $f : A \rightarrow B$ , and consider  $f^{-1} : B \rightarrow A$

- What can we say about  $f^{-1}$  if  $f$  is not one-to-one?
- What can we say about  $f^{-1}$  if  $f$  is one-to-one but not onto?
- What can we say about  $f^{-1}$  if  $f$  is a bijection?

# Proving (or disproving) that a function is Injective

Let  $f: D \rightarrow C$  such that...

Claim:  $f$  is 1 to 1.

Proof:

Let  $a, b \in D$  such that  $f(a) = f(b)$ .

...

$a = b$

Claim:  $f$  is not 1 to 1.

Proof:

[Find two different elements of the domain that are mapped to the same value]

Examples:

- $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = 3x + 7$
- $f: \mathbb{R} - \{1\} \rightarrow \mathbb{R}$  such that  $f(x) = (x+1)/(x-1)$
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$  such that  $f(x) = x \bmod 7$

# Proving (or disproving) that a function is Surjective

Let  $f: D \rightarrow C$  such that...

Claim: f is onto.

Proof:

Let  $c \in C$  (arbitrarily selected).

...

there exists  $d \in D$  such that  $f(d) = c$ .

Claim: f is not onto.

Proof:

[Find an element of the codomain such for all  $d \in D$ ,  $f(d)$  is not equal to that element]

Examples:

- $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = 3x + 7$
- $f: \mathbb{R} \rightarrow \mathbb{Z}$  such that  $f(x) = \lfloor x/2 \rfloor$
- $f: \mathbb{R}^+ \rightarrow \mathbb{R}$  such that  $f(x) = \sqrt{x}$

# Proving a function is a bijection

Let  $f: D \rightarrow C$  such that...

Claim:  $f$  is a bijection.

Proof:


Part 1 [Prove  $f$  is one-to-one]...

Part 2 [Prove  $f$  is onto]...

# Composition of functions

Let  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$

–  $(g \circ f): X \rightarrow Z$  where  $(\forall x \in X)[(g \circ f)(x) = g(f(x))]$





# Composition of functions

Let  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$ .

What can we say about the domain and range of  $g \circ f$ ?

# Composition on finite sets- example

- Example

$$X = \{1,2,3\}, Y = \{a,b,c,d,e\}, Z = \{x,y,z\}$$

$f(1) = c$	$g(a) = y$	$(g \circ f)(1) = ?$
$f(2) = b$	$g(b) = y$	$(g \circ f)(2) = ?$
$f(3) = a$	$g(c) = z$	$(g \circ f)(3) = ?$
	$g(d) = x$	
	$g(e) = x$	

# Composition for infinite sets- example

$$f: \mathbb{Z} \rightarrow \mathbb{Z} \quad f(n) = n + 1$$



$$g: \mathbb{Z} \rightarrow \mathbb{Z} \quad g(n) = n^2$$

$$(g \circ f)(n) = g(f(n)) = g(n+1) = (n+1)^2$$

$$(f \circ g)(n) = f(g(n)) = f(n^2) = n^2 + 1$$

Note:  $g \circ f \neq f \circ g$

# The pigeonhole principle



- **Basic form:**

A function from one finite set to a smaller finite set cannot be one-to-one; there must be at least two elements in the domain that have the same image in the codomain.

# Pigeon Hole Principle

- Using this class as the domain:
  - must two people share a birth month?
  - must two people share a birthday?
- There exist two people in New York City who are not bald, and who have the same number of hairs on their heads.
- Pairs of people are either acquainted with one another or not. In any group of people, there must be two people who have the same number of acquaintances within the group.