

# Announcements

- Homework #2 due tomorrow at 11:00PM.
- Homework #3 will be posted tomorrow.

# One More Basic Example

- Claim:  $\mathbb{Q}$  is dense. (Assuming we already know about the closure of  $\mathbb{Q}$ .)

# More with Cases

- Claim: For all integers,  $x$  and  $y$ ,  $|x/y| = |x|/|y|$
- Claim: For all  $n \in \mathbb{N}$ ,  $3n^2 + n + 14$  is even.

# Fun Proof

An existence proof that is as close to “constructive” as you can get without actually being constructive...

- Claim: There are two irrational numbers,  $a$  and  $b$ , such that  $a^b$  is rational.

# Notation for “divisibility”

Suppose  $ab=c$ , where  $a,b,c \in \mathbb{Z}$  (with  $b \neq 0$ )

- We use the following notation to express “b divides c”

$$b|c$$

- In proofs, we frequently use the following interchangeably:

$$b|c \text{ is the same as } (\exists a \in \mathbb{Z})[c = ab]$$

# Implications

Below are outlines of the standard technique for proving implications:

**Claim:** If P then Q.

**Proof:**

Assume P.

...

Q.

Therefore,  $P \rightarrow Q$ .

**Claim:**  $(\forall x \in D)$  [If P(x) then Q(x)].

**Proof:**

Let  $d \in D$ , selected arbitrarily.

Assume P(d) holds.

...

Q(d).

Therefore,  $P(d) \rightarrow Q(d)$ .

Since d was selected arbitrarily,

$(\forall x \in D) [P(x) \rightarrow Q(x)]$ .

# Examples with implications

- Claim:  $\forall x, y, z \in \mathbb{N}$ : If  $x|y$ , and  $y|z$ , then  $x|z$ .
- Claim:  $\forall x, y \in \mathbb{R}$ , if  $x + y = 7$  and  $xy = 10$ , then  $x^2 + y^2 = 29$

# Proof by Contrapositive

Sometimes implications are easier to prove this way:

**Claim:** If P then Q.

**Proof:**

Assume  $\sim Q$ .

...

$\sim P$ .

Therefore,  $P \rightarrow Q$ .



# Examples using Contrapositive

- $(\forall n \in \mathbb{N})$  If  $3n + 2$  is odd then  $n$  is odd.
- $(\forall n, a, b \in \mathbb{R}^+)$  If  $n = ab$  then  $a \leq \sqrt{n}$  or  $b \leq \sqrt{n}$ .

# Proofs of Equivalence (“If and only if”)

Two techniques:

Claim:  $P \leftrightarrow Q$ .

Proof:

$P \leftrightarrow S1$

$\leftrightarrow S2$

$\leftrightarrow S3$

...

$\leftrightarrow Q$

- Doesn't always work
- Easy to make mistakes
- Maybe less writing

Claim:  $P \leftrightarrow Q$ .

Proof:

**Part I.** [Show  $P \rightarrow Q$ ]

...

**Part II.** [Show  $Q \rightarrow P$ ]

- Works more often
- Less error prone
- Probably more writing

# Be careful!

Critique this “proof”.

**Warning: This proof is invalid!**

**Claim:**  $(\forall n \in \mathbb{N})[n \text{ is odd} \leftrightarrow n^2 \text{ is odd}]$

**Proof:**

Let  $a \in \mathbb{N}$ , selected arbitrarily.

$a \text{ is odd} \leftrightarrow a = 2k + 1$  (some  $k \in \mathbb{N}$ )

$$\leftrightarrow a^2 = (2k + 1)(2k + 1)$$

$$\leftrightarrow a^2 = 4k^2 + 4k + 1$$

$$\leftrightarrow a^2 = 2(2k^2 + 2k) + 1$$

$$\leftrightarrow a^2 \text{ is odd (since } 2k^2 + 2k \in \mathbb{N}, \text{ by closure)}$$

Since  $a$  was selected arbitrarily, the proposition is true for all  $n \in \mathbb{N}$ .

# Proofs of Equivalence (if and only if)

- Claim:  $(\forall n \in \mathbb{N})[n \text{ is odd} \leftrightarrow n^2 \text{ is odd}]$