Announcements

• Homework #6 due on Friday. We may not grade *every* question.

Today's Password

lamb

Review

- (Regular) Induction
- Strong Induction

Another recurrence relation example

• Assume the following definition of a sequence:

$$a_0 = 1 \qquad a_1 = 2$$
$$(\forall k \in Z^{\ge 2}) [a_k = a_{k-1} + a_{k-2}]$$

Prove the following definition property, using strong induction:

$$(\forall n \in Z^{\geq 0}) \left[a_n \leq 2^n \right]$$

Another example- a divisibility property

• Assume the following definition of a recurrence relation:

$$a_0 = 0$$

 $a_1 = 7$
 $(\forall i \ge 2) [a_i = 2a_{i-1} + 3a_{i-2}]$

• Prove using strong induction that all elements in this relation have this property:

$$(\forall n \in N) [a_n \equiv 0 \pmod{7}]$$

Yet another one...

• Assume the following definition of a recurrence relation:

$$a_{0} = 0$$

$$a_{1} = 4$$

$$(\forall i \ge 2) [a_{i} = 6a_{i-1} - 5a_{i-2}]$$

• Prove using strong induction that all elements in this relation have this property:

$$(\forall n \in N) \left[a_n = 5^n - 1 \right]$$

Another example

 Theorem: for all n ≥ 2: n can be expressed as the product of primes. (Note that we consider a single prime factor to be a "product" of primes.)

This is "half" of the Unique Prime Factorization Theorem. The other half would be to show that the prime factorization is *unique*.

Chocolate Bar Division

Suppose you have a chocolate bar that is sectioned off into n squares, arranged in a rectangle. You can break the bar into pieces along the lines separating the squares. (Each break must go all the way across the current piece.)

Claim: It will always take n-1 breaks to separate the bar into individual squares. (No matter how you proceed!)