

Announcements

- Homework #10 is due Friday.
- Exam #2 is on Tuesday 4/30.

Unit 11

Relations

Relations

A **relation** (among sets) is a subset of their Cartesian product.

Relations can involve any number of sets, but frequently they are **binary** (two sets).

Examples of Binary Relations

Let $S = \{\text{Students at Maryland}\}$

Let $F = \{\text{faculty members at Maryland}\}$

Define relation R on $S \times F$ by:

$R = \{ \langle x, y \rangle \in S \times F : x \text{ has been in a class taught by } y \}$

Notation:

aRb means $\langle a, b \rangle \in R$

Examples of Binary Relations

- Any predicate with two free variables (over fixed domains) defines a binary relation over the same domains:


For all $x \in A, y \in B$

$$xRy \text{ iff } P(x, y) \text{ is true}$$

- $<$ is a binary relation on $\mathbb{R} \times \mathbb{R}$, or $\mathbb{Z} \times \mathbb{Z}$, etc.
- Any function can be thought of as a binary relation
(Can any binary relation be thought of as a function?)
- $=$ can be thought of as a (simple) binary relation over any domain

Ways to represent Binary Relations

- Arrow Diagrams





- Set Notation

$R = \{ \langle 5, A \rangle, \langle 2, A \rangle, \langle 12, A \rangle, \langle 12, C \rangle, \langle 12, D \rangle \}$

Ways to represent Binary Relations

- Graphs



Ways to represent Binary Relations

- Matrix Representation

$R = \{(2,1), (3,1), (3,2)\}$ could also be represented as:

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 1 & 1 \end{bmatrix} \end{matrix}$$

Ternary Relations

Examples:

- Let $R \subseteq \mathbb{Z} \times \mathbb{Z} \times \mathbb{N}$ be defined by:

$\langle a, b, c \rangle \in R$ if and only if $a \equiv_c b$

Alternate notation (like a predicate):

$R(a, b, c)$ holds if and only if $a \equiv_c b$.

- Let $R \subseteq \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ be defined by:

$\langle a, b, c \rangle \in R$ iff there could be a triangle with sides of lengths a , b and c .

Unary Relations

What would a **Unary Relation** look like?

Examples?

n-ary Relations

Relations can involve any number of sets.

Example:

Let $n \in \mathbb{N}^+$

Define $R \subseteq \mathbb{R}^n$ ($\mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \dots \times \mathbb{R}$) as:

$$\langle x_1, x_2, x_3, \dots, x_n \rangle \in R \quad \text{if and only if} \quad \sqrt{x_1^2 + x_2^2 + x_3^2 + \dots + x_n^2} \leq 1$$

What is the geometric interpretation for...

$n=2$?

$n=3$?

$n=1$?

$n=4$???

Properties of Binary Relations

Reflexive	$(\forall a \in A) [aRa]$
Irreflexive	$(\forall a \in A) [a \not R a]$
Symmetric	$(\forall a, b \in A) [aRb \rightarrow bRa]$
Antisymmetric	$(\forall a, b \in A) [aRb \wedge bRa \rightarrow a = b]$
Asymmetric	$(\forall a, b \in A) [aRb \rightarrow b \not R a]$
Non-symmetric	$(\forall a, b \in A) [a \neq b \rightarrow (aRb \leftrightarrow b \not R a)]$
Transitive	$(\forall a, b, c \in A) [aRb \wedge bRc \rightarrow aRc]$

Which Properties Hold?

Which of the properties on the previous slide hold for...

- $<$ over \mathbb{R}
- $=$ over the set $\{A, B, C\}$
- R over \mathbb{N} such that aRb iff a is a factor of b
- R over \mathbb{N} such that aRb iff $a \equiv_7 b$
- **R** over {students in this class} such that aRb iff a considers b to be a friend