

Homework 11, Morally due Tue May 7, 3:30PM

THIS HW IS THREE PAGES!!!!!!!!!!

THROUGHOUT THIS HW: Unless I tell you what n is, assume n is large enough so that we can use our approximations and the Ind assumptions we made in class. (If you were not in class then — too bad for you!)

THROUGHOUT THIS HW: Assume the grader does not already know the material so clearly explain what you are doing or how you are setting up your program.

1. (0 points but if you don't show up to the final I will assume you got this problem wrong and you will get 0 points for this entire HW) WHEN IS THE FINAL? WHERE IS THE FINAL?
2. (40 points) We are going to put m balls into n boxes, uniformly at random. You can assume n is large so our approximations work.
 - (a) (20 points) What is the probability (approx) that some box has at least 4 balls in it?
 - (b) (20 points) Make a statement:
If $m \geq XXX$ then the prob that some box has 4 balls in it is $\geq \frac{3}{4}$.
FILL IN THE XXX. XXX will depend on n . Try to make XXX as small as possible. Please provide a number for the constant in-front of n .

GO TO NEXT PAGE

3. (30 points) Assume n is large enough for our approximations to work. Also assume that $n^{0.9}$ is an integer. We are going to put $n^{0.9}$ balls into n boxes, uniformly at random. (Yes you read that right.)
- (a) What is the prob that some box has ≥ 2 balls?
 - (b) What is the prob that some box has ≥ 3 balls?
 - (c) Find a CONSTANT XXX (So XXX is ind of n) such that the prob that some box has $\geq XXX$ balls is $\geq \frac{1}{2}$. Try to make XXX as large as possible. (You may use that $\binom{n}{k} \sim n^k$.)

SHOW all of your work so that if the grader is not familiar with the problem he can still understand your answer.

GO TO NEXT PAGE

4. (30 points) Let p_n be the EXACT answer to the n -person hatcheck problem. Recall that

$$p_n = \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} + \cdots + (-1)^n \frac{1}{n!}.$$

Write a program that does the following

For $2 \leq n \leq 20$

- (a) Compute p_n exactly. (To deal with the precision issues in your code, please keep the numerator and denominator of the computed fractions separately and perform only one division in the end.)
- (b) Compute $\frac{1}{e} - p_n$ (it may be positive or negative).
- (c) Put your results in a nice easy-to-read tables with first column n and the second column p_n and third column $\frac{1}{e} - p_n$.
- (d) Make a conjecture about when the $\frac{1}{e} - p_n$ is positive and when it is negative.
- (e) Make a conjecture about how quickly $|\frac{1}{e} - p_n|$ goes to zero. For example, maybe you think $|\frac{1}{e} - p_n| \sim \frac{\pi}{n}$. (HINT: I do not think that.)

Your answer should have the columns and the conjectures.