BILL AND EMILY RECORD LECTURE!!!!

UNTIMED PART OF FINAL IS MONDAY May 10 9:00AM. DEAD CAT WED May 12 9:00PM

FINAL IS MONDAY May 17 8:00PM-10:15PM

FILL OUT COURSE EVALS for ALL YOUR COURSES!!!

Problems with a Point Exploring Math and Computer Science

Authors: William Gasarch Clyde Kruskal

How This Book Came to Be

Book's Origin

- ► In 2003 Lance Fortnow started Complexity Blog
- ▶ In 2007 Bill Gasarch joined and it was a co-blog.
- In 2015 various book publishers asked us

Can you make a book out of your blog?

Lance declined but Bill said YES.

Book's Point

Bill took the posts that had the following format:

- ► make a point about mathematics
- ▶ do some math to <u>underscore</u> those points and made those into chapters.

Book's Point

Bill took the posts that had the following format:

- make a point about mathematics
- ▶ do some math to <u>underscore</u> those points and made those into chapters.

Caveat: Not every chapter is quite like that.

To quote Ralph Waldo Emerson

A foolish consistency is the hobgoblin of small minds.

Problems with a Point needed a subtitle. I proposed

Problems with a Point needed a subtitle.

I proposed

Problems with a Point: Mathematical Musing and Math to

make those Musings Magnificent

Problems with a Point needed a subtitle.

I proposed

Problems with a Point: Mathematical Musing and Math to make those Musings Magnificent

The publisher said NO!

Problems with a Point needed a subtitle.

I proposed

Problems with a Point: Mathematical Musing and Math to make those Musings Magnificent

The publisher said NO!

I proposed

Problems with a Point: Mathematical Meditations and

Computer Science Cogitations

Problems with a Point needed a subtitle.

I proposed

Problems with a Point: Mathematical Musing and Math to make those Musings Magnificent

The publisher said NO!

I proposed

Problems with a Point: Mathematical Meditations and Computer Science Cogitations

The publisher said NO!

Problems with a Point needed a subtitle.

I proposed

Problems with a Point: Mathematical Musing and Math to make those Musings Magnificent

The publisher said NO!

I proposed

Problems with a Point: Mathematical Meditations and Computer Science Cogitations

The publisher said NO!

The publisher wisely decided to be less cute and more informative: **Problems with a Point: Exploring Math and Computer Science**

After some samples of Bill's writing the publisher said

After some samples of Bill's writing the publisher said

Please Procure People to Polish Prose and Proofs of Problems with a Point

SO

After some samples of Bill's writing the publisher said

Please Procure People to Polish Prose and Proofs of Problems with a Point

SO

Clyde Kruskal became a co-author.

After some samples of Bill's writing the publisher said

Please Procure People to Polish Prose and Proofs of Problems with a Point

SO

Clyde Kruskal became a co-author.

Now onto some samples of the book!

Point: Students Can Give Strange Answers

The Paint Can Problem

From the Year 2000 Maryland Math Competition: There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

Work on it in groups! Prove a General Theorem.

The Paint Can Problem

From the Year 2000 Maryland Math Competition: There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

Work on it in groups! Prove a General Theorem. Answer:

If there are 45 different colors of paint then we are done. Assume there are \leq 44 different colors. If all colors appear \leq 44 times then there are $44 \times 44 = 1936 < 2000$ cans of paint, a contradiction.

Note: this was Problem 1, which is supposed to be easy and indeed 95% got it right. What about the other 5%? Next slide.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- There are at least 45 cans that are different colors.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- There are at least 45 cans that are different colors.

ANSWER:

Paint cans are grey. Hence there are all the same color. Therefore there are 2000 cans that are the same color.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

ANSWER:

Paint cans are grey. Hence there are all the same color. Therefore there are 2000 cans that are the same color.

What do you think:

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

ANSWER:

Paint cans are grey. Hence there are all the same color. Therefore there are 2000 cans that are the same color.

What do you think:

Thats just stupid. 0 points.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

ANSWER:

Paint cans are grey. Hence there are all the same color. Therefore there are 2000 cans that are the same color.

What do you think:

- ► Thats just stupid. 0 points.
- Question says cans of the same color.... The full 30 pts.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- There are at least 45 cans that are different colors.

ANSWER:

Paint cans are grey. Hence there are all the same color. Therefore there are 2000 cans that are the same color.

What do you think:

- ► Thats just stupid. 0 points.
- Question says cans of the same color.... The full 30 pts.
- ▶ Not only does he get 30 points, but everyone else should get 0.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- There are at least 45 cans that are different colors.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

ANSWER:

If you look at a paint color really really carefully there will be differences. Hence, even if two cans seem to both be (say) RED, they are really different. Therefore there are 2000 cans of different colors.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

ANSWER:

If you look at a paint color really really carefully there will be differences. Hence, even if two cans seem to both be (say) RED, they are really different. Therefore there are 2000 cans of different colors.

What do you think:

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- ▶ There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

ANSWER:

If you look at a paint color really really carefully there will be differences. Hence, even if two cans seem to both be (say) RED, they are really different. Therefore there are 2000 cans of different colors.

What do you think:

► Thats just stupid. 0 points.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

ANSWER:

If you look at a paint color really really carefully there will be differences. Hence, even if two cans seem to both be (say) RED, they are really different. Therefore there are 2000 cans of different colors.

What do you think:

- Thats just stupid. 0 points.
- ▶ Well... he's got a point. 30 points in fact.

There are 2000 cans of paint. Show that at least one of the following two statements is true:

- There are at least 45 cans of the same color.
- ▶ There are at least 45 cans that are different colors.

ANSWER:

If you look at a paint color really really carefully there will be differences. Hence, even if two cans seem to both be (say) RED, they are really different. Therefore there are 2000 cans of different colors.

What do you think:

- Thats just stupid. 0 points.
- ▶ Well... he's got a point. 30 points in fact.
- ▶ Not only does he get 30 points, but everyone else should get 0.

A Triangle Problem

From the year 2007 Maryland Math Competition.

QUESTION Let ABC be a fixed triangle. Let COL be any 2-coloring of the plane where each point is colored with red or green. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

A Triangle Problem

From the year 2007 Maryland Math Competition.

QUESTION Let ABC be a fixed triangle. Let COL be any 2-coloring of the plane where each point is colored with red or green. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

Note I think I was assigned to grade it since it **looks like** the kind of problem I would make up, even though I didn't. It was problem 5 (out of 5) and was hard. About 100 students tried it, 8 got full credit, 10 got partial credit

Funny Answers One

QUESTION Let ABC be a fixed triangle. Let COL be any 2-coloring of the plane where each point is colored with red or green. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

Funny Answers One

QUESTION Let ABC be a fixed triangle. Let COL be any 2-coloring of the plane where each point is colored with red or green. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

Funny Answer One

All the vertices are red because I can make them whatever color I want. I can also write at a 30 degree angle to the bottom of this paper (The students answer was written at a 30 degree angle to the bottom of the paper.) if thats what I feel like doing at the moment. Just like 2+2=5 if thats what my math teacher says. Math is pretty subjective anyway.

Was Student One Serious?

All the vertices are red because I can make them whatever color I want. I can also write at a 30 degree angle to the bottom of this paper (The students answer was written at a 30 degree angle to the bottom of the paper.) if thats what I feel like doing at the moment. Just like 2+2=5 if thats what my math teacher says. Math is pretty subjective anyway.

Was Student One Serious?

All the vertices are red because I can make them whatever color I want. I can also write at a 30 degree angle to the bottom of this paper (The students answer was written at a 30 degree angle to the bottom of the paper.) if thats what I feel like doing at the moment. Just like 2+2=5 if thats what my math teacher says. Math is pretty subjective anyway.

Theorem The students is not serious.

Was Student One Serious?

All the vertices are red because I can make them whatever color I want. I can also write at a 30 degree angle to the bottom of this paper (The students answer was written at a 30 degree angle to the bottom of the paper.) if thats what I feel like doing at the moment. Just like 2+2=5 if thats what my math teacher says. Math is pretty subjective anyway.

Theorem The students is not serious.

Proof Assume, by contradiction, that they are serious. Then they really think math is subjective. Hence they don't really understand math. Hence they would not have done well enough on Part I to qualify for Part II. But they took Part II. Contradiction.

QUESTION Let ABC be a fixed triangle. Let COL be any 2-coloring of the plane where each point is colored with red or green. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

QUESTION Let ABC be a fixed triangle. Let COL be any 2-coloring of the plane where each point is colored with red or green. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

I like to think that we live in a world where points are not judged by their color, but by the content of their character. Color should be irrelevant in the the plane. To prove that there exists a group of points where only one color is acceptable is a reprehensible act of bigotry and discrimination.

QUESTION Let ABC be a fixed triangle. Let COL be any 2-coloring of the plane where each point is colored with red or green. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

I like to think that we live in a world where points are not judged by their color, but by the content of their character. Color should be irrelevant in the the plane. To prove that there exists a group of points where only one color is acceptable is a reprehensible act of bigotry and discrimination.

Was Student Two Serious?

QUESTION Let ABC be a fixed triangle. Let COL be any 2-coloring of the plane where each point is colored with red or green. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

I like to think that we live in a world where points are not judged by their color, but by the content of their character. Color should be irrelevant in the the plane. To prove that there exists a group of points where only one color is acceptable is a reprehensible act of bigotry and discrimination.

Was Student Two Serious? Yes.

QUESTION Let ABC be a fixed triangle. Let COL be any 2-coloring of the plane where each point is colored with red or green. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

I like to think that we live in a world where points are not judged by their color, but by the content of their character. Color should be irrelevant in the the plane. To prove that there exists a group of points where only one color is acceptable is a reprehensible act of bigotry and discrimination.

Was Student Two Serious? Yes. About Justice!.

The Real Answer to Points in the Plane Problem

Each point in the plane is colored either red or green. Let ABC be a fixed triangle. Prove that there is a triangle DEF in the plane such that DEF is similar to ABC and the vertices of DEF all have the same color.

Fix a 2-coloring of the plane.


There are 3 equally-spaced mono points on *x*-axis

Proof Clearly there are two points on the x-axis of the same color: p_1 , p_2 are RED. If p_3 , the midpoint of p_1 , p_2 , is RED then p_1 , p_3 , p_2 are all RED. DONE. Hence we assume p_3 is GREEN.

Let p_4 be such that $|p_1 - p_4| = |p_2 - p_1|$. If p_4 is RED then p_4, p_1, p_2 are all RED. DONE. Hence we assume p_4 is GREEN.

Let p_5 be such that $|p_5 - p_2| = |p_2 - p_1|$. If p_5 is RED then p_1, p_2, p_5 are all RED. DONE. Hence we assume p_5 is GREEN.

Only case left p_3, p_4, p_5 are all GREEN. DONE.



Finish Proof By Picture




Figure: Triangle Similar to ABC with Monochromatic Vertices

P, Q, R are RED.

If T or U or S are RED then get RED Triangle similar to ABC.

If not then ALL of T, U, S are GREEN, so get GREEN triangle similar to ABC.



Point: What is a Pattern?

Simple Functions

Bill assigned the following in Discrete Math: For each of the following sequences find a **simple function** A(n) such that the sequence is $A(1), A(2), A(3), \ldots$

- **1**. 10, -17, 24, -31, 38, -45, 52, · · ·
- **2**. -1, 1, 5, 13, 29, 61, 125, · · ·
- **3**. 6, 9, 14, 21, 30, 41, 54, · · ·

Caveat: These are NOT trick questions. **Work on it in groups.**

Simple Functions

Bill assigned the following in Discrete Math: For each of the following sequences find a **simple function** A(n) such that the sequence is $A(1), A(2), A(3), \ldots$

- **1**. 10, -17, 24, -31, 38, -45, 52, · · ·
- 2. -1, 1, 5, 13, 29, 61, 125, ...
- **3**. 6, 9, 14, 21, 30, 41, 54, · · ·

Caveat: These are NOT trick questions.

Work on it in groups.

- 1. 10, -17, 24, -31, 38, -45, 52, \cdots $A(n) = (-1)^{n+1}(7n+3)$.
- 2. -1, 1, 5, 13, 29, 61, 125, \cdots $A(n) = 2^n 3$.
- 3. 6, 9, 14, 21, 30, 41, 54, \cdots $A(n) = n^2 + 5$.

One student, in earnest, emailed Bill the following:

We never defined Simple Function in class so I went to Wikipedia.

One student, in earnest, emailed Bill the following:

We never defined Simple Function in class so I went to Wikipedia. It said that A Simple Function is a linear combination of indicator functions of measurable sets. Is that what you want us to use?

One student, in earnest, emailed Bill the following:

We never defined Simple Function in class so I went to Wikipedia. It said that A Simple Function is a linear combination of indicator functions of measurable sets. Is that what you want us to use?

I doubt the student knows what those terms mean

One student, in earnest, emailed Bill the following:

We never defined Simple Function in class so I went to Wikipedia. It said that A Simple Function is a linear combination of indicator functions of measurable sets. Is that what you want us to use?

I doubt the student knows what those terms mean I doubt Clyde knows what those terms mean.

One student, in earnest, emailed Bill the following:

We never defined Simple Function in class so I went to Wikipedia. It said that A Simple Function is a linear combination of indicator functions of measurable sets. Is that what you want us to use?

I doubt the student knows what those terms mean I doubt Clyde knows what those terms mean. I don't know what these terms mean.

One student, in earnest, emailed Bill the following:

We never defined Simple Function in class so I went to Wikipedia. It said that A Simple Function is a linear combination of indicator functions of measurable sets. Is that what you want us to use?

I doubt the student knows what those terms mean I doubt Clyde knows what those terms mean. I don't know what these terms mean.

I told him NO— all I wanted is an easy-to-describe function.

One student, in earnest, emailed Bill the following:

We never defined Simple Function in class so I went to Wikipedia. It said that A Simple Function is a linear combination of indicator functions of measurable sets. Is that what you want us to use?

I doubt the student knows what those terms mean I doubt Clyde knows what those terms mean.
I don't know what these terms mean.

I told him NO— all I wanted is an easy-to-describe function. I should have told him to use that def to see what he did.

One student, in earnest, emailed Bill the following:

We never defined Simple Function in class so I went to Wikipedia. It said that A Simple Function is a linear combination of indicator functions of measurable sets. Is that what you want us to use?


I doubt the student knows what those terms mean I doubt Clyde knows what those terms mean. I don't know what these terms mean.

I told him NO— all I wanted is an easy-to-describe function. I should have told him to use that def to see what he did. The student got the first one right, but left the other two blank.


When Do Patterns Hold?

The last question brings up the question of when patterns do and don't hold. We looked for cases where a pattern *did not* hold.

What is the max number of regions formed by connecting every pair of n points on a circle. For n = 1, 2, 3, 4, 5:




What is the max number of regions formed by connecting every pair of n points on a circle. For n = 1, 2, 3, 4, 5:




Based on this data what guess is tempting?

What is the max number of regions formed by connecting every pair of n points on a circle. For n = 1, 2, 3, 4, 5:




Based on this data what guess is tempting? 2^{n-1} .

What is the max number of regions formed by connecting every pair of n points on a circle. For n = 1, 2, 3, 4, 5:



Based on this data what guess is tempting? 2^{n-1} . But for n = 6, the number of regions is only 31.

What is the max number of regions formed by connecting every pair of n points on a circle. For n = 1, 2, 3, 4, 5:



Based on this data what guess is tempting? 2^{n-1} . But for n=6, the number of regions is only 31. The actual number of regions for n points is $\binom{n}{4}+\binom{n}{2}+1$.

Second Non-Pattern: Borwein Integrals

$$\int_0^\infty \frac{\sin x}{x} = \frac{\pi}{2}$$

$$\int_0^\infty \frac{\sin x}{x} \frac{\sin \frac{x}{3}}{\frac{x}{3}} = \frac{\pi}{2}$$

$$\vdots$$

$$\int_0^\infty \frac{\sin x}{x} \frac{\sin \frac{x}{3}}{\frac{x}{3}} \frac{\sin \frac{x}{5}}{\frac{x}{5}} \frac{\sin \frac{x}{7}}{\frac{x}{7}} \frac{\sin \frac{x}{9}}{\frac{x}{9}} \frac{\sin \frac{x}{11}}{\frac{x}{11}} \frac{\sin \frac{x}{13}}{\frac{x}{13}} = \frac{\pi}{2}$$

Second Non-Pattern: Borwein Integrals

$$\int_0^\infty \frac{\sin x}{x} = \frac{\pi}{2}$$

$$\int_0^\infty \frac{\sin x}{x} \frac{\sin \frac{x}{3}}{\frac{x}{3}} = \frac{\pi}{2}$$

$$\vdots$$

$$\int_0^\infty \frac{\sin x}{x} \frac{\sin \frac{x}{3}}{\frac{x}{3}} \frac{\sin \frac{x}{5}}{\frac{x}{5}} \frac{\sin \frac{x}{7}}{\frac{x}{7}} \frac{\sin \frac{x}{9}}{\frac{x}{9}} \frac{\sin \frac{x}{11}}{\frac{x}{11}} \frac{\sin \frac{x}{13}}{\frac{x}{13}} = \frac{\pi}{2}$$

But

$$\int_0^\infty \frac{\sin x}{x} \frac{\sin \frac{x}{3}}{\frac{x}{3}} \frac{\sin \frac{x}{5}}{\frac{x}{5}} \frac{\sin \frac{x}{7}}{\frac{x}{7}} \frac{\sin \frac{x}{9}}{\frac{x}{9}} \frac{\sin \frac{x}{11}}{\frac{x}{11}} \frac{\sin \frac{x}{13}}{\frac{x}{13}} \frac{\sin \frac{x}{15}}{\frac{x}{15}} =$$

935615849440640907310521750000



Why the breakdown at 15?

Because

$$\frac{1}{3} + \frac{1}{5} + \dots + \frac{1}{13} < 1$$

but

$$\frac{1}{3} + \frac{1}{5} + \dots + \frac{1}{15} > 1.$$

For more Google

Borwein Integral

Computers to FIND proofs vs Computers to DO Proofs

Colorings and Square Differences

The following are all true:

1. There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

The following are all true:

- 1. There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.
- 2. There exists a number W_3 such that, for all 3-colorings of $\{1, \ldots, W_3\}$ there exists 2 nums, square-apart, same color.

The following are all true:

- 1. There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.
- 2. There exists a number W_3 such that, for all 3-colorings of $\{1, \ldots, W_3\}$ there exists 2 nums, square-apart, same color.
- 3. There exists a number W_4 such that, for all 4-colorings of $\{1, \ldots, W_4\}$ there exists two nums, square-apart, same color.

The following are all true:

- 1. There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.
- 2. There exists a number W_3 such that, for all 3-colorings of $\{1, \ldots, W_3\}$ there exists 2 nums, square-apart, same color.
- 3. There exists a number W_4 such that, for all 4-colorings of $\{1, \ldots, W_4\}$ there exists two nums, square-apart, same color.
- 4. For all c there exists a number $W_c \ldots$

The following are all true:

- 1. There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.
- 2. There exists a number W_3 such that, for all 3-colorings of $\{1, \ldots, W_3\}$ there exists 2 nums, square-apart, same color.
- 3. There exists a number W_4 such that, for all 4-colorings of $\{1, \ldots, W_4\}$ there exists two nums, square-apart, same color.
- 4. For all c there exists a number $W_c \dots$

The proofs in the literature of these theorems give EEEEEEEENORMOUS bounds on W_2 , W_3 , W_4 , W_c . We look at easier proofs with two **points** in mind:

- ▶ Would they be good questions on a HS math competition?
- ▶ What is the role of Computers in these proofs?

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

Work on in groups and try to minimize W_2 .

Let COL be a 2-coloring of $\{1,2,3,\ldots\}$ with colorings R and B. We can assume COL(1)=R.

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

Work on in groups and try to minimize W_2 .

Let COL be a 2-coloring of $\{1,2,3,\ldots\}$ with colorings R and B. We can assume $\mathrm{COL}(1)=R$. Since 1 is a square $\mathrm{COL}(2)=B$.

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

```
Let \mathrm{COL} be a 2-coloring of \{1,2,3,\ldots\} with colorings R and R. We can assume \mathrm{COL}(1)=R. Since 1 is a square \mathrm{COL}(2)=R. Since 1 is a square \mathrm{COL}(3)=R.
```

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

```
Let \mathrm{COL} be a 2-coloring of \{1,2,3,\ldots\} with colorings R and B. We can assume \mathrm{COL}(1)=R. Since 1 is a square \mathrm{COL}(2)=B. Since 1 is a square \mathrm{COL}(3)=R. Since 1 is a square \mathrm{COL}(4)=B.
```

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

```
Let \mathrm{COL} be a 2-coloring of \{1,2,3,\ldots\} with colorings R and B. We can assume \mathrm{COL}(1)=R. Since 1 is a square \mathrm{COL}(2)=B. Since 1 is a square \mathrm{COL}(3)=R. Since 1 is a square \mathrm{COL}(4)=B. Since 1 is a square \mathrm{COL}(5)=R.
```

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

```
Let COL be a 2-coloring of \{1,2,3,\ldots\} with colorings R and B. We can assume COL(1) = R. Since 1 is a square COL(2) = B. Since 1 is a square COL(3) = R. Since 1 is a square COL(4) = B. Since 1 is a square COL(5) = R. AH-HA: COL(1) = COL(5) and 5 - 1 = 4 = 2^2. So W_2 < 5.
```

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

```
Let \mathrm{COL} be a 2-coloring of \{1,2,3,\ldots\} with colorings R and B. We can assume \mathrm{COL}(1)=R. Since 1 is a square \mathrm{COL}(2)=B. Since 1 is a square \mathrm{COL}(3)=R. Since 1 is a square \mathrm{COL}(4)=B. Since 1 is a square \mathrm{COL}(4)=B. Since 1 is a square \mathrm{COL}(5)=R. AH-HA: \mathrm{COL}(1)=\mathrm{COL}(5) and 5-1=4=2^2. So W_2\leq 5. AH-HA: RBRB shows that W_2<5.
```

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

```
Let COL be a 2-coloring of \{1,2,3,\ldots\} with colorings R and B. We can assume \mathrm{COL}(1)=R. Since 1 is a square \mathrm{COL}(2)=B. Since 1 is a square \mathrm{COL}(3)=R. Since 1 is a square \mathrm{COL}(4)=B. Since 1 is a square \mathrm{COL}(4)=B. Since 1 is a square \mathrm{COL}(5)=R. AH-HA: \mathrm{COL}(1)=\mathrm{COL}(5) and 5-1=4=2^2. So W_2\leq 5. AH-HA: RBRB shows that W_2\leq 5. So W_2=4.
```

There exists a number W_2 such that, for all 2-colorings of $\{1, \ldots, W_2\}$ there exists 2 nums, square-apart, same color.

Work on in groups and try to minimize W_2 .

```
Let COL be a 2-coloring of \{1,2,3,\ldots\} with colorings R and B. We can assume \mathrm{COL}(1)=R.
```

Since 1 is a square COL(2) = B.

Since 1 is a square COL(3) = R.

Since 1 is a square COL(4) = B.

Since 1 is a square COL(5) = R.

AH-HA: COL(1) = COL(5) and $5 - 1 = 4 = 2^2$. So $W_2 \le 5$.

AH-HA: RBRB shows that $W_2 \leq 5$.


So $W_2 = 4$.

Upshot Could be easy HS Math Comp Prob. No computer used.

There exists a number W_3 such that, for all 3-colorings of $\{1, \ldots, W_3\}$ there exists 2 nums, square-apart, same color.

There exists a number W_3 such that, for all 3-colorings of $\{1, \ldots, W_3\}$ there exists 2 nums, square-apart, same color.

There exists a number W_3 such that, for all 3-colorings of $\{1, \ldots, W_3\}$ there exists 2 nums, square-apart, same color.



Since $COL(x) = COL(x + 41) \dots$

Use COL(x) = COL(x + 41) to finish the proof and find upper bound on W_3 .

Since $COL(x) = COL(x + 41) \dots$

Use COL(x) = COL(x + 41) to finish the proof and find upper bound on W_3 .

$$\operatorname{COL}(1) = \operatorname{COL}(1+41) = \operatorname{COL}(1+2\times41) = \dots = \operatorname{COL}(1+41\times41)$$

Since
$$COL(x) = COL(x + 41) \dots$$

Use COL(x) = COL(x + 41) to finish the proof and find upper bound on W_3 .

$$COL(1) = COL(1+41) = COL(1+2\times41) = \cdots = COL(1+41\times41)$$

So 1 and 41^2 are a square apart and the same color.

$$W_3 \le 1 + 41^2 = 1682$$

Since
$$COL(x) = COL(x + 41) \dots$$

Use COL(x) = COL(x + 41) to finish the proof and find upper bound on W_3 .

$$COL(1) = COL(1+41) = COL(1+2\times41) = \cdots = COL(1+41\times41)$$

So 1 and 41^2 are a square apart and the same color.

$$W_3 \le 1 + 41^2 = 1682$$

Can we get better bound on W_3 ?

Better Bound on W₃




Figure: If $x \ge 10$ then COL(x) = COL(x+7), so $W_3 \le 59$

1. Problem 5 (so hard) on UMCP HS Math Comp, 2006: Show that for all 3-colorings of $\{1, \ldots, 2006\}$ there exists 2 numbers that are a square apart that are the same color

- 1. Problem 5 (so hard) on UMCP HS Math Comp, 2006: Show that for all 3-colorings of $\{1, \ldots, 2006\}$ there exists 2 numbers that are a square apart that are the same color
- 2. 240 took exam, 40 tried this problem, 10 got it right.


- 1. Problem 5 (so hard) on UMCP HS Math Comp, 2006: Show that for all 3-colorings of $\{1, \ldots, 2006\}$ there exists 2 numbers that are a square apart that are the same color
- 2. 240 took exam, 40 tried this problem, 10 got it right.
- 3. Bill Gasarch and Matt Jordan proved, by hand, $W_3 = 29$.

- 1. Problem 5 (so hard) on UMCP HS Math Comp, 2006: Show that for all 3-colorings of $\{1, \ldots, 2006\}$ there exists 2 numbers that are a square apart that are the same color
- 2. 240 took exam, 40 tried this problem, 10 got it right.
- 3. Bill Gasarch and Matt Jordan proved, by hand, $W_3 = 29$.
- Is there a HS-proof that W₄ exists? Bill wanted to put this problem on the next HS exam to find out. He was (wisely) told NO.

- 1. Problem 5 (so hard) on UMCP HS Math Comp, 2006: Show that for all 3-colorings of $\{1, \ldots, 2006\}$ there exists 2 numbers that are a square apart that are the same color
- 2. 240 took exam, 40 tried this problem, 10 got it right.
- 3. Bill Gasarch and Matt Jordan proved, by hand, $W_3 = 29$.
- Is there a HS-proof that W₄ exists? Bill wanted to put this problem on the next HS exam to find out. He was (wisely) told NO.
- 5. The question still remains: Is there a HS proof that W_4 exists?

- 1. Problem 5 (so hard) on UMCP HS Math Comp, 2006: Show that for all 3-colorings of $\{1, \ldots, 2006\}$ there exists 2 numbers that are a square apart that are the same color
- 2. 240 took exam, 40 tried this problem, 10 got it right.
- 3. Bill Gasarch and Matt Jordan proved, by hand, $W_3 = 29$.
- Is there a HS-proof that W₄ exists? Bill wanted to put this problem on the next HS exam to find out. He was (wisely) told NO.
- 5. The question still remains: Is there a HS proof that W_4 exists? YES. Discovered by Zach Price in 2019 via clever computer search. Next slide.

W_4 Exists: COL(x) = COL(x + 290, 085, 290)



Reflection on W₄

Reflection on W₄

1. Zach's proof shows $W_4 \le 1 + 299,085,290^2$. **PRO** Proof is easy to verify **CON** Number is large, proof does not generalize to W_5 .

Reflection on W_4

- 1. Zach's proof shows $W_4 \le 1 + 299,085,290^2$. PRO Proof is easy to verify CON Number is large, proof does not generalize to W_5 .
- The classical proof.
 PRO Gives bounds for W_c.
 CON Bounds are GINORMOUS, even for W₂.

Reflection on W₄

- 1. Zach's proof shows $W_4 \le 1 + 299,085,290^2$. PRO Proof is easy to verify CON Number is large, proof does not generalize to W_5 .
- The classical proof.
 PRO Gives bounds for W_c.
 CON Bounds are GINORMOUS, even for W₂.
- A Computer Search showed that W₄ = 58.
 PRO Get exact value.
 CON not human-verifiable. Does not generalize to W₅.

Which do you prefer?

How Amazon Prices Books

Our Book on Amazon?

```
The book cost more used than new!
https://www.amazon.com/
Problems-Point-Exploring-Computer-Science/dp/
9813279974
```

Our Book on Amazon?

The book cost more used than new! https://www.amazon.com/ Problems-Point-Exploring-Computer-Science/dp/ 9813279974

Chapter 1 of

Problems with two Points:

More Explorations of Math and Computer Science

will be

The Mathematics of Book Pricing on Amazon

BILL AND EMILY STOP RECORDING LECTURE!!!!

UNTIMED PART OF FINAL IS MONDAY May 10 9:00AM. DEAD CAT **WED MAY 12 9:00PM**

FINAL IS MONDAY May 17 8:00PM-10:15PM

FILL OUT COURSE EVALS for ALL YOUR COURSES!!!