

Homework 10 MORALLY Due April 28

1. (0 points) What is your name.

GO TO THE NEXT PAGE

2. (10 points)

(a) (10 points) Show that

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

with a combinatorial proof.

(Hint: Show that the RHS solves the question of how many ways to choose k objects out of n objects.)

(b) (0 points but you'll need this for a later problem on this HW set and maybe later in life as well.)

By convention $(\forall n \geq 0)[\binom{n}{0} = 1]$ and $(\forall k \geq 1)[\binom{0}{k} = 0]$.

From Part 1 that $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$.

Use these two equations to write a program that will do the following:

Given N, K outputs $\binom{k}{n}$ for all $0 \leq k \leq K$ and $0 \leq n \leq N$.

Run this program for $N = 52$ and $K = 6$.

GO TO THE NEXT PAGE

3. (30 points) Oliver-Poker is poker with a normal deck, but each player gets SIX cards.

For all questions here answer it BOTH in terms of binomial coefficient (e.g.,

$$\frac{\binom{12}{8}}{\binom{52}{6}}$$

)

and as an actual number to four places (e.g., 0.2192).

(For that you will use the output of the program you wrote in Problem 2.)

- (a) (10 points) What is the probability of getting three 2-of-a-kinds?
Example: $(2\heartsuit, 2\spadesuit, 4\heartsuit, 4\diamondsuit, 8\spadesuit, 8\clubsuit)$ is three 2-of-a-kind.
Counterexample: $(2\heartsuit, 2\spadesuit, 2\clubsuit, 2\diamondsuit, 8\spadesuit, 8\clubsuit)$ DOES NOT count as three 2-of-a-kind.
- (b) (10 points) What is the probability of getting a 2-of-a-kind and a 4-of-a-kind?
Example: $(2\heartsuit, 2\heartsuit, 2\diamondsuit, 8\spadesuit, 8\clubsuit)$ is a 4-of-a-kind and a 2-of-a-kind.
- (c) (10 points) What is the probability of getting a two 3-of-a-kind?
Example: $(2\heartsuit, 2\spadesuit, 2\heartsuit, 8\diamondsuit, 8\spadesuit, 8\clubsuit)$ is two 3-of-a-kind.
- (d) (0 points) Once its approved Oliver (also called *Poptart*) will be teaching a STIC on Poker (real poker, not this problem). Consider taking it!!

GO TO THE NEXT PAGE

4. (30 points) Show that there is no way to load two 8-sided dice to get fair sums.

GO TO THE NEXT PAGE

5. (30 points)

- (a) (0 points but you have to do it) Bill throws 2 fair 8-sided dice that are labeled in the normal way (each dice has faces with 1,2,3,4,5,6,7,8). For each $0 \leq i \leq 16$, give the probability of the sum of the dice being i .
- (b) (30 points) Give two 8-sided dice that are NOT labelled (1, 2, 3, 4, 5, 6, 7, 8). which, when rolled, give the same probabilities from Part *a*. (The labels have to all be natural numbers that are ≥ 1 .)