

# Review For Midterm One

# Review of Propositional Logic

# Propositional Logic

1.  $\wedge$ ,  $\vee$ ,  $\neg$ , *goes*

# Propositional Logic

1.  $\wedge$ ,  $\vee$ ,  $\neg$ , *goes*
2. Truth Tables (TT):

# Propositional Logic

1.  $\wedge$ ,  $\vee$ ,  $\neg$ , *goes*
2. Truth Tables (TT):
  - 2.1 Given a formula, can find the TT for it.

# Propositional Logic

1.  $\wedge$ ,  $\vee$ ,  $\neg$ , *goes*
2. Truth Tables (TT):
  - 2.1 Given a formula, can find the TT for it. Have class do some examples of this.

# Propositional Logic

1.  $\wedge$ ,  $\vee$ ,  $\neg$ , *goes*
2. Truth Tables (TT):
  - 2.1 Given a formula, can find the TT for it. Have class do some examples of this.
  - 2.2 Given a TT, can find a DNF formula for it.

# Propositional Logic

1.  $\wedge$ ,  $\vee$ ,  $\neg$ , *goes*
2. Truth Tables (TT):
  - 2.1 Given a formula, can find the TT for it. Have class do some examples of this.
  - 2.2 Given a TT, can find a DNF formula for it.  
Have class do some examples of this.
  - 2.3 A formula with  $n$  variables has a TT with  $2^n$  rows.

# Circuits

Recall from the last slide:

# Circuits

Recall from the last slide:

**Given a TT, can find a DNF formula for it.**

# Circuits

Recall from the last slide:

**Given a TT, can find a DNF formula for it.**

Easy: Given a Formula, you can make a circuit out of it.

# Circuits

Recall from the last slide:

**Given a TT, can find a DNF formula for it.**

Easy: Given a Formula, you can make a circuit out of it.

**Upshot** given a TT, can make a circuit for it.

# Circuits

Recall from the last slide:

**Given a TT, can find a DNF formula for it.**

Easy: Given a Formula, you can make a circuit out of it.

**Upshot** given a TT, can make a circuit for it.

Can use this to make circuits that compute arithmetic functions.

# Circuits

Recall from the last slide:

**Given a TT, can find a DNF formula for it.**

Easy: Given a Formula, you can make a circuit out of it.

**Upshot** given a TT, can make a circuit for it.

Can use this to make circuits that compute arithmetic functions.

**Discuss** how you write a circuit that tells, given an 8-bit number, outputs Y if prime and N if not prime.

# Circuits: Half-Adders and Full-Adders

# Circuits: Half-Adders and Full-Adders

1. Can do a circuit that inputs 2 bits and outputs the sum and the carry. Called a Half-Adder (HA).

# Circuits: Half-Adders and Full-Adders

1. Can do a circuit that inputs 2 bits and outputs the sum and the carry. Called a Half-Adder (HA).
2. Can do a circuit that inputs 3 bits and outputs the sum and the carry. Called a Full-Adder (HA).

## Circuits: Half-Adders and Full-Adders

1. Can do a circuit that inputs 2 bits and outputs the sum and the carry. Called a Half-Adder (HA).
2. Can do a circuit that inputs 3 bits and outputs the sum and the carry. Called a Full-Adder (FA).
3. Can combine HA and FAs to get create a circuit that adds two  $n$ -bit numbers.

# Logical Equivalence

# Logical Equivalence

1. Two formulas are equivalent if they have the same TT.

# Logical Equivalence

1. Two formulas are equivalent if they have the same TT.
2. Some well known equivalences:
  - 2.1  $\neg\neg p \equiv p$
  - 2.2  $\neg(p \vee q) \equiv \neg p \wedge \neg q.$
  - 2.3  $\neg(p \wedge q) \equiv \neg p \vee \neg q.$
  - 2.4 There are others.

# SATisfiability (SAT)

**Def**  $\phi(\vec{x}) \in \text{SAT}$  if there is  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .

If  $\vec{b}$  exists it is called a **SATisfying (SAT) Assignment**.

# SATisfiability (SAT)

**Def**  $\phi(\vec{x}) \in \text{SAT}$  if there is  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .

If  $\vec{b}$  exists it is called a **SATisfying (SAT) Assignment**.

$$(x_1 \vee x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2) \wedge (\neg x_2 \vee \neg x_3) \in \text{SAT?}$$

# SATisfiability (SAT)

**Def**  $\phi(\vec{x}) \in \text{SAT}$  if there is  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .

If  $\vec{b}$  exists it is called a **SATisfying (SAT) Assignment**.

$$(x_1 \vee x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2) \wedge (\neg x_2 \vee \neg x_3) \in \text{SAT?}$$

**Yes**  $x_1 = T, x_2 = F, x_3 = F$ .

# SATisfiability (SAT)

**Def**  $\phi(\vec{x}) \in \text{SAT}$  if there is  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .

If  $\vec{b}$  exists it is called a **SATisfying (SAT) Assignment**.

$$(x_1 \vee x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2) \wedge (\neg x_2 \vee \neg x_3) \in \text{SAT?}$$

**Yes**  $x_1 = T, x_2 = F, x_3 = F$ .

$$(x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_3) \wedge (\neg x_2 \vee \neg x_3) \wedge x_2 \in \text{SAT?}$$

# SATisfiability (SAT)

**Def**  $\phi(\vec{x}) \in \text{SAT}$  if there is  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .

If  $\vec{b}$  exists it is called a **SATisfying (SAT) Assignment**.

$$(x_1 \vee x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2) \wedge (\neg x_2 \vee \neg x_3) \in \text{SAT?}$$

**Yes**  $x_1 = T, x_2 = F, x_3 = F$ .

$$(x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_3) \wedge (\neg x_2 \vee \neg x_3) \wedge x_2 \in \text{SAT?}$$

**NO** Any SAT assignment needs  $x_2 = T$ . So question is:

# SATisfiability (SAT)

**Def**  $\phi(\vec{x}) \in \text{SAT}$  if there is  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .

If  $\vec{b}$  exists it is called a **SATisfying (SAT) Assignment**.

$$(x_1 \vee x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2) \wedge (\neg x_2 \vee \neg x_3) \in \text{SAT?}$$

**Yes**  $x_1 = T, x_2 = F, x_3 = F$ .

$$(x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_3) \wedge (\neg x_2 \vee \neg x_3) \wedge x_2 \in \text{SAT?}$$

**NO** Any SAT assignment needs  $x_2 = T$ . So question is:

$$x_1 \wedge (\neg x_1 \vee x_3) \wedge \neg x_3 \in \text{SAT?}$$

# SATisfiability (SAT)

**Def**  $\phi(\vec{x}) \in \text{SAT}$  if there is  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .

If  $\vec{b}$  exists it is called a **SATisfying (SAT) Assignment**.

$$(x_1 \vee x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2) \wedge (\neg x_2 \vee \neg x_3) \in \text{SAT?}$$

**Yes**  $x_1 = T, x_2 = F, x_3 = F$ .

$$(x_1 \vee \neg x_2) \wedge (\neg x_1 \vee x_3) \wedge (\neg x_2 \vee \neg x_3) \wedge x_2 \in \text{SAT?}$$

**NO** Any SAT assignment needs  $x_2 = T$ . So question is:

$$x_1 \wedge (\neg x_1 \vee x_3) \wedge \neg x_3 \in \text{SAT?}$$

In any SAT assignment need  $x_1 = T$  and  $x_3 = F$  so  $\neg x_1 \vee x_3$  is  $F$ .  
Hence NOT in SAT.

# Is There a Class of Formulas for Which SAT is in P?

We define several variants of SAT:

# Is There a Class of Formulas for Which SAT is in P?

We define several variants of SAT:

1. SAT is the set of all formulas that are in SAT.

# Is There a Class of Formulas for Which SAT is in P?

We define several variants of SAT:

1. SAT is the set of all formulas that are in SAT. That is,  
 $\phi(\vec{x}) \in SAT$  if there exists a vector  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .

# Is There a Class of Formulas for Which SAT is in P?

We define several variants of SAT:

1. SAT is the set of all formulas that are in SAT. That is,  $\phi(\vec{x}) \in SAT$  if there exists a vector  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .
2. CNFSAT is the set of all formulas in SAT of the form  $C_1 \wedge \cdots \wedge C_m$  where each  $C_i$  is an  $\vee$  of literals.

# Is There a Class of Formulas for Which SAT is in P?

We define several variants of SAT:

1. SAT is the set of all formulas that are in SAT. That is,  $\phi(\vec{x}) \in SAT$  if there exists a vector  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .
2. CNFSAT is the set of all formulas in SAT of the form  $C_1 \wedge \cdots \wedge C_m$  where each  $C_i$  is an  $\vee$  of literals.
3.  $k$ -CNFSAT is the set of all formulas in SAT of the form  $C_1 \wedge \cdots \wedge C_m$  where each  $C_i$  is an  $\vee$  of exactly  $k$  literals. Called  $kSAT$ .

# Is There a Class of Formulas for Which SAT is in P?

We define several variants of SAT:

1. SAT is the set of all formulas that are in SAT. That is,  $\phi(\vec{x}) \in \text{SAT}$  if there exists a vector  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .
2. CNFSAT is the set of all formulas in SAT of the form  $C_1 \wedge \cdots \wedge C_m$  where each  $C_i$  is an  $\vee$  of literals.
3.  $k$ -CNFSAT is the set of all formulas in SAT of the form  $C_1 \wedge \cdots \wedge C_m$  where each  $C_i$  is an  $\vee$  of exactly  $k$  literals. Called  $k$ SAT.
4. DNFSAT is the set of all formulas in SAT of the form  $C_1 \vee \cdots \vee C_m$  where each  $C_i$  is an  $\wedge$  of literals.

# Is There a Class of Formulas for Which SAT is in P?

We define several variants of SAT:

1. SAT is the set of all formulas that are in SAT. That is,  $\phi(\vec{x}) \in \text{SAT}$  if there exists a vector  $\vec{b}$  such that  $\phi(\vec{b}) = T$ .
2. CNFSAT is the set of all formulas in SAT of the form  $C_1 \wedge \cdots \wedge C_m$  where each  $C_i$  is an  $\vee$  of literals.
3.  $k$ -CNFSAT is the set of all formulas in SAT of the form  $C_1 \wedge \cdots \wedge C_m$  where each  $C_i$  is an  $\vee$  of exactly  $k$  literals. Called  $k$ SAT.
4. DNFSAT is the set of all formulas in SAT of the form  $C_1 \vee \cdots \vee C_m$  where each  $C_i$  is an  $\wedge$  of literals.
5.  $k$ -DNFSAT is the set of all formulas in SAT of the form  $C_1 \vee \cdots \vee C_m$  where each  $C_i$  is an  $\wedge$  of exactly  $k$  literals.

# Easy and Hard Forms for SAT

1. 2SAT is in P

# Easy and Hard Forms for SAT

1. 2SAT is in P Use 3 vars.

# Easy and Hard Forms for SAT

1. 2SAT is in P Use 3 vars.

Give an example of a formula in 2-CNF that is in SAT.

# Easy and Hard Forms for SAT

1. 2SAT is in P Use 3 vars.

Give an example of a formula in 2-CNF that is in SAT.

Give an example of a formula in 2-CNF that is not in SAT.

2. DNFSAT is in P

# Easy and Hard Forms for SAT

1. 2SAT is in P Use 3 vars.

Give an example of a formula in 2-CNF that is in SAT.

Give an example of a formula in 2-CNF that is not in SAT.

2. DNFSAT is in P

Give an example of a formula in DNF that is in SAT.

# Easy and Hard Forms for SAT

1. 2SAT is in P Use 3 vars.  
Give an example of a formula in 2-CNF that is in SAT.  
Give an example of a formula in 2-CNF that is not in SAT.
2. DNFSAT is in P  
Give an example of a formula in DNF that is in SAT.  
Give an example of a formula in DDF that is not in SAT.
3. 3SAT is thought to not be in P (NP-complete).

# Easy and Hard Forms for SAT

1. 2SAT is in P Use 3 vars.  
Give an example of a formula in 2-CNF that is in SAT.  
Give an example of a formula in 2-CNF that is not in SAT.
2. DNFSAT is in P  
Give an example of a formula in DNF that is in SAT.  
Give an example of a formula in DNF that is not in SAT.
3. 3SAT is thought to not be in P (NP-complete).  
Give an example of a formula in 3-CNF that is in SAT.

# Easy and Hard Forms for SAT

1. 2SAT is in P Use 3 vars.  
Give an example of a formula in 2-CNF that is in SAT.  
Give an example of a formula in 2-CNF that is not in SAT.
2. DNFSAT is in P  
Give an example of a formula in DNF that is in SAT.  
Give an example of a formula in DNF that is not in SAT.
3. 3SAT is thought to not be in P (NP-complete).  
Give an example of a formula in 3-CNF that is in SAT.  
Give an example of a formula in 3-CNF that is not in SAT.

# SAT is interesting

# SAT is interesting

The following is known:

$(3\text{-SAT is in P}) \leftrightarrow (\text{TSP is in P}) \leftrightarrow (\text{SCHEM is in P})$ .

# SAT is interesting

The following is known:

$(3\text{-SAT is in P}) \leftrightarrow (\text{TSP is in P}) \leftrightarrow (\text{SCHEG is in P})$ .

There are **thousands** of problems are equiv to SAT. Hence:

# SAT is interesting

The following is known:

$(3\text{-SAT is in P}) \leftrightarrow (\text{TSP is in P}) \leftrightarrow (\text{SCHEG is in P})$ .

There are **thousands** of problems are equiv to SAT. Hence:

- ▶ The complexity of 3-SAT is **important** since it relates to the complexity of many other problems.

# SAT is interesting

The following is known:

$(3\text{-SAT is in P}) \leftrightarrow (\text{TSP is in P}) \leftrightarrow (\text{SCHEG is in P})$ .

There are **thousands** of problems are equiv to SAT. Hence:

- ▶ The complexity of 3-SAT is **important** since it relates to the complexity of many other problems.
- ▶ Many of the problems 3-SAT is equivalent to have been worked on for 90 or more years; hence, it is unlikely they are in P. Hence it is unlikely that 3-SAT is in P.

# Review of Quantifier Logic

# Quantifiers

# Quantifiers

1.  $\exists x$  means exists  $x$ . Domain matters!

# Quantifiers

1.  $\exists x$  means exists  $x$ . Domain matters!  $\forall x$  means for all  $x$ .  
Domain matters!

# Quantifiers

1.  $\exists x$  means exists  $x$ . Domain matters!  $\forall x$  means for all  $x$ .  
Domain matters!
1. Give a  $D$  so that  $(\forall x)(\exists y)[x + y = 5]$  is true.

# Quantifiers

1.  $\exists x$  means exists  $x$ . Domain matters!  $\forall x$  means for all  $x$ .  
Domain matters!
1. Give a  $D$  so that  $(\forall x)(\exists y)[x + y = 5]$  is true.
2. Give a  $D$  so that  $(\forall x)(\exists y)[x + y = 1]$  is false.

# Quantifiers

1.  $\exists x$  means exists  $x$ . Domain matters!  $\forall x$  means for all  $x$ .  
Domain matters!
1. Give a  $D$  so that  $(\forall x)(\exists y)[x + y = 5]$  is true.
2. Give a  $D$  so that  $(\forall x)(\exists y)[x + y = 1]$  is false.
3. Give a  $D$  so that  $(\forall x)(\exists y)[xy = 5]$  is true.

# Quantifiers

1.  $\exists x$  means exists  $x$ . Domain matters!  $\forall x$  means for all  $x$ .  
Domain matters!
1. Give a  $D$  so that  $(\forall x)(\exists y)[x + y = 5]$  is true.
2. Give a  $D$  so that  $(\forall x)(\exists y)[x + y = 1]$  is false.
3. Give a  $D$  so that  $(\forall x)(\exists y)[xy = 5]$  is true.
4. Give a  $D$  so that  $(\forall x)(\exists y)[xy = 5]$  is false.

$$\neg(\exists x)[P(x)] \\ \equiv \\ (\forall x)[\neg P(x)]$$

# Quantifiers and Math

The domain is  $\mathbb{Z}$ .

# Quantifiers and Math

The domain is  $\mathbb{Z}$ .

How you would you write with quantifies:

**For all but a finite number of  $x$ ,  $P(x)$  happens.**

# Quantifiers and Math

The domain is  $\mathbb{Z}$ .

How you would you write with quantifies:

**For all but a finite number of  $x$ ,  $P(x)$  happens.**

How you would you write with quantifies:

**For an infinite number of  $x$ ,  $P(x)$  happens.**

# Quantifiers and Math

The domain is  $\mathbb{Z}$ .

How you would you write with quantifies:

**For all but a finite number of  $x$ ,  $P(x)$  happens.**

How you would you write with quantifies:

**For an infinite number of  $x$ ,  $P(x)$  happens.**

How you would you write with quantifies:

**For all but a finite number of  $x,y$   $P(x,y)$  happens.**

# Quantifiers and Math

The domain is  $\mathbb{Z}$ .

How you would you write with quantifies:

**For all but a finite number of  $x$ ,  $P(x)$  happens.**

How you would you write with quantifies:

**For an infinite number of  $x$ ,  $P(x)$  happens.**

How you would you write with quantifies:

**For all but a finite number of  $x, y$   $P(x, y)$  happens.**

How you would you write with quantifies:

**For an infinite number of  $x, y$ ,  $P(x, y)$  happens.**

# Quantifiers and Math

The domain is  $\mathbb{N}$ .

# Quantifiers and Math

The domain is  $\mathbb{N}$ .

How you would you write the following with quantifiers:

$x$  is the sum of 8 cubes.

# Quantifiers and Math

The domain is  $\mathbb{N}$ .

How you would you write the following with quantifiers:

$x$  is the sum of 8 cubes.

How you would you write the following with quantifiers:

# Quantifiers and Math

The domain is  $\mathbb{N}$ .

How you would you write the following with quantifiers:

$x$  is the sum of 8 cubes.

How you would you write the following with quantifiers:

**All but a finite number of numbers can be written as the sum of 7 cubes**

# Quantifiers and Math

The domain is  $\mathbb{N}$ .

How would you write the following with quantifiers:

$x$  is the sum of 8 cubes.

How would you write the following with quantifiers:

**All but a finite number of numbers can be written as the sum of 7 cubes**

**All but a finite number of numbers can be written as the sum of 7 cubes**  
**An infinite number of numbers cannot be written as the sum of 3 cubes**

# Quantifiers and Math

The domain is  $\mathbb{N}$ .

How would you write the following with quantifiers:

$x$  is the sum of 8 cubes.

How would you write the following with quantifiers:

**All but a finite number of numbers can be written as the sum of 7 cubes**

**All but a finite number of numbers can be written as the sum of 7 cubes  
An infinite number of numbers cannot be written as the sum of 3 cubes**

The above statements are true but hard to prove.

# Quantifiers and Math

The domain is  $\mathbb{N}$ .

How you would you write the following with quantifiers:

$x$  is the sum of 8 cubes.

How you would you write the following with quantifiers:

**All but a finite number of numbers can be written as the sum of 7 cubes**

**All but a finite number of numbers can be written as the sum of 7 cubes**  
**An infinite number of numbers cannot be written as the sum of 3 cubes**

The above statements are true but hard to prove.

The point of the math-with-quant slides is that we can STATE math of interest clearly using quantifiers.

# Vacuously True

Domain is  $\mathbb{N}$ .

# Vacuously True

Domain is  $\mathbb{N}$ .

$$(\forall x)[x < -1 \rightarrow x \text{ is prime}].$$

# Vacuously True

Domain is  $\mathbb{N}$ .

$$(\forall x)[x < -1 \rightarrow x \text{ is prime}].$$

True or False?

# Vacuously True

Domain is  $\mathbb{N}$ .

$$(\forall x)[x < -1 \rightarrow x \text{ is prime}].$$

True or False?

True.

# Vacuously True

Domain is  $\mathbb{N}$ .

$$(\forall x)[x < -1 \rightarrow x \text{ is prime}].$$

True or False?

True.

If you want to show this is false then have to find an  $x \in \mathbb{N}$  such that

$$x < -1 \text{ and } x \text{ is not prime}$$

# Vacuously True

Domain is  $\mathbb{N}$ .

$$(\forall x)[x < -1 \rightarrow x \text{ is prime}].$$

True or False?

True.

If you want to show this is false then have to find an  $x \in \mathbb{N}$  such that

$$x < -1 \text{ and } x \text{ is not prime}$$

You can't! So the statement is true.

# An Interesting HW Question

For this problem we use the following standard terminology:

## An Interesting HW Question

For this problem we use the following standard terminology:

A domain is *dense* if  $(\forall x, y)[x < y \implies (\exists z)[x < z < y]]$ .

## An Interesting HW Question

For this problem we use the following standard terminology:

A domain is *dense* if  $(\forall x, y)[x < y \implies (\exists z)[x < z < y]]$ .

A domain has a *min element* if  $(\exists x)(\forall y)[x \leq y]$ .

## An Interesting HW Question

For this problem we use the following standard terminology:

A domain is *dense* if  $(\forall x, y)[x < y \implies (\exists z)[x < z < y]]$ .

A domain has a *min element* if  $(\exists x)(\forall y)[x \leq y]$ .

A domain has a *max element* if  $(\exists x)(\forall y)[y \leq x]$ .

## An Interesting HW Question

For this problem we use the following standard terminology:

A domain is *dense* if  $(\forall x, y)[x < y \implies (\exists z)[x < z < y]]$ .

A domain has a *min element* if  $(\exists x)(\forall y)[x \leq y]$ .

A domain has a *max element* if  $(\exists x)(\forall y)[y \leq x]$ .

In this problem we list conditions on a domain. EITHER give a domain that satisfies the conditions OR state that there is NO such domain (no proof required).

## An Interesting HW Question

For this problem we use the following standard terminology:

A domain is *dense* if  $(\forall x, y)[x < y \implies (\exists z)[x < z < y]]$ .

A domain has a *min element* if  $(\exists x)(\forall y)[x \leq y]$ .

A domain has a *max element* if  $(\exists x)(\forall y)[y \leq x]$ .

In this problem we list conditions on a domain. EITHER give a domain that satisfies the conditions OR state that there is NO such domain (no proof required).

The list of conditions is on the next slide.

# Conditions

1)  $D$  is finite and dense.

# Conditions

1)  $D$  is finite and dense.

DOES EXIST. The domain is  $\emptyset$ . Dense vacuously.

## Conditions

1)  $D$  is finite and dense.

DOES EXIST. The domain is  $\emptyset$ . Dense vacuously.

2)  $D$  is finite and dense and has  $\geq 2$  elements.

## Conditions

1)  $D$  is finite and dense.

DOES EXIST. The domain is  $\emptyset$ . Dense vacuously.

2)  $D$  is finite and dense and has  $\geq 2$  elements.

DOES NOT EXIST. Since  $|D| \geq 2$  let  $x < y$  be in  $D$ . Then there exists  $z \in D$ ,  $x < z < y$ . Then there is a point between  $x$  and  $z$  and between  $z$  and  $y$ . Keep doing this.  $D$  is infinite.

## Conditions

1)  $D$  is finite and dense.

DOES EXIST. The domain is  $\emptyset$ . Dense vacuously.

2)  $D$  is finite and dense and has  $\geq 2$  elements.

DOES NOT EXIST. Since  $|D| \geq 2$  let  $x < y$  be in  $D$ . Then there exists  $z \in D$ ,  $x < z < y$ . Then there is a point between  $x$  and  $z$  and between  $z$  and  $y$ . Keep doing this.  $D$  is infinite.

3)  $D$  is infinite and not dense

## Conditions

1)  $D$  is finite and dense.

DOES EXIST. The domain is  $\emptyset$ . Dense vacuously.

2)  $D$  is finite and dense and has  $\geq 2$  elements.

DOES NOT EXIST. Since  $|D| \geq 2$  let  $x < y$  be in  $D$ . Then there exists  $z \in D$ ,  $x < z < y$ . Then there is a point between  $x$  and  $z$  and between  $z$  and  $y$ . Keep doing this.  $D$  is infinite.

3)  $D$  is infinite and not dense

DOES EXIST:  $\mathbb{Z}$ .

## Conditions

1)  $D$  is finite and dense.

DOES EXIST. The domain is  $\emptyset$ . Dense vacuously.

2)  $D$  is finite and dense and has  $\geq 2$  elements.

DOES NOT EXIST. Since  $|D| \geq 2$  let  $x < y$  be in  $D$ . Then there exists  $z \in D$ ,  $x < z < y$ . Then there is a point between  $x$  and  $z$  and between  $z$  and  $y$ . Keep doing this.  $D$  is infinite.

3)  $D$  is infinite and not dense

DOES EXIST:  $\mathbb{Z}$ .

4)  $D$  is infinite, has a min element, has a max element, and is dense.

## Conditions

1)  $D$  is finite and dense.

DOES EXIST. The domain is  $\emptyset$ . Dense vacuously.

2)  $D$  is finite and dense and has  $\geq 2$  elements.

DOES NOT EXIST. Since  $|D| \geq 2$  let  $x < y$  be in  $D$ . Then there exists  $z \in D$ ,  $x < z < y$ . Then there is a point between  $x$  and  $z$  and between  $z$  and  $y$ . Keep doing this.  $D$  is infinite.

3)  $D$  is infinite and not dense

DOES EXIST:  $\mathbb{Z}$ .

4)  $D$  is infinite, has a min element, has a max element, and is dense.

DOES EXIST:  $[0, 1]$ .

## Conditions

1)  $D$  is finite and dense.

DOES EXIST. The domain is  $\emptyset$ . Dense vacuously.

2)  $D$  is finite and dense and has  $\geq 2$  elements.

DOES NOT EXIST. Since  $|D| \geq 2$  let  $x < y$  be in  $D$ . Then there exists  $z \in D$ ,  $x < z < y$ . Then there is a point between  $x$  and  $z$  and between  $z$  and  $y$ . Keep doing this.  $D$  is infinite.

3)  $D$  is infinite and not dense

DOES EXIST:  $\mathbb{Z}$ .

4)  $D$  is infinite, has a min element, has a max element, and is dense.

DOES EXIST:  $[0, 1]$ .

5)  $D$  is infinite, has a min element, has a max element, and is NOT dense.

## Conditions

1)  $D$  is finite and dense.

DOES EXIST. The domain is  $\emptyset$ . Dense vacuously.

2)  $D$  is finite and dense and has  $\geq 2$  elements.

DOES NOT EXIST. Since  $|D| \geq 2$  let  $x < y$  be in  $D$ . Then there exists  $z \in D$ ,  $x < z < y$ . Then there is a point between  $x$  and  $z$  and between  $z$  and  $y$ . Keep doing this.  $D$  is infinite.

3)  $D$  is infinite and not dense

DOES EXIST:  $\mathbb{Z}$ .

4)  $D$  is infinite, has a min element, has a max element, and is dense.

DOES EXIST:  $[0, 1]$ .

5)  $D$  is infinite, has a min element, has a max element, and is NOT dense.

DOES EXIST:  $\{-1, -\frac{1}{2}, -\frac{1}{3}, \dots\} \cup \{\dots, \frac{1}{3}, \frac{1}{2}, 1\}$

# Sets

# Set Operations

$A, B, U$  are all sets.  $U$  is the universe we are in (e.g.,  $\mathbb{R}$ ).

# Set Operations

$A, B, U$  are all sets.  $U$  is the universe we are in (e.g.,  $\mathbb{R}$ ).

1.  $A \cup B = \{x : x \in A \vee x \in B\}$ .

# Set Operations

$A, B, U$  are all sets.  $U$  is the universe we are in (e.g.,  $\mathbb{R}$ ).

1.  $A \cup B = \{x : x \in A \vee x \in B\}$ .

2.  $A \cap B = \{x : x \in A \wedge x \in B\}$ .

# Set Operations

$A, B, U$  are all sets.  $U$  is the universe we are in (e.g.,  $\mathbb{R}$ ).

1.  $A \cup B = \{x : x \in A \vee x \in B\}$ .

2.  $A \cap B = \{x : x \in A \wedge x \in B\}$ .

3.  $\bar{A} = \{x \in U : x \notin A\}$ .

Note that  $\bar{A}$  only makes sense if you have a Universe.

# Empty Set

# Empty Set

1.  $\emptyset$  is the empty set.

1.1  $A \cup \emptyset = A$

# Empty Set

1.  $\emptyset$  is the empty set.

1.1  $A \cup \emptyset = A$

1.2  $A \cap \emptyset = \emptyset$ .

# Power Set

**Definition** Of  $A$  is a set then **the powerset of  $A$** , denoted  $P(A)$ , is the set of all subsets of  $A$ .

# Power Set

**Definition** Of  $A$  is a set then **the powerset of  $A$** , denoted  $P(A)$ , is the set of all subsets of  $A$ .

If  $A = \{4, 6, 9\}$  then  
 $P(A)$  is

$$\{\emptyset, \{4\}, \{6\}, \{9\}, \{4, 6\}, \{4, 9\}, \{6, 9\}, \{4, 6, 9\}\}$$

# Mods