

**Ramsey's Theorem**  
**for the Infinite Complete Graph and**  
**the Infinite Complete Hypergraph**  
Exposition by William Gasarch

## 1 Introduction

In this document we define notation for graphs and hypergraphs that we use for the course and then look at Ramsey's theorem and the Canonical Ramsey theory on  $\mathbb{N}$ . Why start with  $\mathbb{N}$ ? Because Joel Spencer said

**Infinite Ramsey Theory is easier than Finite Ramsey Theory**  
**because all of the messy constants go away.**

## 2 Notation

Recall that a graph is a set of vertices and a set of edges which are *unordered pairs* of vertices. Why pairs? We will generalize this by allowing edges to be sets of size 1, 2 (the usual case), 3, general  $a$  and not have any restriction on size.

### Notation 2.1

1. If  $n \geq 1$  then  $[n] = \{1, \dots, n\}$ .
2. If  $a \in \mathbb{N}$  and  $A$  is a set then  $\binom{A}{a}$  is the set of all subsets of  $A$  of size  $a$ . We commonly use  $\binom{[n]}{a}$  and  $\binom{\mathbb{N}}{a}$ .

**Def 2.2** Let  $a \in \mathbb{N}$  (note that  $a = 0$  is allowed). A  *$a$ -hypergraph* is a set of vertices  $V$  and a set of edges which is a subset of  $\binom{V}{a}$ .

### Examples

1. A 0-hypergraph is just a set of vertices. This is just stupid but we'll keep it around in case we need some edge case.
2. A 1-hypergraph is a set of vertices together with edges which are also vertices. So its just a set of vertices but some are also called edges.

3. A 2-hypergraph is the usual graphs you know and love.
4. A 3-hypergraph. Edges are sets of 3 vertices.  $V = \mathbf{N}$  and the edges are all  $(a, b, c)$  such that  $a + b + c \equiv 0 \pmod{9}$ . I *could not* have said  $a + 2b + 3c \equiv 0 \pmod{9}$  since then the order would matter. We are dealing with unordered hypergraphs. I could have said all  $(a, b, c)$  with  $a < b < c$  such that  $a + 2b + 3c \equiv 0 \pmod{9}$ .
5. Another example of a 3-hypergraph: let  $V$  be some set of points in the plane. Let the edges be all 3-sets of points that form non-degenerate triangles.

**Def 2.3** A *hypergraph* (notice the lack of a parameter) is a set of vertices  $V$  together with edges which are subsets of  $V$ .

**Example**

1.  $V = \mathbf{N}$  and we take the set of all finite subsets of  $\mathbf{N}$  whose sum is  $\equiv 0 \pmod{9}$ . Note that the empty set would be an edge.
2.  $V$  is a set of points in the plane. The edges are all of the lines in the plane.
3. Any  $a$ -hypergraph is also a hypergraph.

We are all familiar with the complete graph on  $\mathbf{N}$ :

**Notation 2.4**  $K_{\mathbf{N}}$  is the graph  $(V, E)$  where

$$\begin{aligned} V &= \mathbf{N} \\ E &= \binom{\mathbf{N}}{2} \end{aligned}$$

Here is the complete  $a$ -hypergraph on  $\mathbf{N}$ :

**Notation 2.5**  $K_{\mathbf{N}}^a$  is the hypergraph  $(V, E)$  where

$$\begin{aligned} V &= \mathbf{N} \\ E &= \binom{\mathbf{N}}{a} \end{aligned}$$

**Convention 2.6** In this course unless otherwise noted (1) a *coloring of a graph* is a coloring of the edges of the graph. and (2) a *coloring of a hypergraph* is a coloring of the edges of the hypergraph.

### 3 Ramsey Theory on the Complete 1-Hypergraph on $\mathbb{N}$

The following theorem is too obvious to prove but I want to state it:

**Theorem 3.1** *For every 2-coloring of  $\mathbb{N}$  there is an infinite  $A \subseteq \mathbb{N}$  that is the same color.*

Even though this is an easy theorem here are some questions:

1. Is there a finite version of this theorem?
2. If you are given a program that computes the coloring can you determine which color (or perhaps both) appears infinitely often?
3. What if you are given a simple computational device (e.g., a DFA with output). Then can you determine which color? What is the complexity of the problem?

What if I allow an infinite number of colors?

**Theorem 3.2** *For every coloring of  $\mathbb{N}$  there is either (1) an infinite  $A \subseteq \mathbb{N}$  that is the same color, or (2) an infinite  $A \subseteq \mathbb{N}$  that all have different colors (called a rainbow set).*

**Proof:** Let COL be a coloring of  $\mathbb{N}$ . We define an infinite sequence of vertices,

$$x_1, x_2, \dots,$$

and an infinite sequence of sets of vertices,

$$V_0, V_1, V_2, \dots,$$

that are based on COL.

Here is the intuition: Either COL(1) appears infinitely often (so we are done) or not. If not then we get rid of the finite number of vertices colored COL(1) except 1. We then do the same for COL(2). We will either find some color that appears infinitely often or create a sequence of all different colors.

We now describe it formally.

$$V_0 = \mathbf{N}$$

$$x_1 = 1$$

$$c_1 = \text{DONE if } |\{v \in V_0 \mid \text{COL}(v) = \text{COL}(x_1)\}| \text{ is infinite. And you are DONE! STOP}$$

$$= \text{COL}(x_1) \text{ otherwise}$$

$$V_1 = \{v \in V_0 \mid \text{COL}(v) \neq c_1\} \text{ (note that } |V_1| \text{ is infinite)}$$

Let  $i \geq 2$ , and assume that  $V_{i-1}$  is defined. We define  $x_i$ ,  $c_i$ , and  $V_i$ :

$$x_i = \text{the least number in } V_{i-1}$$

$$c_i = \text{DONE if } |\{v \in V_{i-1} \mid \text{COL}(v) = \text{COL}(x_i)\}| \text{ is infinite. And you are DONE! STOP}$$

$$= \text{COL}(x_i) \text{ otherwise}$$

$$V_i = \{v \in V_{i-1} \mid \text{COL}(v) \neq c_i\} \text{ (note that } |V_i| \text{ is infinite)}$$

How long can this sequence go on for? If ever it stops then we are done as we have found a color appearing infinitely often. If not then the sequence

$$x_1, x_2, \dots,$$

is infinite and each number in it is a different color, so we have found a rainbow set. ■

1. Is there a finite version of this theorem?
2. If you are given a program that computes the coloring can you determine which color (if any) appears infinitely often?
3. What if you are given a simple computational device (e.g., a DFA with output). Then can you determine which color? What is the complexity of the problem?

## 4 A Bit More Notation

For the case of the 1-hypergraph we didn't need notions of complete graphs or homog sets, though that is what we were talking about. For  $a$ -hypergraphs we will.

**Def 4.1** Let  $\text{COL}: \binom{\mathbb{N}}{2} \rightarrow [2]$ . Let  $V \subseteq \mathbb{N}$ . The set  $V$  is *homog* if there exists a color  $c$  such that every elements of  $\binom{V}{2}$  is colored  $c$ .

**Def 4.2** Let  $\text{COL}: \binom{\mathbb{N}}{k} \rightarrow [c]$ . Let  $V \subseteq \mathbb{N}$ . The set  $V$  is *homog* if there exists a color  $c$  such that every elements of  $\binom{V}{k}$  is colored  $c$ .

## 5 Ramsey's Theorem for the Infinite Complete Graphs

The following is Ramsey's Theorem for  $K_{\mathbb{N}}$ .

**Theorem 5.1** *For every 2-coloring of the edges of  $K_{\mathbb{N}}$  there is an infinite homog set.*

**Proof:** Let  $\text{COL}$  be a 2-coloring of  $K_{\mathbb{N}}$ . We define an infinite sequence of vertices,

$$x_1, x_2, \dots,$$

and an infinite sequence of sets of vertices,

$$V_0, V_1, V_2, \dots,$$

that are based on  $\text{COL}$ .

Here is the intuition: Vertex  $x_1 = 1$  has an infinite number of edges coming out of it. Some are RED, and some are BLUE. Hence there are an infinite number of RED edges coming out of  $x_1$ , or there are an infinite number of BLUE edges coming out of  $x_1$  (or both). Let  $c_1$  be a color such that  $x_1$  has an infinite number of edges coming out of it that are colored  $c_1$ . Let  $V_1$  be the set of vertices  $v$  such that  $\text{COL}(v, x_1) = c_1$ . Then keep iterating this process.

We now describe it formally.

$$V_0 = \mathbb{N}$$

$$x_1 = 1$$

$$\begin{aligned} c_1 &= \text{RED} \text{ if } |\{v \in V_0 \mid \text{COL}(v, x_1) = \text{RED}\}| \text{ is infinite} \\ &= \text{BLUE} \text{ otherwise} \end{aligned}$$

$$V_1 = \{v \in V_0 \mid \text{COL}(v, x_1) = c_1\} \text{ (note that } |V_1| \text{ is infinite)}$$

Let  $i \geq 2$ , and assume that  $V_{i-1}$  is defined. We define  $x_i$ ,  $c_i$ , and  $V_i$ :

$$\begin{aligned} x_i &= \text{the least number in } V_{i-1} \\ c_i &= \text{RED if } |\{v \in V_{i-1} \mid \text{COL}(v, x_i) = \text{RED}\}| \text{ is infinite} \\ &= \text{BLUE otherwise} \\ V_i &= \{v \in V_{i-1} \mid \text{COL}(v, x_i) = c_i\} \text{ (note that } |V_i| \text{ is infinite)} \end{aligned}$$

(NOTE- look at the step where we define  $c_i$ . We are using the fact that if you 2-color  $\mathbb{N}$  you are guaranteed some color appears infinitely often; we are using the 1-hypergraph Ramsey Theorem. Later when we prove Ramsey on 3-hypergraphs we will use Ramsey on 2-hypergraphs.)

How long can this sequence go on for? Well,  $x_i$  can be defined if  $V_{i-1}$  is nonempty. We can show by induction that, for every  $i$ ,  $V_i$  is infinite. Hence the sequence

$$x_1, x_2, \dots$$

is infinite.

Consider the infinite sequence

$$c_1, c_2, \dots$$

Each of the colors in this sequence is either RED or BLUE. Hence there must be an infinite sequence  $i_1, i_2, \dots$  such that  $i_1 < i_2 < \dots$  and

$$c_{i_1} = c_{i_2} = \dots$$

Denote this color by  $c$ , and consider the vertices

$$H = \{x_{i_1}, x_{i_2}, \dots\}$$

We leave it to the reader to show that  $H$  is homog. ■

**Exercise 1** Show that, for all  $c \geq 3$ , for every  $c$ -coloring of the edges of  $K_{\mathbb{N}}$ , there is a an infinite homog set.

Questions to ponder:

1. Is there a finite version?
2. What if we allow an infinite number of colors?
3. Computational and Complexity issues.

## 6 Ramsey's Theorem for 3-Hypergraphs: First Proof

**Theorem 6.1** *For all  $\text{COL}: \binom{\mathbb{N}}{3} \rightarrow [2]$  there exists an infinite 3-homog set.*

**Proof:**

**CONSTRUCTION**

**PART ONE**

$$V_0 = \mathbb{N}$$

$$x_0 = 1.$$

Assume  $x_1, \dots, x_{i-1}$  defined,  $V_{i-1}$  defined and infinite.

$$x_i = \text{the least number in } V_{i-1}$$

$$V_i = V_{i-1} - \{x_i\} \text{ (Will change later without changing name.)}$$

$$\text{COL}^*(x, y) = \text{COL}(x_i, x, y) \text{ for all } \{x, y\} \in \binom{V_i}{2}$$

$$V_i = \text{an infinite 2-homogeneous set for } \text{COL}^*$$

$$c_i = \text{the color of } V_i$$

KEY: for all  $y, z \in V_i$ ,  $\text{COL}(x_i, y, z) = c_i$ .

**END OF PART ONE**

**PART TWO**

We have vertices

$$x_1, x_2, \dots,$$

and associated colors

$$c_1, c_2, \dots,$$

There are only two colors, hence, by the 1-homog Ramsey Theorem there exists  $i_1, i_2, \dots$ , such that  $i_1 < i_2 < \dots$  and

$$c_{i_1} = c_{i_2} = \dots$$

We take this color to be RED. Let

$$H = \{x_{i_1}, x_{i_2}, \dots\}.$$

We leave it to the reader to show that  $H$  is homog.

END OF PART TWO  
END OF CONSTRUCTION

■

### Exercise 2

1. Show that, for all  $c$ , for all  $c$ -coloring of  $K_{\mathbb{N}}^3$  there exists an infinite 3-homog set.
2. State and prove a theorem about  $c$ -coloring  $\binom{\mathbb{N}}{a}$ .
3. What if we allow an infinite number of colors?

## 7 Ramsey's Theorem for 3-Hypergraphs: Second Proof

In the last section the proof went as follows:

- Color a *node* by using 2-hypergraph Ramsey. This is done  $\omega$  times.
- After the nodes are colored we use 1-hypergraph. This is done once.

We given an alternative proof where:

- Color an *edge* by using 1-hypergraph Ramsey This is done  $\omega$  times.
- After *all* the edges of an infinite complete graph are colored we use 2-hypergraph Ramsey. This is done once.

We now proceed formally.

**Theorem 7.1** *For all COL:  $\binom{\mathbb{N}}{3} \rightarrow [2]$  there exists an infinite 3-homog set.*

**Proof:**

Let COL be a 2-coloring of  $\binom{\mathbb{N}}{3}$ . We define a sequence of vertices,

$$x_1, x_2, \dots,$$

Here is the intuition: Let  $x_1 = 1$  and  $x_2 = 2$ . The vertices  $x_1, x_2$  induces the following coloring of  $\{3, 4, \dots\}$ .

$$\text{COL}^*(y) = \text{COL}(x_1, x_2, y).$$

Let  $V_1$  be an infinite 1-homogeneous for  $\text{COL}^*$ . Let  $\text{COL}^{**}(x_1, x_2)$  be the color of  $V_1$ . Let  $x_3$  be the least vertex left (bigger than  $x_2$ ).

The number  $x_3$  induces *two* colorings of  $V_1 - \{x_3\}$ :

$$(\forall y \in V_1 - \{x_3\})[\text{COL}_1^*(y) = \text{COL}(x_1, x_3, y)]$$

$$(\forall y \in V_1 - \{x_3\})[\text{COL}_2^*(y) = \text{COL}(x_2, x_3, y)]$$

Let  $V_2$  be an infinite 1-homogeneous for  $\text{COL}_1^*$ . Let  $\text{COL}^{**}(x_1, x_3)$  be the color of  $V_2$ . Restrict  $\text{COL}_2^*$  to elements of  $V_2$ , though still call it  $\text{COL}_2^*$ . We reuse the variable name  $V_2$  to be an infinite 1-homogeneous for  $\text{COL}_2^*$ . Let  $\text{COL}^{**}(x_1, x_3)$  be the color of  $V_2$ . Let  $x_4$  be the least element of  $V_2$ . Repeat the process.

We describe the construction formally.

## CONSTRUCTION PART ONE:

$$\begin{aligned} x_1 &= 1 \\ V_1 &= \mathbf{N} - \{x_1\} \end{aligned}$$

Let  $i \geq 2$ . Assume that  $x_1, \dots, x_{i-1}, V_{i-1}$ , and  $\text{COL}^{**} : \binom{\{x_1, \dots, x_{i-1}\}}{2} \rightarrow \{\text{RED}, \text{BLUE}\}$  are defined.

$$\begin{aligned} x_i &= \text{the least element of } V_{i-1} \\ V_i &= V_{i-1} - \{x_i\} \text{ (We will change this set without changing its name).} \end{aligned}$$

We define  $\text{COL}^{**}(x_1, x_i), \text{COL}^{**}(x_2, x_i), \dots, \text{COL}^{**}(x_{i-1}, x_i)$ . We will also define smaller and smaller sets  $V_i$  (not smaller by size – they are all infinite – but smaller by being subsets). We will keep the variable name  $V_i$  throughout.

For  $j = 1$  to  $i - 1$

1.  $\text{COL}_j^* : V_i \rightarrow \{\text{RED}, \text{BLUE}\}$  is defined by  $\text{COL}_j^*(y) = \text{COL}(x_j, x_i, y)$ .

2. Let  $V_i$  be redefined as an infinite 1-homogeneous set for  $\text{COL}^*$ . Note that  $V_i$  is still infinite.
3.  $\text{COL}^{**}(x_j, x_i)$  is the color of  $V_i$ .

**END OF PART ONE**

**PART TWO:**

From PART ONE we have a set of vertices  $X$

$$X = \{x_1, x_2, \dots\}$$

and a 2-coloring  $\text{COL}^{**}$  of  $\binom{X}{2}$ . By the 2-hypergraph Ramsey Theorem there exists an infinite homog (with respect to  $\text{COL}^{**}$ ) set

$$H = \{y_1, y_2, \dots\}$$

Assume that the homog color is  $R$ . Then for  $i < j < k$

$$\text{COL}(y_i, y_j, y_k) = \text{COL}^{**}(y_i, y_j) = R$$

So  $H$  is homog for  $\text{COL}$ . **END OF PART TWO**  
**END OF CONSTRUCTION ■**