

If L is ANY set then $SUBSEQ(L)$ is Regular
Exposition by William Gasarch (gasarch@cs.umd.edu)

1 Introduction

Definition 1.1 Let Σ be a finite alphabet.

1. Let $w \in \Sigma^*$. $SUBSEQ(w)$ is the set of all strings you get by replacing some of the symbols in w with the empty string.
2. Let $L \subseteq \Sigma^*$. $SUBSEQ(L) = \bigcup_{w \in L} SUBSEQ(w)$.

The following are easy to show:

1. If L is regular than $SUBSEQ(L)$ is regular.
2. If L is context free than $SUBSEQ(L)$ is context free.
3. If L is c.e. than $SUBSEQ(L)$ is c.e.

Note that one of the obvious suspects is missing. Is the following true:

If L is decidable then $SUBSEQ(L)$ is decidable.

We will show something far stronger. We will show that

If L is ANY subset of Σ^ WHATSOEVER then $SUBSEQ(L)$ is regular.*

Higman [2] first proved this theorem. His proof is the one we give here; however, he used different terminology.

The proofs that if L is regular (context free, c.e.) then $SUBSEQ(L)$ is regular (context free, c.e.) are constructive. That is, given the DFA (CFG, TM) for L you could produce the DFA (CFG, TM) for $SUBSEQ(L)$. (In the case of c.e. you are given M such that $L = DOM(M)$ and you can produce a TM M' such that $SUBSEQ(L) = DOM(M')$). The proof that if L is any language whatsoever then $SUBSEQ(L)$ is regular will be nonconstructive. We will discuss this later.

Definition 1.2 A set together with an ordering (X, \preceq) is a *well quasi ordering* (wqo) if for any sequence x_1, x_2, \dots there exists i, j such that $i < j$ and $x_i \preceq x_j$. We call this i, j an *uptick*

Note 1.3 If (X, \preceq) is a wqo then its both well founded and has no infinite antichains.

Lemma 1.4 *Let (X, \preceq) be a wqo. For any sequence x_1, x_2, \dots there exists an infinite ascending subsequence.*

Proof: Let x_1, x_2, \dots , be an infinite sequence. Define the following coloring:
 $COL(i, j) =$

- UP if $x_i \preceq x_j$.
- DOWN if $x_j \prec x_i$.
- INC if x_i and x_j are incomparable.

By Ramsey's theorem there is either an infinite homog UP-set, an infinite homog DOWN-set or an infinite homog INC-set. We show the last two cannot occur.

If there is an infinite homog DOWN-set then take that infinite subsequence. That subsequence violates the definition of well quasi ordering.

If there is an infinite homog INC-set then take that infinite subsequence. That subsequence violates the definition of well quasi ordering. ■

We now redefine wqo.

Definition 1.5 A set together with an ordering (X, \preceq) . is a *well quasi ordering* (wqo) if one of the following equivalent conditions holds.

- For any sequence x_1, x_2, \dots there exists i, j such that $i < j$ and $x_i \preceq x_j$.
- For any sequence x_1, x_2, \dots there exists an *infinite* ascending subsequence.

Definition 1.6 If (X, \preceq_1) and (Y, \preceq_2) are wqo then we define \preceq on $X \times Y$ as $(x, y) \preceq (x', y')$ if $x \preceq_1 y'$ and $x' \preceq_2 y$.

Lemma 1.7 If (X, \preceq_1) and (Y, \preceq_2) are wqo then $(X \times Y, \preceq)$ is a wqo (\preceq defined as in the above definition).

Proof: Let $(x_1, y_1), (x_2, y_2), (x_3, y_3), \dots$ be an infinite sequence of elements from $A \times B$.

Define the following coloring:

$COL(i, j) =$

- UP-UP if $x_i \preceq x_j$ and $y_i \preceq y_j$.
- UP-DOWN if $x_i \preceq x_j$ and $y_j \preceq y_i$.
- UP-INC if $x_i \preceq x_j$ and y_j, y_i are incomparable.
- DOWN-UP, DOWN-DOWN, DOWN-INC, INC-UP, INC-DOWN, INC-INC are defined similarly.

By Ramsey's theorem there is a homog set in one of those colors. If the color has a DOWN in it then there is an infinite descending sequence within either x_1, x_2, \dots , or y_1, y_2, \dots which violates either X or Y being a wqo. If the color has an INC in it then there is an infinite antichain within either x_1, x_2, \dots , or y_1, y_2, \dots which violates either X or Y being a wqo. Hence the color must be UP-UP. This shows that there is an infinite ascending sequence. ■

2 Subsets of Well Quasi Orders that are Closed Downward

Lemma 2.1 *Let (X, \preceq) be a countable wqo and let $Y \subseteq X$. Assume that Y is closed downward under \preceq . Then there exists a finite set of elements $\{z_1, \dots, z_k\} \subseteq X - Y$ such that*

$$y \in Y \text{ iff } (\forall i)[z_i \not\preceq y].$$

(The set $\{z_1, \dots, z_k\}$ is called an obstruction set.)

Proof: Let OBS be the set of elements z such that

1. $z \notin Y$.
2. Every $y \preceq z$ is in Y .

Claim 1: OBS is finite

Proof: We first show that every $z, z' \in OBS$ are incomparable. Assume, by way of contradiction, that $z \preceq z'$. Then $z \in Y$ by part 2 of the definition of OBS . But if $z \in Y$ then $z \notin OBS$. Contradiction.

Assume that OBS is infinite. Then the elements of OBS (in any order) form an infinite anti-chain. This violates the property of \preceq being a wqo. Contradiction.

End of Proof

Let $OBS = \{z_1, z_2, \dots\}$. The order I put the elements in is arbitrary.

Claim 2: For all y :

$$y \in Y \text{ iff } (\forall i)[z_i \not\preceq y].$$

Proof of Claim 2:

We prove the contrapositive

$$y \notin Y \text{ iff } (\exists i)[z_i \preceq y].$$

Assume $y \notin Y$. If $y \in OBS$ then we are done. If $y \notin OBS$ then, by the definition of OBS there must be some z such that $z \notin Y$ and $z \prec y$. If $z \in OBS$ then we are done. If not then repeat the process with z . The process cannot go on forever since then we would have an infinite descending sequence, violating the wqo property. Hence, after a finite number of steps, we arrive at an element of OBS . Therefore there is a $z \in OBS$ with $z \preceq y$.

Assume $(\exists i)[z_i \preceq y]$. Since Y is closed downward under \preceq and $z_i \notin Y$, this implies that $y \notin Y$.

■

3 $(\Sigma^*, \preceq_{\text{subseq}})$ is a Well Quasi Ordering

Definition 3.1 *The subsequence order, which we denote \preceq_{subseq} , is defined as $x \preceq_{\text{subseq}} y$ if x is a subsequence of y .*

IDEA: We will show that $(\Sigma^*, \preceq_{\text{subseq}})$ is a wqo. Note that if $A \subseteq \Sigma^*$ then $SUBSEQ(A)$ is closed under \preceq_{subseq} . Hence by the Lemma 2.1 there exists strings z_1, \dots, z_n such that

$$x \in SUBSEQ(A) \text{ iff } (\forall i)[z_i \not\preceq x]$$

For fixed z the set $\{x \mid z \not\preceq x\}$ is regular. Hence $SUBSEQ(A)$ is the intersection of a finite number of regular sets and is hence regular.

Theorem 3.2 (Σ^*, \preceq) is a wqo.

Proof: Assume not. Then there exists (perhaps many) sequences x_1, x_2, \dots such that for all $i < j$, $x_i \not\preceq x_j$. We call such these *bad sequences*.

Look at ALL of the bad sequences. Look at ALL of the first elements of those bad sequences. Let y_1 be the *shortest* such element (if there is a tie then pick one of them arbitrarily).

Assume that y_1, y_2, \dots, y_n have been picked. Look at ALL of the bad sequences that begin y_1, \dots, y_n (there will be at least one). Look at ALL of the $n + 1$ st elements of those sequences. Let y_{n+1} be the shortest such element (if there is a tie then pick one of them arbitrarily). We have a sequence

$$y_1, y_2, \dots$$

This is referred to as a *minimal bad sequence*.

Let $y_i = y'_i \sigma_i$ where $\sigma_i \in \Sigma$. (note that none of the y_i are empty since if they were they would not be part of any bad sequence).

Let $Y = \{y'_1, y'_2, \dots\}$.

Claim: Y is a wqo.

Proof of Claim:

Assume not. Then there is a bad sequence $y'_{k_1}, y'_{k_2}, \dots$. We know that $y_{k_i} = y'_{k_i} \sigma_{k_i}$. Lets say the bad sequence is

$$y'_{84}, y'_{12}, y'_4, y'_{1001}, y'_{32}, \dots \text{ (no pattern is intended).}$$

Lets say that y'_1, y'_2, y'_3 never appear. So y'_4 is the least indexed element. We will remove all the elements before y'_4 . Hence we can assume that the sequence starts with y'_4 .

More generally, we will start the sequence at the least indexed element. We just assume this, so we assume that $k_1 \leq \{k_2, k_3, \dots\}$. Consider the following sequence:

$$y_1, y_2, \dots, y_{k_1-1}, y'_{k_1}, y'_{k_2}, \dots$$

We show this is a BAD sequence.

There cannot be an $i < j \leq k_1 - 1$ such that $y_i \preceq y_j$ since that would mean that y_1, y_2, \dots is not a bad sequence.

There cannot be an $i < j$ with $y'_{k_i} \preceq y'_{k_j}$ since that would mean that $y'_{k_1}, y'_{k_2}, \dots$ is not a bad sequence.

And now for the interesting case. There cannot be an $i \leq k_1 - 1$ and a k_j such that $y_i \preceq y'_{k_j}$. If we had this then we would have

$$y_i \preceq y'_{k_j} \preceq y'_{k_j} \sigma_{k_j} = y_{k_j}.$$

But we made sure that $i < k_j$, so this would imply that y_1, y_2, \dots is not a bad sequence.

OKAY, so this is a bad sequence. So what? Well look— its a bad sequence that begins $y_1, y_2, \dots, y_{k_1-1}$ but its k_1 th element is y'_{k_1} which is SHORTER than y_{k_1} . This contradicts y_1, y_2, \dots , being a MINIMAL bad sequence.

End of Proof of Claim

So we know that Y is a wqo. We also know that Σ with any ordering is a wqo. By Lemma 1.7 $Y \times \Sigma$ is a wqo.

Look at the sequence

$$(y'_1, \sigma_1), (y'_2, \sigma_2), \dots$$

where $y_i = y'_i \sigma_i$.

Since Y is a wqo there exists $i < j$ such that

$$(y'_i, \sigma_i) \preceq_{\text{subseq}} (y'_j, \sigma_j), \dots$$

Clearly $y_i \preceq_{\text{subseq}} y_j$. ■

4 Main Result

Theorem 4.1 *Let Σ be a finite alphabet. If $L \subseteq \Sigma^*$ then $SUBSEQ(L)$ is regular.*

Proof: Let $L \subseteq \Sigma^*$. The set $SUBSEQ(L)$ is closed under the \preceq_{subseq} ordering. By Theorem 3.2 \preceq_{subseq} is a wqo. By Lemma 2.1 $SUBSEQ(L)$ has a finite obstruction set. From this it is easy to show that $SUBSEQ(L)$ is regular. ■

5 Nonconstructive?

One can ask: Given a DFA, CFG, P-machine, NP-machine, TM (decidable), TM (c.e.) for a language L , can one actually obtain a DFA for $SUBSEQ(L)$. For that matter, can you obtain a CFG, etc for $SUBSEQ(L)$.

	$SUBSEQ(REG)$	$SUBSEQ(CFGP)$	$SUBSEQ(P)$	$SUBSEQ(DEC)$	$SUBSEQ(C.E.)$
<i>REG</i>	<i>CON</i>	<i>CON</i>	<i>CON</i>	<i>CON</i>	<i>CON</i>
<i>CFG</i>	<i>CON</i>	<i>CON</i>	<i>CON</i>	<i>CON</i>	<i>CON</i>
<i>P</i>	<i>NONCON</i>	<i>NONCON</i>	<i>NONCON</i>	<i>NONCON</i>	<i>CON</i>
<i>NP</i>	<i>NONCON</i>	<i>NONCON</i>	<i>NONCON</i>	<i>NONCON</i>	<i>CON</i>
<i>DEC</i>	<i>NONCON</i>	<i>NONCON</i>	<i>NONCON</i>	<i>NONCON</i>	<i>CON</i>
<i>C.E.</i>	<i>NONCON</i>	<i>NONCON</i>	<i>NONCON</i>	<i>NONCON</i>	<i>CON</i>

Gasarch, Fenner, Postow [1] showed all of the NONCON results. Leeuwen [3] showed that, given a CFG for L , you can obtain a DFA for $SUBSEQ(L)$ (it also appears in [1] which is online). All the rest of the results are easy.

References

- [1] S. Fenner, W. Gasarch, and B. Postow. The complexity of finding $SUBSEQ(A)$. *Theory of Computing Systems*, 45(3):577–612, October 2009. 10.1007/s00224-008-9111-4.
- [2] A. G. Higman. Ordering by divisibility in abstract algebra. *Proceedings of the London Mathematical Society*, 3:326–336, 1952.
- [3] J. van Leeuwen. Effective constructions in well-partially-ordered free monoids. *Discrete Mathematics*, 21:237–252, 1978.