

$$R(k) \geq \Omega(k^2)$$

$$R(k) \geq \Omega(k^2)$$

Color the edges of $K_{(k-1)^2}$ as follows:

$$R(k) \geq \Omega(k^2)$$

Color the edges of $K_{(k-1)^2}$ as follows:

Partition the vertices into $k - 1$ sets of $k - 1$ each.

$$R(k) \geq \Omega(k^2)$$

Color the edges of $K_{(k-1)^2}$ as follows:

Partition the vertices into $k - 1$ sets of $k - 1$ each.

If two vertices are in the same set color then edge RED.

$$R(k) \geq \Omega(k^2)$$

Color the edges of $K_{(k-1)^2}$ as follows:

Partition the vertices into $k - 1$ sets of $k - 1$ each.

If two vertices are in the same set color then edge RED.

If two vertices are in different sets color then edge BLUE.

$$R(k) \geq \Omega(k^2)$$

Color the edges of $K_{(k-1)^2}$ as follows:

Partition the vertices into $k - 1$ sets of $k - 1$ each.

If two vertices are in the same set color then edge RED.

If two vertices are in different sets color then edge BLUE.

Exercise Show there is no homog set of size k .

$$R(k) \geq \Omega(k^2)$$

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

Color the edges of K_{n^2} as follows:

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

Color the edges of K_{n^2} as follows:

Partition the vertices into n sets of n each.

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

Color the edges of K_{n^2} as follows:

Partition the vertices into n sets of n each.

For each set use the coloring without a k -homog set.

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

Color the edges of K_{n^2} as follows:

Partition the vertices into n sets of n each.

For each set use the coloring without a k -homog set.

For colorings between sets view each set as a vertex, so have n such vertices.

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

Color the edges of K_{n^2} as follows:

Partition the vertices into n sets of n each.

For each set use the coloring without a k -homog set.

For colorings between sets view each set as a vertex, so have n such vertices.

Color all edges between set i and set j with the color of (i, j) in COL'.

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

Color the edges of K_{n^2} as follows:

Partition the vertices into n sets of n each.

For each set use the coloring without a k -homog set.

For colorings between sets view each set as a vertex, so have n such vertices.

Color all edges between set i and set j with the color of (i, j) in COL'.

Exercise Show there is no homog set of size k^2 .

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

Color the edges of K_{n^2} as follows:

Partition the vertices into n sets of n each.

For each set use the coloring without a k -homog set.

For colorings between sets view each set as a vertex, so have n such vertices.

Color all edges between set i and set j with the color of (i, j) in COL'.

Exercise Show there is no homog set of size k^2 .

Since $R(4) > 17$ get $R(4^2) > 17^2$, and for m a power of two get

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

Color the edges of K_{n^2} as follows:

Partition the vertices into n sets of n each.

For each set use the coloring without a k -homog set.

For colorings between sets view each set as a vertex, so have n such vertices.

Color all edges between set i and set j with the color of (i, j) in COL'.

Exercise Show there is no homog set of size k^2 .

Since $R(4) > 17$ get $R(4^2) > 17^2$, and for m a power of two get $R(4^m) > 17^m = (4^m)^{\log_4 17}$.

$$R(k) \geq \Omega(k^2)$$

Assume $R(k) > n$. We show $R(k^2) > n^2$.

We assume there is a coloring COL' of K_n with no k -homog set.

Color the edges of K_{n^2} as follows:

Partition the vertices into n sets of n each.

For each set use the coloring without a k -homog set.

For colorings between sets view each set as a vertex, so have n such vertices.

Color all edges between set i and set j with the color of (i, j) in COL'.

Exercise Show there is no homog set of size k^2 .

Since $R(4) > 17$ get $R(4^2) > 17^2$, and for m a power of two get $R(4^m) > 17^m = (4^m)^{\log_4 17}$.

$R(k) > k^{\log_4 17} = k^{2.043}$.