
High School Proofs for Better Bounds on the Quadratic van der Waerden Numbers

William Gasarch, Clyde Kruskal, Justin Kruskal, Zach Price

Abstract. A corollary of the polynomial van der Waerden theorem is that, for any polynomial $p(x) \in \mathbb{Z}[x]$ with constant term 0, for any $c \in \mathbb{N}$, there exists $W \in \mathbb{N}$ such that, for all c -colorings of $\{1, \dots, W\}$ there exists a, d such that a and $a + p(d)$ are the same color. The proof of the polynomial van der Waerden theorem, and even of these corollaries, is difficult and gives enormous upper bounds for W . We consider just quadratic polynomials. For $c = 2, 3$ we obtain reasonable bounds, and for $c = 4$ for some quadratics we obtain reasonable bounds, using simple methods.

1. INTRODUCTION We use the following standard definition.

Definition. Let \mathbb{Z} be the set of integers, \mathbb{N} be the set of non-negative integers, and \mathbb{N}^+ be the set of positive integers. Let $[W]$ be the set $\{1, \dots, W\}$ (where $W \in \mathbb{N}$).

In this paper we will give *High School Proofs (HS Proof)* of theorems. The term *High School Proof* is not a formal term. We use it to mean a proof that can be explained to a bright high school student. We use the term *High School Proof* since (1) the term *elementary* is ambiguous, and (2) the term *Combinatorial* is not quite right since (a) the rather difficult proof of Szemerédi's Theorem is combinatorial, and (b) the rather difficult proof of Gower's bound is mostly combinatorial.

Recall van der Waerden's Theorem [1, 2] (see also the books by Graham-Rothchild-Spencer [3] and Landman-Robertson [4]).

Theorem 1. For any $k \in \mathbb{N}$, for any $c \in \mathbb{N}$, there exists $W = W(k, c)$, such that for any c -coloring of $[W]$, there exists $a, d \in \mathbb{N}$, $d \neq 0$, such that $a, a + d, \dots, a + (k - 1)d$ are all the same color.

The original proof by van der Waerden was HS but yielded bounds on W that were not primitive recursive [3]. Shelah [5] gave a HS proof that yielded primitive recursive bounds on W . These bounds were still quite large in that they really cannot be written down nicely. Gowers [6] gave a non-HS proof that yielded bounds that can be written down:

$$W(k, c) \leq 2^{2^{c2^{k+9}}}$$

We discuss a known generalization of van der Waerden's theorem. Note that the conclusion of van der Waerden's theorem is that

$$a, a + d, a + 2d, \dots, a + (k - 1)d \text{ are the same color.}$$

Can we replace $d, 2d, \dots, (k - 1)d$ by other functions of d ? Yes. We can replace them with polynomials in $\mathbb{Z}[x]$ that have no constant term. Here is the Polynomial van der Waerden Theorem:

Theorem 2. *Let $p_1, \dots, p_k \in \mathbb{Z}[x]$ such that, for $1 \leq i \leq k$, $p_i(0) = 0$. Let $c \in \mathbb{N}$. Then there exists $W = W(p_1, \dots, p_k; c)$ such that, for any c -coloring of $[W]$, there exists $a, d \in \mathbb{N}$, $d \neq 0$, such that $a, a + p_1(d), \dots, a + p_k(d)$ are all the same color.*

For $k = 1$, this theorem was proven independently by Furstenberg [7] and Sárközy [8]. Bergelson and Leibman [9] proved the general result using ergodic methods (not a HS proof). These proofs yielded no upper bounds on $W(p_1, \dots, p_k; c)$. Walters [10] obtained a HS proof of Theorem 2, but the bounds on W were not primitive recursive. Shelah [11] gave a (non HS) proof that yielded primitive recursive bounds on W . These bounds were still quite large in that they really cannot be written down nicely. Nobody has obtained a proof that yields bounds one can write down.

Peluse [12] and Peluse and Prediville [13] proved density results that can be translated into bounds for some polynomial van der Waerden numbers.

1. Peluse and Prediville [13] showed that there exists a d such that for large n , $W(x, x^2; (\log \log n)^d) \leq n$.
2. Peluse [12] showed that if $p_1, \dots, p_m \in \mathbb{Z}[x]$ are polynomials of different degrees then there exists a constant d (which depends on p_1, \dots, p_m) such that, for large n , $W(p_1(x), \dots, p_m(x); (\log \log n)^d) \leq n$.
3. Peluse and Prediville [14] showed that there exists a d such that for large n , $W(x, x^2; (\log n)^d) \leq n$.

These proofs are not HS.

We are interested in the case of $W(ax^2 + bx; c)$ where $c = 2, 3, 4$. Furstenberg's proof showed that $W(x^2; c)$ exists; however, his proof gave no upper bounds. Sárközy's proof showed that $W(x^2; c) \leq 2^{O(c^3)}$. Pintz, Steiger, and Szemerédi [15] (see also [16] for exposition) showed that $W(x^2; c) \leq 2^{O(c^{0.0001})}$. The 0.0001 can be replaced with any smaller constant; however, in that case the constant associated with the big-O will increase. It is possible that either Sárközy's proof of $W(x^2; c) \leq 2^{O(c^3)}$ or Pintz, Steiger, and Szemerédi proof of $W(x^2; c) \leq 2^{O(c^{0.0001})}$ could be modified with a fixed value of c such as 4. That may lead to an improvement on our bound on $W(x^2; 4)$; however, such a proof would not be HS.

Harnel, Lyall, and Rice [17] showed that there exists a function $f : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{N}$ such that

$$W(ax^2 + bx; c) \leq 2^{f(a,b)c^{0.0001}}$$

(the 0.0001 can be replaced with any smaller constant; however, in that case the function f will be bigger).

Later Rice [16] showed that, for all k , there exists a function $f : \mathbb{Z}^k \rightarrow \mathbb{N}$ such that

$$W(a_k x^k + \dots + a_1 x; c) \leq 2^{f(a_k, \dots, a_1)c^{0.0001}}$$

(the 0.0001 can be replaced with any smaller constant; however, in that case the function f will be bigger). Rice (personal communication) later obtained the following more precise result: for all $\epsilon > 0$, for all $a_1, \dots, a_k \in \mathbb{Z}$, for $J = |a_1| + \dots + |a_k|$:

$$W(a_k x^k + \dots + a_1 x; c) \leq 2^{2^{2^{100k^2/\epsilon}}} + 2^{2^{(100k^4 \log J)^{100}}} + 2^{c^\epsilon}$$

In summary, the known bounds on $W(ax^2 + bx; c)$ are large.

In this paper we show that, for some $p \in \mathbb{Z}[x]$ and $c = 2, 3, 4$, one can obtain much better bounds on $W(p(x); c)$. Our proofs will be purely combinatorial and much easier than those of Walters, Shelah, and Peluse. We hasten to point out that they proved the full polynomial van der Waerden theorem whereas we only prove it in very special cases.

We will show the following.

- For all $a \in \mathbb{Z}$, $W(ax; c) = |ac| + 1$. (Theorem 5)
- For all $a, b \in \mathbb{Z}$, $W(ax^2 + bx; 2) \leq 12|a| + 6|b|$. (Theorem 6 We actually obtain more precise bounds than that depending on how a, b are related to each other. In Appendix A is a table of some exact values of $W(ax^2 + bx; 2)$.)
- For all $a \in \mathbb{N}$, $a \geq 1$, $W(ax^2 + (a - 1)x; 2) = 8a - 3$. (Theorem 8)
- $W(x^2; 3) = 29$ and, for all $a \in \mathbb{Z}$, $W(ax^2; 3) = 28a + 1$. (Theorem 10)
- For $a, b \in \mathbb{Z}$, $W(ax^2 + bx; 3) = O(ab^6 + a^5b^2)$. (Theorem 15) In Appendix B is a table of some exact values of $W(ax^2 + bx; 3)$.
- $W(x^2; 4) \leq 84,149,474,894,213,522$. (Theorem 16) In Appendix C is a table of some upper bounds on $W(ax^2 + bx; 4)$.

2. PRELIMINARIES

Definition. Let $c \in \mathbb{N}^+$ and $W \in \mathbb{N}^+$.

1. A c -coloring of $[W]$ is a mapping $[W] \rightarrow [c]$.
2. Let $p \in \mathbb{Z}[x]$. A $(p; c)$ -proper coloring of $[W]$ is a c -coloring of $[W]$ such that, for all distinct $x, y \in [W]$, if $y - x = p(d)$ for some $d \in \mathbb{Z}$, then x and y have different colors. When the context is clear, we will often write *proper c -coloring* or simply *proper coloring*.

Note that the polynomial van der Waerden number, $W = W(p(x); c)$, is the least number such that there is no $(p; c)$ -proper coloring of $[W]$.

Although we care about proper $(p; c)$ -colorings, we need a more general notion:

Definition. Let $F \subseteq \mathbb{Z}$, $c \in \mathbb{N}^+$, and $W \in \mathbb{N}^+$.

- An $(F; c)$ -proper coloring of $[W]$ is a c -coloring of $[W]$ such that, for all $x, y \in [W]$ with $y - x \in F$, x and y have different colors.
- $W = W(F; c)$ is the least number such that there is no $(F; c)$ -proper coloring of $[W]$. If no such number exists, we set $W(F; c) = \infty$.
- In the above definitions F is the set of *forbidden distances*. We will use this term for polynomial van der Waerden numbers as well. For example, if looking at $W(3x^2; c)$ the forbidden distances are $3 \times 1^2, 3 \times 2^2, \dots$

We leave the following easy lemma to the reader.

Lemma 3. Let $c \in \mathbb{N}^+$.

1. If $0 \in F$ then $W(F; c) = 1$.
2. Assume $f \in F$. Let $F' = F \cup \{-f\}$. Then $W(F; c) = W(F'; c)$.

Lemma 4. Let $p \in \mathbb{Z}[x]$, $a \in \mathbb{N}^+$, and $c \in \mathbb{N}$. Then

$$W(ap; c) = a(W(p; c) - 1) + 1.$$

Proof.

1) $W(ap; c) \leq a(W(p; c) - 1) + 1$:

Assume, by way of contradiction, that $W(ap; c) \geq a(W(p; c) - 1) + 2$. Hence there exists COL, an $(ap; c)$ -proper coloring of $[a(W(p; c) - 1) + 1]$. Note that, for all x , $ap(x)$ is a forbidden distance for COL.

We use COL to define COL', a proper $(p; c)$ -coloring of $[W(p; c)]$; which contradicts the definition of $W(p; c)$.

For $1 \leq i \leq W(p; c)$ let

$$\text{COL}'(i) = \text{COL}(a(i - 1) + 1).$$

Suppose $j - i$ is a forbidden distance for COL'. Then there exists x such that $j - i = p(x)$. Then

$$a(j - 1) + 1 - (a(i - 1) + 1) = a(j - i) = ap(x), \text{ a forbidden distance for COL.}$$

Hence $\text{COL}(a(j - 1) + 1) \neq \text{COL}(a(i - 1) + 1)$, so $\text{COL}'(j) \neq \text{COL}'(i)$. Therefore COL' is a proper $(p; c)$ -coloring of $[W(p; c)]$.

2) $W(ap; c) \geq a(W(p; c) - 1) + 1$:

To show $W(ap; c) \geq a(W(p; c) - 1) + 1$ we need to give a proper $(ap; c)$ -coloring of $[a(W(p; c) - 1)]$.

Let X be a number to be named later. Let COL' be a proper $(p; c)$ -coloring of $[X]$. The reader can easily verify that COL, defined below, is a proper $(ap; c)$ -coloring of $[aX]$.

- Color $1, \dots, a$ with $\text{COL}'(1)$.
- Color $a + 1, \dots, 2a$ with $\text{COL}'(2)$.
- \vdots
- Color $(X - 1)a + 1, \dots, Xa$ with $\text{COL}'(X)$.

Take $X = W(p; c) - 1$. By definition there exists COL', a proper $(p; c)$ -coloring of $[X]$. Hence COL is a proper $(ap; c)$ -coloring of $[aX] = [a(W(p; c) - 1)]$ which is what we need. ■

3. THE EXACT VALUE OF $W(ax; 2)$

For completeness we cover linear polynomials, for which we obtain a complete solution. The proof is very similar to the proof of Lemma 4.

Theorem 5. *Let $a \in \mathbb{Z}$ and $c \in \mathbb{N}^+$. Then*

$$W(ax; c) = |ac| + 1.$$

Proof. By Lemma 3.1 we have the case of $a = 0$. We will assume $a \geq 1$. The case where $a \leq -1$ is similar. By Lemma 3.2 we can assume that a is a forbidden distance. $W(ax; c) \leq ac + 1$:

By setting $x = 1, 2, \dots, c$ we get forbidden distances $a, 2a, \dots, ca$. So $1, a + 1, 2a + 1, \dots, ca + 1$ must all be different colors, but there are only c colors.

$W(ax; c) \geq ac + 1$:

We can properly c -color $[ca]$:

- Color $1, \dots, a$ with 1.
- Color $a + 1, \dots, 2a$ with 2.
- \vdots
- Color $(c - 1)a + 1, \dots, ca$ with c .

■

4. UPPER BOUNDS ON $W(ax^2 + bx; 2)$

Theorem 6. *Let $a, b \in \mathbb{N}$.*

1. $W(ax^2 + bx, 2) \leq 12a + 6b + 1$.
2. *If $b \geq 3a$ then $W(-ax^2 + bx, 2) \leq 6b - 12a + 1$.*
3. *If $2a \leq b \leq 3a$ then $W(-ax^2 + bx, 2) \leq 3b - 3a + 1$.*
4. *If $a \leq b \leq 2a$ then $W(-ax^2 + bx, 2) \leq 9a - 3b + 1$.*
5. *If $0 \leq b \leq a$ then $W(-ax^2 + bx, 2) \leq 12a - 6b + 1$.*
6. *One can obtain bounds for $W(ax^2 - bx, 2)$ easily since it equals $W(-ax^2 + bx, 2)$.*
7. *One can obtain bounds for $W(-ax^2 - bx, 2)$ easily since it equals $W(ax^2 + bx, 2)$.*
8. *For all $a, b \in \mathbb{Z}$, $W(ax^2 + bx; 2) \leq 12|a| + 6|b|$. (This follows from the other parts.)*

Proof. If $a = 0$ then Theorem 5 yields the results. Hence we assume $a \geq 1$.

We will need the following claim.

Claim: If COL is a 2-coloring of an initial segment of \mathbb{N}^+ . Let d be a forbidden distance for COL. then $3d$ is a forbidden distance for COL.

Proof of Claim: Let y and $y + 3d$ be in the domain of COL. Hence $y + d, y + 2d$ are also in the domain of COL. We can assume $\text{COL}(y) = R$. Then

$$\text{COL}(y) = R \implies \text{COL}(y + d) = B \implies \text{COL}(y + 2d) = R \implies \text{COL}(y + 3d) = B.$$

End of Proof of Claim

1) $W(ax^2 + bx; 2)$. By plugging in $x = 1, 2, 3$ we find forbidden distances:

$$\{a + b, 4a + 2b, 9a + 3b\}.$$

By the Claim the following are forbidden distances:

$$\{3a + 3b, 3(4a + 2b), 9a + 3b\} = \{3a + 3b, 12a + 6b, 9a + 3b\}.$$

Assume there is a proper $W(x^2; 2)$ -coloring of $[12a + 6b + 1]$. We will get a contradiction. We can assume that $\text{COL}(1) = R$. Note that

$$\text{COL}(1) = R \implies \text{COL}(1 + (3a + 3b)) = B \implies \text{COL}(1 + (3a + 3b) + (9a + 3b)) = R.$$

We simplify to obtain $\text{COL}(12a + 6b + 1) = R$.

$$\text{COL}(12a + 6b + 1) = R \implies \text{COL}(12a + 6b + 1 - (12a + 6b)) = B.$$

We simplify to obtain $\text{COL}(1) = B$ which is a contradiction.

The key to the last proof was that

- $(3a + 3b) + (9a + 3b) - (12a + 6b) = 0$.
- COL is defined on $(3a + 3b) + (9a + 3b) + 1 = 12a + 6b + 1$.

For all later proofs we just give nonnegative forbidden distances d_1, d_2, d_3 such that $d_1 + d_2 - d_3 = 0$, and conclude that the bound is $d_1 + d_2 + 1$. We abbreviate *Forbidden Distances* by FD.

We now consider $W(-ax^2 + bx; 2)$:

2) $b \geq 3a$. $\{3b - 3a, 6b - 12a, 3b - 9a\}$ are FDs. Hence $(3b - 3a) + (3b - 9a) - (6b - 12a) = 0$ is a FD.

3) $2a \leq b \leq 3a$. $\{3b - 3a, 6b - 12a, 9a - 3b\}$ are FDs. Hence $(6b - 12a) + (9a - 3b) - (3b - 3a) = 0$ is a FD.

4) $a \leq b \leq 2a$. $\{3b - 3a, 12a - 6b, 9a - 3b\}$ are FDs. Hence $(3b - 3a) + (12a - 6b) - (9a - 3b) = 0$ is a FD.

5) $0 \leq b \leq a$. $\{3a - 3b, 12a - 6b, 9a - 3b\}$ are FDs. Hence $(3a - 3b) + (9a - 3b) - (12a - 6b) = 0$ is a FD.

■

Corollary 7. For all $a, b \in \mathbb{Z}$, $W(ax^2 + bx; 2) \leq 12|a| + 6|b| + 1$.

The bounds on $W(ax^2 + bx; 2)$ (and the others) from Theorem 6 hold for all a, b ; however, for particular a, b better bounds can often be found. We give a class of examples.

Theorem 8. Let $a \in \mathbb{N}$ with $a \geq 1$. Then $W(ax^2 + (a - 1)x; 2) = 8a - 3$.

Proof.

1) $W(ax^2 + (a - 1)x; 2) \leq 8a - 3$.

Let $\text{COL}: [8a - 3] \rightarrow \{R, B\}$.

By plugging in $x = 1, 2$ we find forbidden distances: $\{2a - 1, 6a - 2\}$. Since $2a - 1$ is a forbidden distance, so is $3(2a - 1) = 6a - 3$. We will use forbidden distances $\{6a - 3, 6a - 2\}$.

Let $y \leq 2a - 1$. Assume $\text{COL}(y) = R$. Then

$$\text{COL}(y) = R \implies \text{COL}(y + (6a - 2)) = B \implies \text{COL}(y + (6a - 2) - (6a - 3)) = R.$$

Since $y + (6a - 2) - (6a - 3) = y + 1$ we have the following which is the keys fact needed for our proof:

$$y \leq 2a - 1 \implies \text{COL}(y) = \text{COL}(y + 1).$$

(We needed $y \leq 2a - 1$ since we needed $y + (6a - 2) \leq 8a - 3$ so that $y + (6a - 2)$ is in the domain of COL.)

Assume $\text{COL}(1) = R$. Then by applying the above we get $\text{COL}(2) = R, \dots, \text{COL}(2a) = R$. However, since $\text{COL}(1) = R$ and $2a - 1$ is a forbidden distance, $\text{COL}(2a) = B$. This is a contradiction.

2) $W(ax^2 + (a - 1)x; 2) \geq 8a - 3$.

We give a coloring COL of $[8a - 4]$ such that, for all $x, y \in [8a - 4]$ with $|x - y| \in \{2a - 1, 6a - 2\}$, $\text{COL}(x) \neq \text{COL}(y)$. All other forbidden distances are larger than $8a - 4$ and hence irrelevant.

Here is the coloring:

1. For $1 \leq y \leq 2a - 1$, $\text{COL}(y) = R$.
2. For $2a \leq y \leq 4a - 2$, $\text{COL}(y) = B$.
3. For $4a - 1 \leq y \leq 6a - 3$, $\text{COL}(y) = R$.
4. For $6a - 2 \leq y \leq 8a - 4$, $\text{COL}(y) = B$.


The reader can verify that this coloring suffices. ■

In Appendix A is a table of some exact values of $W(ax^2 + bx; 2)$.

5. $W(ax^2; 3) = 28a + 1$

In this section we will show that $W(x^2; 3) = 29$ and then $W(ax^2; 3) = 28a + 1$. We first show a weaker theorem which will be a good warm-up to our work on 4-colorings in Section 7.

Theorem 9. $W(x^2; 3) \leq 68$.



Proof.

Figure 1. In any proper $(x^2, 3)$ -coloring, $\text{COL}(10) = \text{COL}(17)$

Assume, by way of contradiction, that COL is an $(x^2; 3)$ -proper coloring of [68]. Figure 1 shows some constraints on COL: COL restricted to $\{10, 1, 26, 17\}$ has to be

a proper 3-coloring of the graph (no vertices that have an edge between them are the same color).

We can assume $\text{COL}(10) = R$ and $\text{COL}(1) = B$. By looking at Figure 1 we see that $\text{COL}(26) \notin \{R, B\}$, hence $\text{COL}(26) = G$. Again by looking at Figure 1 we have that $\text{COL}(17) \notin \{B, G\}$, hence $\text{COL}(17) = R$.

Note that we have shown that $\text{COL}(10) = \text{COL}(17)$. More generally we have shown that, for all x , $\text{COL}(x) = \text{COL}(x + 7)$. Not quite. We need that (1) $x - 9$ is in the domain of the coloring, so $x \geq 10$, and $x + 16$ is in the domain of the coloring, so $x \leq 52$. To restate: if $10 \leq x \leq 52$ then $\text{COL}(x) = \text{COL}(x + 7)$.

$$\text{COL}(10) = \text{COL}(17) = \text{COL}(24) = \text{COL}(31) = \text{COL}(38) = \text{COL}(45) = \text{COL}(52) = \text{COL}(59).$$

Since $59 - 10 = 49 = 7^2$, this contradicts COL being an $(x^2; 3)$ -proper coloring. ■

The bound in Theorem 9 is not tight. The next theorem gives a tight bound. The following theorem was proven by Matthew Jordan and William Gasarch.

Theorem 10.

1. $W(x^2; 3) = 29$.
2. For all $a \in \mathbb{Z}$, $W(ax^2; 3) = 28a + 1$. This follows from Part 1 and Lemma 4.

Proof. $W(x^2; 3) \leq 29$: Assume, by way of contradiction, that there exists COL , a proper $(x^2, 3)$ -coloring of $\{1, \dots, 29\}$. Figure 1 shows some constraints on COL : COL restricted to $\{1, 10, 17, 26\}$ has to be a proper 3-coloring of the graph (no vertices that have an edge between them are the same color).

By Figure 1, $\text{COL}(10) = \text{COL}(17)$. By similar reasoning one can show that

$$(\forall x)[10 \leq x \leq 13 \implies \text{COL}(x) = \text{COL}(x + 7)].$$

We refer to this fact as FORCE-SEVEN since the value of $\text{COL}(x)$ forces the value of $\text{COL}(x + 7)$.

We can assume $\text{COL}(10) = R$. Since $11 - 10 = 1^2$ we know that $\text{COL}(10) \neq \text{COL}(11)$, so we can assume $\text{COL}(11) = B$.

17: By FORCE-SEVEN $\text{COL}(17) = \text{COL}(10) = R$

18: By FORCE-SEVEN $\text{COL}(18) = \text{COL}(11) = B$.

| | | | | | | | | | | |
|----|----|----|----|----|----|----|----|----|----|----|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| R | B | | | | | | R | B | | |

19: Since $\text{COL}(10) = R$, $\text{COL}(19) = \text{COL}(10 + 3^2) \neq R$. Since $\text{COL}(18) = B$, $\text{COL}(19) = \text{COL}(18 + 1^2) \neq B$. Hence $\text{COL}(19) \notin \{R, B\}$, so $\text{COL}(19) = G$.

12: By FORCE-SEVEN $\text{COL}(12) = \text{COL}(19) = G$.

| | | | | | | | | | | |
|----|----|----|----|----|----|----|----|----|----|----|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| R | B | G | | | | | R | B | G | |

20: Since $\text{COL}(11) = B$ and $\text{COL}(19) = G$, $\text{COL}(20) = R$.

13: By FORCE-SEVEN $\text{COL}(13) = \text{COL}(20) = R$.

| | | | | | | | | | | |
|----------|----------|----------|----------|----|----|----|----------|----------|----------|----------|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| <i>R</i> | <i>B</i> | <i>G</i> | <i>R</i> | | | | <i>R</i> | <i>B</i> | <i>G</i> | <i>R</i> |

Now we have that $\text{COL}(17) = \text{COL}(13) = R$. But $17 - 13 = 2^2$. This is a contradiction.

$W(x^2, 3) \geq 29$:

We present a proper 3-coloring:

| | | | | | | | | | | | | | |
|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
| <i>B</i> | <i>G</i> | <i>R</i> | <i>G</i> | <i>R</i> | <i>B</i> | <i>G</i> | <i>B</i> | <i>G</i> | <i>R</i> | <i>B</i> | <i>G</i> | <i>B</i> | <i>G</i> |

| | | | | | | | | | | | | | |
|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
| <i>R</i> | <i>B</i> | <i>R</i> | <i>B</i> | <i>G</i> | <i>R</i> | <i>B</i> | <i>R</i> | <i>B</i> | <i>G</i> | <i>R</i> | <i>G</i> | <i>R</i> | <i>B</i> |

We can assume $\text{COL}(10) = R$. Since $11 - 10 = 1^2$ we know that $\text{COL}(10) \neq \text{COL}(11)$, so we can assume $\text{COL}(11) = B$.

17: By FORCE-SEVEN $\text{COL}(17) = \text{COL}(10) = R$

18: By FORCE-SEVEN $\text{COL}(18) = \text{COL}(11) = B$.

| | | | | | | | | | | |
|----------|----------|----|----|----|----|----|----------|----------|----|----|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| <i>R</i> | <i>B</i> | | | | | | <i>R</i> | <i>B</i> | | |

19: Since $\text{COL}(10) = R$, $\text{COL}(19) = \text{COL}(10 + 3^2) \neq R$. Since $\text{COL}(18) = B$, $\text{COL}(19) = \text{COL}(18 + 1^2) \neq B$. Hence $\text{COL}(19) \notin \{R, B\}$, so $\text{COL}(19) = G$.

12: By FORCE-SEVEN $\text{COL}(12) = \text{COL}(19) = G$.

| | | | | | | | | | | |
|----------|----------|----------|----|----|----|----|----------|----------|----------|----|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| <i>R</i> | <i>B</i> | <i>G</i> | | | | | <i>R</i> | <i>B</i> | <i>G</i> | |

20: Since $\text{COL}(11) = B$ and $\text{COL}(19) = G$, $\text{COL}(20) = R$.

13: By FORCE-SEVEN $\text{COL}(13) = \text{COL}(20) = R$.

| | | | | | | | | | | |
|----------|----------|----------|----------|----|----|----|----------|----------|----------|----------|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| <i>R</i> | <i>B</i> | <i>G</i> | <i>R</i> | | | | <i>R</i> | <i>B</i> | <i>G</i> | <i>R</i> |

Now we have that $\text{COL}(17) = \text{COL}(13) = R$. But $17 - 13 = 2^2$. This is a contradiction.

$W(x^2, 3) \geq 29$:

We present a proper 3-coloring:

| | | | | | | | | | | | | | |
|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
| <i>B</i> | <i>G</i> | <i>R</i> | <i>G</i> | <i>R</i> | <i>B</i> | <i>G</i> | <i>B</i> | <i>G</i> | <i>R</i> | <i>B</i> | <i>G</i> | <i>B</i> | <i>G</i> |

| | | | | | | | | | | | | | |
|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 |
| <i>R</i> | <i>B</i> | <i>R</i> | <i>B</i> | <i>G</i> | <i>R</i> | <i>B</i> | <i>R</i> | <i>B</i> | <i>G</i> | <i>R</i> | <i>G</i> | <i>R</i> | <i>B</i> |

■

6. UPPER BOUNDS ON $W(ax^2 + bx; 3)$

We will obtain upper bounds on $W(ax^2 + bx; 3)$ in the case where $a \in \mathbb{N}^+$ and $b \in \mathbb{Z} - \{0\}$. For $b = 0$, Theorem 10.3 yields $W(ax^2; 3) = 28a + 1$. For $a = 0$, Theorem 5 yields $W(bx; 3) = bc + 1$. For $a \leq -1$ we can use $W(ax^2 + b; 3) = W(-ax^2 - b; 3)$.

Definition.

- (a) A coloring of $[w]$ has *repeat distance* r if x and $x + r$ have the same color, for all $1 \leq x \leq w - r$.
- (b) A coloring of $[w]$ has *repeat distance* r *under one-sided boundary condition* b if x and $x + r$ have the same color, for all $1 \leq x \leq w - r - b$.
- (c) A coloring of $[w]$ has *repeat distance* r *under two-sided boundary condition* b if x and $x + r$ have the same color, for all $b + 1 \leq x \leq w - r - b$.

Lemma 11. *In any 3-coloring of $[w]$ with forbidden distances $s, t, s + t$, where $0 < s < t$:*

- (a) $2s + t$ is a repeat distance.
- (b) $t - s$ is a repeat distance under two-sided boundary condition s .
- (c) $3s$ is a repeat distance under one-sided boundary condition t .

Proof. Let $u = s + t$.

- (a) Consider a 3-coloring satisfying the conditions of the lemma. Let

$$1 \leq x \leq w - (2s + t).$$

Without loss of generality, we can assume that x is *R*. Then $x + s$ is not *R*, say *B*, and $x + u = (x + s) + t$ cannot be *R* or *B* so it must be *G*. Then $(x + s) + u = (x + u) + s$ cannot be *B* or *G* so it must be *R*. Since x and $x + u + s$ are both *R*,

$$(x + u + s) - x = u + s = 2s + t$$

is a repeat distance.

- (b) Consider a 3-coloring satisfying the conditions of the lemma. Let

$$s < x \leq w - (t - s) - s.$$

Without loss of generality, we can assume that x is *R*. Then $x - s$ is not *R*, say *B*, and $(x - s) + u = x + t$ cannot be *R* or *B* so it must be *G*. Then $(x - s) + t = (x + t) - s$ cannot be *B* or *G*, so it must be *R*. This process requires that $x - s > 0$ and $x + t \leq w$. So $(x + t - s) - x = t - s$ is a repeat distance under two-sided boundary condition s .

- (c) Take $2s + t$ from part (a) and subtract $t - s$ from part (b). The repeat distance is $(2s + t) - (t - s) = 3s$. There is a one-sided boundary of size $(t - s) + s = t$ from one side of part (b). ■

Lemma 12. Assume $[w]$ has a proper 3-coloring COL where s is a forbidden distance and r is repeat distance under either one-sided or two-sided boundary condition b . If $r|s$ then

$$w \leq s + 2b + 1.$$

Proof. We prove the lemma for 2-sided boundary condition. The case of 1-sided boundary condition follows immediately.

Assume, by way of contradiction, that $w \geq s + 2b + 2$. By (a) the definition of repeat distance under two sided boundary condition, (b) $r|s$, and (c) $b + 1 + (\frac{s}{r} - 1)r \leq w - r - b$ (this is equivalent to $w \geq 2b + s + 1$ which follows from $w \geq 2b + s + 2$) we have:

$$\begin{aligned} \text{COL}(b + 1) &= \text{COL}(b + 1 + r) = \text{COL}(b + 1 + 2r) = \dots \\ &= \text{COL}\left(b + 1 + \left(\frac{s}{r} - 1\right)r\right) = \text{COL}\left(b + 1 + \left(\frac{s}{r}\right)r\right) = \text{COL}(b + 1 + s). \end{aligned}$$

But s is a forbidden distance so $b + 1$ and $s + b + 1$ cannot have the same color. Contradiction. ■

We use Lemma 12 to get upper bounds on several quadratic van der Waerden numbers. For one of them we have an exact value.

Theorem 13.

1. For $a, b > 0$ and $a|b$, $W(ax^2 + bx; 3) \leq \frac{72b^2}{a} + 1$.
2. $W(x^2 + x; 3) = 73$.

Proof.

1) Let $p(x) = ax^2 + bx$. Let

$$x = \frac{5b}{a}, \quad y = \frac{6b}{a}, \quad z = \frac{8b}{a}.$$

Then

$$p(x) = \frac{30b^2}{a}, \quad p(y) = \frac{42b^2}{a}, \quad p(z) = \frac{72b^2}{a}.$$

Since $p(x) + p(y) = p(z)$, by Lemma 11b, $p(y) - p(x) = \frac{12b^2}{a}$ is a repeat distance under two-sided boundary condition $\frac{30b^2}{a}$. But $p(\frac{3b}{a}) = \frac{12b^2}{a}$ is a forbidden distance.

Thus, by Lemma 12, $W(ax^2 + bx; 3) \leq \frac{12b^2}{a} + 2 \cdot \frac{30b^2}{a} + 1 = \frac{72b^2}{a} + 1$.

2) By Part 1 $W(x^2 + x; 3) \leq 73$. We show $W(x^2 + x; 3) \geq 73$ by giving a $(x^2 + x; 3)$ -proper coloring of $\{1, \dots, 72\}$.

| | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|----|----|----|----|----|----|----|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 |
| R | R | G | G | R | R | B | B | R | R | B | B | G | G | B | B | G | G |

| | | | | | | | | | | | | | | | | | |
|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 |
| R | R | G | G | R | R | B | B | R | R | B | B | G | G | B | B | G | G |

| | | | | | | | | | | | | | | | | | |
|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 |
| R | R | G | G | R | R | B | B | R | R | B | B | G | G | B | B | G | G |

| | | | | | | | | | | | | | | | | | |
|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 |
| R | R | G | G | R | R | B | B | R | R | B | B | G | G | B | B | G | G |

■

We need one more lemma before getting an upper bound for $W(p(x); 3)$ where $p(x) = ax^2 + bx$.

Lemma 14. *Let*

$$q(a, b) = (4a^3 + 4a^2 + 3a + 1)b^2 \quad \text{and} \quad r(a, b) = (4a^5 + 8a^4 + 8a^3 + 6a^2 + 3a + 1)b^2.$$

Then, for all $a \in \mathbb{N}^+$, $b \in \mathbb{Z} - \{0\}$, $\gcd(2q(a, b) + r(a, b), 3q(a, b))$ divides $18b^2$.

Proof.

The only values of (a, b) for which $q(a, b) = 0$ either involve $a \notin \mathbb{N}^+$ or $b = 0$. Hence, for the domain we are concerned with, $q(a, b) \neq 0$. The only values of (a, b) for which $r(a, b) = 0$ either involve $a \notin \mathbb{R}$, $a = -1$, or $b = 0$. Hence, for the domain we are concerned with, $r(a, b) \neq 0$. Therefore $\gcd(q(a, b), r(a, b))$ always exists. We use this implicitly.

1) We examine $\gcd(q(a, b), r(a, b))$.

Let $a \in \mathbb{N}^+$ and $b \in \mathbb{Z} - \{0\}$. Let $d_1 = \gcd(q(a, b), r(a, b))$.

The reader can verify the following equation:

$$(-20a^4 - 12a^3 - 4a^2 - 10a - 1)q(a, b) + (20a^2 - 8a + 7)r(a, b) = 6b^2.$$

Since d_1 divides the LHS, d_1 divides the RHS. Hence d_1 divides $6b^2$.

2) We examine $\gcd(2q(a, b) + r(a, b), 3q(a, b))$.

Let $a \in \mathbb{N}^+$ and $b \in \mathbb{Z} - \{0\}$. Let $d = \gcd(2q(a, b) + r(a, b), 3q(a, b))$.

The reader can verify the following equation:

$$3(2q(a, b) + r(a, b)) - 2(3q(a, b)) = 3r(a, b).$$

Since d divides the LHS, d divides the RHS, hence d divides $3r(a, b)$. Since d also divides $3q(a, b)$, d divides

$$\gcd(3r(a, b), 3q(a, b)) = 3 \gcd(r(a, b), q(a, b)) = 3d_1.$$

Since d_1 divides $6b^2$, d divides $18b^2$. ■

Theorem 15. Let $a \in \mathbb{N}^+$ and $b \in \mathbb{Z} - \{0\}$. Let $p(x) = ax^2 + bx$. Then $W(p(x); 3) = O(ab^6 + a^5b^2)$.

Proof. Let w be such that there is a proper $(p; 3)$ -coloring $\text{COL}: [w] \rightarrow [3]$. We will show that $w = O(ab^6 + a^5b^2)$.

Let

$$x_0 = (2a + 1)b, \quad y_0 = (2a^2 + 2a + 1)b, \quad z_0 = (2a^2 + 2a + 2)b.$$

Then

$$\begin{aligned} p(x_0) &= (4a^3 + 4a^2 + 3a + 1)b^2 \\ p(y_0) &= (4a^5 + 8a^4 + 8a^3 + 6a^2 + 3a + 1)b^2 \\ p(z_0) &= (4a^5 + 8a^4 + 12a^3 + 10a^2 + 6a + 2)b^2 \end{aligned}$$

Thus $p(x_0) + p(y_0) = p(z_0)$. Note that $p(x_0)$, $p(y_0)$, and $p(z_0)$ are (1) positive since $a \in \mathbb{N}^+$ and the only occurrence of b is in b^2 , and (2) forbidden distances.

- 1) By Lemma 11a, $2p(x_0) + p(y_0)$ is a repeat distance.
- 2) By Lemma 11c, $3p(x_0)$ is a repeat distance under one-sided boundary condition $p(y_0)$.
- 3) By Lemma 14 $\gcd(2p(x_0) + p(y_0), 3p(x_0)) = d \leq 18b^2$.

Claim: Let $a \in \mathbb{N}^+$ and $b \in \mathbb{Z} - \{0\}$.

1. $3p(x_0)$ does not divide $2p(x_0) + p(y_0)$.
2. $2p(x_0) + p(y_0)$ does not divide $3p(x_0)$.
3. There is a linear combination over \mathbb{Z} of $2p(x_0) + p(y_0)$ and $3p(x_0)$ that sums to d where one coefficient is < 0 and the other coefficient is > 0 .

Proof of Claim

Note that

$$\begin{aligned} 2p(x_0) + p(y_0) &= (4a^5 + 8a^4 + 16a^3 + 14a^2 + 9a + 3)b^2 \\ 3p(x_0) &= (12a^3 + 12a^2 + 9a + 3)b^2 \end{aligned}$$

- 1) For all $a \in \mathbb{N}^+$, for all $b \in \mathbb{Z} - \{0\}$, $3p(x_0)$ does not divide $2p(x_0) + p(y_0)$.
If we divide $3p(x_0)$ into $2p(x_0) + p(y_0)$ as polynomials in a, b we get the following:

$$(4a^5 + 8a^4 + 16a^3 + 14a^2 + 9a + 3)b^2 =$$

$$\left(\frac{a^2}{3} + \frac{a}{3} + \frac{3}{4}\right)(12a^3 + 12a^2 + 9a + 3)b^2 + \left(a^2 + \frac{5a}{4} + \frac{3}{4}\right)b^2.$$

Since for all $a \in \mathbb{N}^+$, $b \in \mathbb{Z} - \{0\}$, $(a^2 + \frac{5a}{4} + \frac{3}{4})b^2 \neq 0$, $3p(x_0)$ does not divide $2p(x_0) + p(y_0)$.

2) For all $a \in \mathbb{N}^+$, $b \in \mathbb{Z} - \{0\}$, $2p(x_0) + p(y_0)$ does not divide $3p(x_0)$.

For all $a \in \mathbb{N}^+$, $3p(x_0) < 2p(x_0) + p(y_0)$, hence $2p(x_0) + p(y_0)$ does not divide $3p(x_0)$.

3) Since $\gcd(2p(x_0) + p(y_0), 3p(x_0)) = d$, there is a linear combination over \mathbb{Z} of these two quantities that sums to d . Since both of these quantities are $\geq d$ one coefficient must be ≤ 0 and one must be ≥ 0 . By Part 1 and 2, neither coefficient can be 0. Hence one coefficient is < 0 and the other is > 0 .

End of Proof of Claim

By Claim (part 3) there exists $j, k \in \mathbb{N}$ such that

$$j(2p(x_0) + p(y_0)) - k(3p(x_0)) = d.$$

By starting at 1 and adding repeat distance $2p(x_0) + p(y_0)$ j times and subtracting repeat distance $3p(x_0)$ k times, we see that d is a repeat distance; however, we need to be careful about the boundary condition. By interspersing the adds and subtracts so that we subtract whenever the sum is greater than $2p(x_0) + p(y_0)$, the one-sided boundary condition is $(2p(x_0) + p(y_0)) + p(y_0) = 2(p(x_0) + p(y_0))$. Hence

4) d is a repeat distance with one-sided boundary condition

$$(2p(x_0) + p(y_0)) + p(y_0) = 2(p(x_0) + p(y_0)).$$

5) $p(db) = ad^2b^2 + b^2d = (ad + 1)db^2 = O(ad^2b^2) = O(ab^6)$ is a forbidden distance. (We use $d \leq 18b^2$.)

By Lemma 12 with $s = p(db) = (ad + 1)db^2$, $r = d$, $b = 2(p(x_0) + p(y_0))$ we get

$$w \leq s + 2b + 1 = (ad + 1)db^2 + 4(p(x_0) + p(y_0)) + 1.$$

Since $p(x_0) = O(a^3b^2)$ and $p(y_0) = O(a^5b^2)$, $4(p(x_0) + p(y_0)) + 1 = O(a^5b^2)$. Hence

$$w \leq s + 2b + 1 = (ad + 1)db^2 + 4(p(x_0) + p(y_0)) + 1 = O(ab^6 + a^5b^2).$$



In Appendix B is a table of some exact values of $W(ax^2 + bx; 3)$.

7. UPPER BOUNDS ON $W(x^2; 4)$

Recall that Figure 1 was the key to showing $W(x^2; 3) \leq 68$. We now derive parameters for a new figure that will be the key to an upper bound on $W(x^2; 4)$.

We need to find $a, b, c, d, e, f, x, y, z \in \mathbb{N}^+$ such that the following figure can be drawn:

Hence we need to find solutions in \mathbb{N}^+ to the following system of equations:




Figure 2. In any $(x^2; 4)$ -proper coloring, $\text{COL}(1) = \text{COL}(1 + w)$

$$\begin{aligned} x^2 + a^2 &= y^2 \\ x^2 + b^2 &= z^2 \\ y^2 + c^2 &= z^2 \\ x^2 + d^2 &= w \\ y^2 + e^2 &= w \\ z^2 + f^2 &= w \end{aligned}$$

The first three equations are overlapping Pythagorean triples—we have three numbers (x, y, z) whose squares have all square pairwise differences. From the first three equations one can derive the following:

$$\begin{aligned} c^2 + f^2 &= e^2 \\ b^2 + f^2 &= d^2 \\ a^2 + c^2 &= b^2 \end{aligned}$$

We give one example by deriving $c^2 + f^2 = e^2$ algebraically. From $y^2 + c^2 = z^2$ and $z^2 + f^2 = w$ we get $y^2 + c^2 = w - f^2$, and hence

$$c^2 + f^2 = w - y^2.$$

From $y^2 + e^2 = w$ we get that the left hand side is e^2 . Hence

$$c^2 + f^2 = e^2.$$

Since the first three equations are Pythagorean triples, they can be generated by using Euclid’s formula: all Pythagorean triples are of the form $(k(m^2 - n^2), k(2mn), k(m^2 + n^2))$ where $\text{gcd}(m, n) = 1$ and $m \not\equiv n \pmod{2}$. We can use the Farey sequence as an efficient algorithm to generate coprime pairs m, n . (See Routledge [18] or the Wikipedia entry on Farey sequences for the definition of Farey sequences and the algorithm.)

We used a computer program and obtained the following:

Theorem 16. $W(x^2; 4) \leq 1 + (290,085,289)^2 = 84,149,474,894,213,522$




Figure 3. In any $(x^2; 4)$ -proper coloring, $\text{COL}(1) = \text{COL}(1 + 290,085,290)$

Assume, by way of contradiction, that there exists COL, a proper $(x^2; 4)$ -coloring of $[1 + (290,085,289)^2]$. Figure 3 shows some constraints on COL: COL restricted to the numbers on the vertices has to be a proper 4-coloring of the graph (no vertices that have an edge between them are the same color).

By Figure 3 we know that

$$\text{COL}(1) = \text{COL}(1 + 290,085,289^2).$$

More generally we have shown that, for all x ,

$$\text{COL}(x) = \text{COL}(x + 290,085,289^2).$$

Hence

$$\text{COL}(1) = \text{COL}(1 + 290,085,289) = \text{COL}(1 + 2 \times 290,085,289) = \dots = \text{COL}(1 + (290,085,289)^2).$$

This contradicts COL being an $(x^2; 4)$ -proper coloring. ■

Theorem 16 gave an enormous upper bound on $W(x^2; 4)$. The proof was found by a computer program; however, it is a HS proof and human-verifiable. Four colors

seems to be at the limit of what computers can find. That is, we have been unable to use a program to find a human-verifiable proof for a bound on $W(x^2; 5)$.

Usually a HS proof gives better bounds than a proof that uses advanced mathematics. However, our HS proof of a bound on $W(x^2; 4)$ gives such a large bound that its possible a proof using more advanced mathematics would yield a better result. In particular, its possible that if the proofs of the results of Sarkozy [8], Pintz-Steiger-Szemerédi [15], or Harnel-Lyall-Rice [17] were looked at more carefully then one could obtain better bounds for $W(x^2; c)$ for some small values of c . However, these would not be HS proofs.

8. UPPER BOUNDS ON $W(Ax^2 + Bx; 4)$

To find upper bounds on $W(Ax^2 + Bx; 4)$ we have several overlapping equations of the form

$$(Ax^2 + Bx) + (Ay^2 + By) = (Az^2 + Bz).$$

We need a way to generate such triples (x, y, z) much like the generation of Pythagorean triples. First, we use the quadratic formula to express z in terms of x and y .

$$z = \frac{-B + \sqrt{4A^2(x^2 + y^2) + 4AB(x + y) + B^2}}{2A}$$

We rewrite as

$$4A^2(x^2 + y^2) + 4AB(x + y) + B^2 = (2Az + B)^2.$$

Simple algebra allows us to rewrite this as:

$$(2Ax + B)^2 + (2Ay + B)^2 = (2Az + B)^2 + B^2.$$

If $m = 2Ax + B$, $n = 2Ay + B$, and $k = (2Az + B)$ then we can rewrite this as $m^2 + n^2 = k^2 + B^2$. A parameterization of $m^2 + n^2 = k^2 + B^2$ would imply one for (x, y, z) , and luckily this equation is easier. Using the *Bramagupta-Fibonacci identity* with $bc - ad = B$, we get:

$$(ac - bd)^2 + (ad + bc)^2 = (ac + bd)^2 + B^2$$

So, with parameters a, b, c, d and some tedious algebra we get

$$x = \frac{ac - bd - B}{2A}, y = \frac{ad + bc - B}{2A}, z = \frac{ac + bd - B}{2A}$$

with constraints $bc - ad = B$, $ac - bd > B$, $2A|ac - bd - B$, $2A|ad + bc - B$.

Rather than searching all (a, b, c, d) , we can eliminate parts of the parameter space that do not contain solutions. For fixed a and d , the first constraint implies that bc is some factorization of $ad + B$. We can pre-compute a table of factorizations and use

that to cut the search space down to almost $O(n^2)$. You can see the code for this on GitHub at <https://github.com/zaprice/polyvdw>

We can get bounds for $W(x^2 + Bx; 4)$ with this method with rather large values of B , but only a few bounds for the more general $Ax^2 + Bx$ case; if such configurations exist, it seems the numbers involved are much larger. See Appendix C for some of the upper bounds we have. We note two things about these upper bounds:

1. The largest upper bound on $W(x^2 + Bx; 4)$ that we found was when $B = 0$. Note that these are just the upper bounds we found. It is not clear how the real values compares.
2. For $W(2x^2 + Bx; 4)$ and $W(3x^2 + Bx; 4)$ the B for which we could find an upper bound seem scattered and arbitrary. For example, we were not able to find an upper bound for any of $W(2x^2 + Bx; 4)$ for $0 \leq B \leq 56$, but were able to for 57. And then not again until $B = 95$. Again, this may be a limit to our methods and not a statement about the actual values of $W(2x^2 + Bx; 4)$.

ACKNOWLEDGMENT. We thank Sean Prediville and Alex Rice for discussions about prior known upper bounds on $W(p; c)$.

We thank Rob Brady, Steven Brown, Nathan Hayes, Emily Kaplitz, Gary Peng, Yuang Shen, and Zan Xu for proofreading and discussion.

REFERENCES

1. B. van der Waerden, Beweis einer Baudetschen Vermutung (in dutch), *Nieuw Arch. Wisk.* 15 (1927) 212–216.
2. B. van der Waerden, How the proof of Baudet’s conjecture was found, in: L. Mirsky (Ed.), *Studies in Pure Math*, Academic Press, 1971, pp. 251–260.
3. R. Graham, B. Rothschild, J. Spencer, *Ramsey Theory*, Wiley, New York, 1990.
4. B. Landman, A. Robertson, *Ramsey Theory on the integers*, AMS, Providence, 2004.
5. S. Shelah, Primitive recursive bounds for van der Waerden numbers, *Journal of the American Math Society* 1 (1988) 683–697, <http://www.jstor.org/view/08940347/di963031/96p0024f/0>.
6. W. Gowers, A new proof of Szemerédi’s theorem, *Geometric and Functional Analysis* 11 (2001) 465–588, <http://www.dpms.cam.ac.uk/~wtg10/papers/html>.
7. H. Fürstenberg, Ergodic behavior of diagonal measures and a theorem of Szemerédi’s on arithmetic progressions, *Journal of d’Analyse Mathématique* 31 (1977) 204–256, <http://www.cs.umd.edu/~gasarch/TOPICS/vdw/furstenbergsz.pdf>.
8. A. Sárközy, On differences of sets of sequences of integers I, *Acta Math. Sci. Hung.* 31 (1977) 125–149, <http://www.cs.umd.edu/~gasarch/TOPICS/vdw/vdw.html>.
9. V. Bergelson, A. Leibman, Set-polynomials and polynomial extension of the Hales-Jewett theorem, *Annals of Mathematics* 150 (1999) 33–75, <http://www.math.ohio-state.edu/~vitaly/>.
10. M. Walters, Combinatorial proofs of the polynomial van der Waerden theorem and the polynomial Hales-Jewett theorem, *Journal of the London Mathematical Society* 61 (2000) 1–12, <http://jllms.oxfordjournals.org/cgi/reprint/61/1/1>.
11. Shelah, A partition theorem, *Scientiae Math Japonicae* 56 (2002) 413–438, <https://shelah.logic.at/files/199205/679.pdf>.
12. S. Peluse, Bounds for sets with no polynomial progressions, *Forum of Mathematics Pi* 8 (2020) e16, <https://arxiv.org/abs/1909.00309>.
13. S. Peluse, S. Prendville, Quantitative bounds in the non-linear Roth theorem, <http://arxiv.org/abs/1903.02592> (2019).
14. S. Peluse, S. Prendville, A polylogarithmic bound in the nonlinear Roth theorem, *International Math Research Notes* 8 (2022) 5658–5684, <https://arxiv.org/abs/2003.04122>.
15. J. Pintz, W. Steiger, E. Szemerédi, On sets of natural numbers whose difference set contains no squares, *Journal of the London Mathematical Society* 37 (1988) 219–231,

- <http://jllms.oxfordjournals.org/>.
16. A. Rice, A maximal extension of the best-known bounds for the Furstenberg-Sarkozy Theorem, *Acta Arithmetica* (2019) 1–41
<http://arxiv.org/abs/1903.02592>.
 17. M. Harnel, N. Lyall, A. Rice, Improved bounds on Sarkozy’s Theorem for quadratic polynomials, *International Math Research Notices* 8 (2013) 1761–1782,
<https://arxiv.org/abs/1111.5786>.
 18. N. Routledge, Computing Farey sequences, *The Mathematical Gazette*
<https://www.jstor.org/stable/27821716>.

WILLIAM GASARCH *Department of Computer Science, University of MD at College Park*
gasarch@umd.edu

CLYDE KRUSKAL *Department of Computer Science, University of MD at College Park*
ckruskal@umd.edu

JUSTIN KRUSKAL *EPIC Computing*
tinsuj@gmail.com

ZACH PRICE *Department of Mathematics, George Mason University*
zprice3@masonlive.gmu.edu

A. SOME EXACT VALUES OF $W(ax^2 + bx; 2)$

We present a table of $W(p(x); 2)$ for $p(x) = ax^2 + bx$ for $0 \leq a \leq 10$ and $-10 \leq b \leq 10$.

The values for $a, b \geq 0$ were obtained by using our formulas for an upper bound and then searching for a 2-coloring for the lower bound.

| | | a | | | | | | | | | | |
|-----|-----|-----|----|----|----|----|----|----|----|-----|-----|----|
| | | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| b | -10 | 21 | 1 | 1 | 9 | 9 | 1 | 25 | 11 | 13 | 17 | 1 |
| | -9 | 19 | 1 | 9 | 1 | 7 | 5 | 7 | 37 | 15 | 1 | 23 |
| | -8 | 17 | 1 | 1 | 7 | 1 | 7 | 9 | 13 | 1 | 21 | 25 |
| | -7 | 15 | 1 | 7 | 5 | 5 | 25 | 11 | 1 | 19 | 61 | 29 |
| | -6 | 13 | 1 | 1 | 1 | 5 | 9 | 1 | 17 | 21 | 25 | 73 |
| | -5 | 11 | 1 | 5 | 13 | 7 | 1 | 15 | 49 | 25 | 29 | 31 |
| | -4 | 9 | 1 | 1 | 5 | 1 | 13 | 17 | 23 | 25 | 33 | 37 |
| | -3 | 7 | 1 | 3 | 1 | 11 | 37 | 19 | 25 | 31 | 73 | 41 |
| | -2 | 5 | 1 | 1 | 9 | 13 | 19 | 49 | 29 | 33 | 39 | 41 |
| | -1 | 3 | 1 | 7 | 25 | 17 | 21 | 27 | 61 | 37 | 41 | 47 |
| | 0 | 1 | 5 | 9 | 13 | 17 | 21 | 25 | 29 | 33 | 37 | 41 |
| 1 | 3 | 13 | 13 | 17 | 23 | 49 | 33 | 37 | 43 | 85 | 53 | |
| 2 | 5 | 11 | 25 | 21 | 25 | 31 | 33 | 41 | 45 | 51 | 97 | |
| 3 | 7 | 13 | 19 | 37 | 29 | 33 | 37 | 73 | 49 | 49 | 59 | |
| 4 | 9 | 17 | 21 | 27 | 49 | 37 | 41 | 47 | 49 | 57 | 61 | |
| 5 | 11 | 25 | 25 | 29 | 35 | 61 | 45 | 49 | 55 | 97 | 61 | |
| 6 | 13 | 23 | 25 | 31 | 37 | 43 | 73 | 53 | 57 | 61 | 65 | |
| 7 | 15 | 25 | 31 | 49 | 41 | 45 | 51 | 85 | 61 | 65 | 71 | |
| 8 | 17 | 29 | 33 | 39 | 41 | 49 | 53 | 59 | 97 | 69 | 73 | |
| 9 | 19 | 37 | 37 | 37 | 47 | 73 | 55 | 61 | 67 | 109 | 77 | |
| 10 | 21 | 35 | 49 | 45 | 49 | 51 | 57 | 65 | 69 | 75 | 121 | |

The numbers tend to increase with increasing a and $|b|$. Some of the diagonals have patterns which likely can be used to make conjectures that are almost surely true. For example:

$$(\forall a \geq 0)[W(ax^2 - (a - 1)x; 2) = 2a + 3].$$

B. SOME EXACT VALUES OF $W(ax^2 + bx; 3)$

We present a table of $W(p(x); 3)$ for $p(x) = ax^2 + bx$ for $0 \leq a \leq 5$ and $-5 \leq b \leq 5$.

The values were obtained by computer.

| | | <i>a</i> | | | | | |
|----------|----|----------|----|-----|-----|-----|-----|
| | | 0 | 1 | 2 | 3 | 4 | 5 |
| <i>b</i> | -5 | 16 | 1 | 64 | 61 | 217 | 1 |
| | -4 | 13 | 1 | 1 | 91 | 1 | 289 |
| | -3 | 10 | 1 | 10 | 1 | 135 | 171 |
| | -2 | 7 | 1 | 1 | 68 | 97 | 171 |
| | -1 | 4 | 1 | 49 | 105 | 190 | 183 |
| | 0 | 1 | 29 | 57 | 85 | 113 | 141 |
| | 1 | 4 | 73 | 76 | 65 | 156 | 253 |
| | 2 | 7 | 64 | 145 | 123 | 151 | ? |
| | 3 | 10 | 37 | 95 | 217 | ? | ? |
| | 4 | 13 | 65 | 127 | ? | 289 | ? |
| 5 | 16 | 55 | ? | 109 | ? | 361 | |

C. SOME UPPER BOUNDS ON $W(ax^2 + bx; 4)$

We give bounds for $W(g; 4)$ where g is of the form $Ax^2 + Bx$. Only bounds for coprime coefficients (A, B) are presented. Each row of the table gives g, x, y, z, w (as in Figure 4), and the bound. We give four such tables.




Figure 4. In any $(g(x); 4)$ -proper coloring, $\text{COL}(1) = \text{COL}(1 + w)$

Table for $x^2 + Bx$ where $0 \leq B \leq 20$.

| g | x | y | z | w | $W(g(x); 4) \leq$ |
|-------------|--------|--------|--------|-------------|------------------------|
| x^2 | 10,608 | 13,108 | 16,133 | 290,085,289 | 84,149,474,894,213,522 |
| $x^2 + x$ | 299 | 302 | 327 | 113,262 | 12,828,393,907 |
| $x^2 + 2x$ | 91 | 127 | 211 | 257,463 | 66,287,711,296 |
| $x^2 + 3x$ | 35 | 43 | 53 | 3,308 | 10,952,789 |
| $x^2 + 4x$ | 80 | 84 | 92 | 10,197 | 104,019,598 |
| $x^2 + 5x$ | 70 | 81 | 100 | 11,250 | 126,618,751 |
| $x^2 + 6x$ | 70 | 86 | 106 | 13,232 | 175,165,217 |
| $x^2 + 7x$ | 638 | 785 | 923 | 988,338 | 976,818,920,611 |
| $x^2 + 8x$ | 160 | 168 | 184 | 40,788 | 1,663,987,249 |
| $x^2 + 9x$ | 35 | 37 | 44 | 3,242 | 10,539,743 |
| $x^2 + 10x$ | 144 | 150 | 165 | 36,075 | 1,301,766,376 |
| $x^2 + 11x$ | 364 | 472 | 727 | 1,263,252 | 1,595,819,511,277 |
| $x^2 + 12x$ | 140 | 172 | 212 | 52,928 | 2,802,008,321 |
| $x^2 + 13x$ | 119 | 129 | 143 | 38,016 | 1,445,710,465 |
| $x^2 + 14x$ | 66 | 96 | 135 | 25,395 | 645,261,556 |
| $x^2 + 15x$ | 120 | 138 | 215 | 54,364 | 2,956,259,957 |
| $x^2 + 16x$ | 75 | 99 | 141 | 45,177 | 2,041,684,162 |
| $x^2 + 17x$ | 123 | 165 | 255 | 232,908 | 54,250,095,901 |
| $x^2 + 18x$ | 70 | 74 | 88 | 12,968 | 168,402,449 |
| $x^2 + 19x$ | 65 | 66 | 69 | 6,852 | 47,080,093 |
| $x^2 + 20x$ | 84 | 96 | 115 | 24,261 | 589,081,342 |

Table for $x^2 + Bx$ where $1980 \leq B \leq 2000$.

| g | x | y | z | w | $W(g(x); 4) \leq$ |
|----------------|-------|-------|-------|------------|-----------------------|
| $x^2 + 1,980x$ | 1,683 | 2,145 | 2,915 | 25,524,829 | 651,567,434,640,662 |
| $x^2 + 1,981x$ | 1,674 | 1,735 | 2,026 | 14,236,652 | 202,710,462,976,717 |
| $x^2 + 1,982x$ | 1,248 | 1,495 | 1,731 | 6,882,723 | 47,385,517,451,716 |
| $x^2 + 1,983x$ | 3,498 | 3,549 | 3,664 | 24,967,678 | 623,434,455,617,159 |
| $x^2 + 1,984x$ | 860 | 975 | 2,585 | 12,424,497 | 154,392,775,905,058 |
| $x^2 + 1,985x$ | 867 | 1,098 | 2,365 | 11,200,200 | 125,466,712,437,001 |
| $x^2 + 1,986x$ | 1,900 | 2,432 | 2,908 | 19,712,552 | 388,623,855,480,977 |
| $x^2 + 1,987x$ | 3,048 | 3,393 | 3,987 | 39,165,018 | 1,533,976,455,831,091 |
| $x^2 + 1,988x$ | 508 | 738 | 1,194 | 6,489,996 | 42,132,950,192,065 |
| $x^2 + 1,989x$ | 2,023 | 2,288 | 3,094 | 18,950,528 | 359,160,204,078,977 |
| $x^2 + 1,990x$ | 1,364 | 1,610 | 2,100 | 13,163,856 | 173,313,300,862,177 |
| $x^2 + 1,991x$ | 1,330 | 1,519 | 1,814 | 7,817,030 | 61,121,521,727,631 |
| $x^2 + 1,992x$ | 975 | 1,065 | 1,871 | 10,120,498 | 102,444,639,800,021 |
| $x^2 + 1,993x$ | 1,985 | 2,349 | 4,373 | 68,596,488 | 4,705,614,878,734,729 |
| $x^2 + 1,994x$ | 1,246 | 1,350 | 1,716 | 8,551,440 | 73,144,177,644,961 |
| $x^2 + 1,995x$ | 891 | 1,185 | 1,464 | 10,543,450 | 111,185,372,085,251 |
| $x^2 + 1,996x$ | 705 | 995 | 1,793 | 7,390,317 | 54,631,536,433,222 |
| $x^2 + 1,997x$ | 1,081 | 1,136 | 1,391 | 8,040,026 | 64,658,074,012,599 |
| $x^2 + 1,998x$ | 1,292 | 1,732 | 3,704 | 39,649,768 | 1,572,183,322,690,289 |
| $x^2 + 1,999x$ | 1,235 | 1,757 | 2,789 | 14,633,322 | 214,163,364,766,363 |
| $x^2 + 2,000x$ | 184 | 280 | 984 | 5,592,000 | 31,281,648,000,001 |

Table for $2x^2 + Bx$ for assorted B .

| g | x | y | z | w | $W(g(x); 4) \leq$ |
|-----------------|--------|--------|--------|---------------|---------------------------|
| $2x^2 + 57x$ | 3,969 | 4,035 | 4,295 | 38,199,155 | 2,918,353,062,779,886 |
| $2x^2 + 95x$ | 707 | 758 | 1,008 | 14,365,638 | 412,744,475,029,699 |
| $2x^2 + 171x$ | 11,907 | 12,105 | 12,885 | 343,792,395 | 236,386,480,508,171,596 |
| $2x^2 + 285x$ | 2,121 | 2,274 | 3,024 | 129,290,742 | 33,432,228,781,682,599 |
| $2x^2 + 399x$ | 27,783 | 28,245 | 30,065 | 1,871,758,595 | 7,006,961,222,744,427,456 |
| $2x^2 + 455x$ | 3,320 | 3,663 | 4,170 | 39,229,128 | 3,077,866,816,534,009 |
| $2x^2 + 511x$ | 2,772 | 3,367 | 6,282 | 131,899,720 | 34,795,139,672,913,721 |
| $2x^2 + 627x$ | 43,659 | 44,385 | 47,245 | 4,622,097,755 | 5,834,090,064,188,269,204 |
| $2x^2 + 805x$ | 1,210 | 1,303 | 2,920 | 87,446,025 | 15,293,684,970,651,376 |
| $2x^2 + 855x$ | 5,548 | 7,087 | 13,262 | 530,042,423 | 561,890,393,545,693,524 |
| $2x^2 + 1,011x$ | 5,164 | 6,568 | 9,889 | 318,517,859 | 202,907,575,025,443,212 |
| $2x^2 + 1,153x$ | 12,705 | 12,726 | 12,970 | 352,488,525 | 248,496,726,932,620,576 |
| $2x^2 + 1,199x$ | 8,245 | 8,710 | 9,748 | 221,108,291 | 97,778,017,806,722,272 |
| $2x^2 + 1,295x$ | 14,030 | 14,355 | 22,244 | 1,162,712,925 | 2,703,804,197,637,349,126 |
| $2x^2 + 1,301x$ | 25,622 | 26,105 | 28,172 | 1,638,880,116 | 5,371,858,201,423,377,829 |
| $2x^2 + 1,365x$ | 9,960 | 10,989 | 12,510 | 353,062,152 | 249,306,248,279,579,689 |
| $2x^2 + 1,459x$ | 954 | 1,174 | 1,379 | 58,465,486 | 6,836,511,407,576,467 |
| $2x^2 + 1,545x$ | 11,298 | 11,815 | 12,860 | 425,440,418 | 361,999,755,841,475,259 |
| $2x^2 + 1,685x$ | 10,695 | 10,968 | 11,570 | 289,144,125 | 167,209,137,251,881,876 |
| $2x^2 + 1,753x$ | 3,586 | 5,236 | 8,232 | 181,967,394 | 66,224,583,947,144,155 |
| $2x^2 + 1,851x$ | 50,031 | 51,441 | 55,164 | 6,379,649,159 | 7,612,882,297,751,201,408 |
| $2x^2 + 1,913x$ | 2,261 | 3,366 | 5,324 | 81,424,299 | 13,259,988,699,966,790 |

Table for $3x^2 + Bx$ for assorted B .

| g | x | y | z | w | $W(g(x); 4) \leq$ |
|-----------------|--------|--------|--------|---------------|---------------------------|
| $3x^2 + x$ | 42,273 | 42,660 | 43,375 | 5,738,872,934 | 6,570,267,294,984,419,923 |
| $3x^2 + 143x$ | 13,244 | 13,332 | 13,442 | 554,651,696 | 922,915,590,942,221,777 |
| $3x^2 + 172x$ | 4,452 | 4,712 | 5,189 | 88,862,311 | 23,689,546,233,099,656 |
| $3x^2 + 200x$ | 1,896 | 2,204 | 5,004 | 115,177,723 | 39,797,746,661,938,788 |
| $3x^2 + 235x$ | 11,155 | 11,270 | 11,610 | 583,594,418 | 1,021,747,471,306,964,403 |
| $3x^2 + 274x$ | 9,322 | 11,610 | 16,903 | 1,125,018,929 | 3,797,003,080,080,107,670 |
| $3x^2 + 344x$ | 8,904 | 9,424 | 10,378 | 355,449,244 | 379,032,617,455,054,545 |
| $3x^2 + 361x$ | 3,540 | 4,658 | 7,703 | 397,333,094 | 473,620,906,200,085,443 |
| $3x^2 + 400x$ | 3,792 | 4,408 | 10,008 | 460,710,892 | 636,763,762,306,663,793 |
| $3x^2 + 407x$ | 2,806 | 3,401 | 6,131 | 122,898,626 | 45,312,266,837,804,411 |
| $3x^2 + 412x$ | 2,077 | 2,829 | 5,839 | 392,773,686 | 462,813,667,064,838,421 |
| $3x^2 + 520x$ | 7,616 | 9,244 | 12,716 | 515,261,395 | 796,483,183,467,963,476 |
| $3x^2 + 556x$ | 9,400 | 9,408 | 9,451 | 273,674,799 | 224,693,838,986,259,448 |
| $3x^2 + 592x$ | 15,744 | 16,472 | 17,944 | 994,061,387 | 2,964,474,711,857,432,412 |
| $3x^2 + 643x$ | 50,932 | 51,357 | 52,351 | 8,273,167,696 | 2,421,731,687,255,606,001 |
| $3x^2 + 688x$ | 17,808 | 18,848 | 20,756 | 1,421,796,976 | 6,064,520,901,084,553,217 |
| $3x^2 + 725x$ | 3,172 | 3,185 | 3,278 | 34,869,750 | 3,647,723,675,756,251 |
| $3x^2 + 728x$ | 16,744 | 17,360 | 18,928 | 1,174,742,491 | 4,140,060,615,695,188,692 |
| $3x^2 + 797x$ | 2,847 | 3,082 | 3,524 | 148,907,272 | 66,520,245,642,541,737 |
| $3x^2 + 814x$ | 5,612 | 6,802 | 12,262 | 491,594,504 | 724,995,869,246,944,305 |
| $3x^2 + 932x$ | 1,820 | 2,229 | 2,799 | 37,745,311 | 4,274,160,686,090,016 |
| $3x^2 + 1,085x$ | 1,190 | 1,344 | 1,540 | 10,401,450 | 324,581,771,880,751 |
| $3x^2 + 1,087x$ | 9,800 | 9,909 | 11,434 | 604,108,526 | 1,094,841,990,223,645,791 |
| $3x^2 + 1,112x$ | 18,800 | 18,816 | 18,902 | 1,094,699,196 | 3,595,100,206,474,645,201 |