

CMSC351 Test 1 Reference**Asymptotic Notations.**

$\Theta(g(n)) = \{f(n) : \text{there exist positive constants } c_1, c_2, \text{ and } n_0 \text{ such that } 0 \leq c_1 g(n) \leq f(n) \leq c_2 g(n) \text{ for all } n \geq n_0\}$.

$O(g(n)) = \{f(n) : \text{there exist positive constants } c \text{ and } n_0 \text{ such that } 0 \leq f(n) \leq cg(n) \text{ for all } n \geq n_0\}$.

$\Omega(g(n)) = \{f(n) : \text{there exist positive constants } c \text{ and } n_0 \text{ such that } 0 \leq cg(n) \leq f(n) \text{ for all } n \geq n_0\}$.

$f(n) = o(g(n))$ if $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$.

$f(n) = \omega(g(n))$ if $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$.

$f(n) \sim g(n)$ if $f(n) = g(n) + o(g(n))$.

Logarithms.

$$\begin{aligned}
 a &= b^{\log_b a} \\
 \log_c(ab) &= \log_c a + \log_c b \\
 \log_b a^n &= n \log_b a \\
 \log_b a &= \frac{\log_c a}{\log_c b} \\
 \log_b(1/a) &= -\log_b a \\
 \log_b a &= \frac{1}{\log_a b} \\
 a^{\log_b n} &= n^{\log_b a} \\
 a^{f(n)} &= e^{f(n) \ln a} \\
 (\log_a f(n))' &= \frac{f'(n)}{f(n) \ln a} \\
 (a^{f(n)})' &= \ln a f'(n) a^{f(n)}
 \end{aligned}$$

Quadratic Formula.

$$ax^2 + bx + c = 0 \Rightarrow x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Summations.

Simple Arithmetic Series:

$$\sum_{k=1}^n k = 1 + 2 + \cdots + n = \frac{n(n+1)}{2}$$

$$\sum_{k=1}^n k^2 = 1 + 4 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{k=1}^n k^3 = 1 + 8 + \cdots + n^3 = \frac{n^2(n+1)^2}{4}$$

General Arithmetic Series:

$$\sum_{k=m}^n k = 1 + 2 + \cdots + n = \frac{(n-m+1)(n+m)}{2}$$

Geometric series:

$$\sum_{k=0}^n x^k = 1 + x + x^2 + \cdots + x^n = \frac{x^{n+1} - 1}{x - 1} \quad x \neq 1$$

$$\sum_{k=0}^{\infty} x^k = \frac{1}{1-x} \quad |x| < 1$$

Recurrences “ SMT: Simplified Master Theorem”:

$$T(n) = \begin{cases} aT(n/b) + n^k & n > 1 \\ 1 & n = 1 \end{cases}$$

implies

$$T(n) = \begin{cases} \Theta(n^{\log_b a}) & a > b^k \\ \Theta(n^k) & a < b^d \\ \Theta(n^k \log_b n) & a = b^k \end{cases} .$$