

START

RECORDING

Discrete Probability

CMSC 250

Axiomatic Definitions, Basic Problems with Cards

Informal Definition of Probability

- Probability that *blah* happens:


$$\frac{\# \text{ possibilities that } \textit{blah} \text{ happens}}{\# \text{ all possibilities}}$$

Informal Definition of Probability

- Probability that **blah** happens:

$$\frac{\# \text{ possibilities that } \textit{blah} \text{ happens}}{\# \text{ all possibilities}}$$

- This definition is owed to [Andrey Kolmogorov](#), and assumes *that all possibilities are equally likely!*




First Examples

- Experiment #1: Tossing the same coin 3 times.


First Examples

- Experiment #1: Tossing the same coin 3 times.
 - What is the probability that I don't get any heads?




First Examples

- Experiment #1: Tossing the same coin 3 times.
 - What is the probability that I don't get any heads?
 - Why?
 - Set of different *events*?
 - $\{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$ (8 of them)
 - Set of events with **no heads**:
 - $\{TTT\}$ (1 of them)
 - Hence the answer: $\frac{1}{8}$



First Examples

- Experiment #1: Tossing the same coin 3 times.
 - What is the probability that I don't get any heads?
 - Why?
 - Set of different *events*?
 - $\{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$ (8 of them)
 - Set of events with **no heads**:
 - $\{TTT\}$ (1 of them)
 - Hence the answer: $\frac{1}{8}$



Implicit assumption: all individual outcomes (HHH, HHT, HTH, ...) **are considered equally likely** (probability $\frac{1}{8}$)

Practice

- Experiment #2: I roll two dice.
 - Probability that I hit **seven** = ?

$$\frac{1}{12}$$

$$\frac{1}{6}$$

$$\frac{7}{12}$$

Something
else

Practice

- Experiment #2: I roll two dice.

- Probability that I hit **seven** = ?

- **Why?**


- Set of different *events*?

- $\{(1, 1), (1, 2), \dots, (6, 1)\}$ (36 of them)

- Set of events where we hit 7.

- $\{(2, 5), (5, 2), (3, 4), (4, 3), (1, 6), (6, 1)\}$ (6 of them)

- Hence the answer: $\frac{6}{36} = \frac{1}{6}$



Practice

- Experiment #2: I roll two dice.

- Probability that I hit **seven** = ?

- **Why?**

- Set of different *events*?


- $\{(1, 1), (1, 2), \dots, (6, 1)\}$ (36 of them)

- Set of events where we hit 7.

- $\{(2, 5), (5, 2), (3, 4), (4, 3), (1, 6), (6, 1)\}$ (6 of them)

- Hence the answer: $\frac{6}{36} = \frac{1}{6}$

- Probability that I hit **two** = ?



Practice

- Experiment #2: I roll two dice.

- Probability that I hit **seven** = ?

- **Why?**

- Set of different *events*?

- $\{(1, 1), (1, 2), \dots, (6, 1)\}$ (36 of them)


- Set of events where we hit 7.

- $\{(2, 5), (5, 2), (3, 4), (4, 3), (1, 6), (6, 1)\}$ (6 of them)

- Hence the answer: $\frac{6}{36} = \frac{1}{6}$

- Probability that I hit **two** = ?


- Same procedure



$$\frac{1}{36}$$


Poker Practice

- Full deck = 52 cards, 13 of each suit:



Poker Practice

- Full deck = 52 cards, 13 of each suit:
- **Flush:** 5 cards of the same suit
- What is the probability of getting a flush?



Probability of a Flush

- How many 5-card hands are there?

Probability of a Flush

- How many 5-card hands are there? $\binom{52}{5}$

Probability of a Flush

- How many 5-card hands are there? $\binom{52}{5}$
- How many 5-card hands are flushes?

Probability of a Flush

- How many 5-card hands are there? $\binom{52}{5}$
- How many 5-card hands are flushes?
 - Choose a suit in one of 4 ways...

Probability of a Flush

- How many 5-card hands are there? $\binom{52}{5}$
- How many 5-card hands are flushes?
 - Choose a suit in one of 4 ways...
 - Given suit choose any 5 cards out of 13...

Probability of a Flush

- How many 5-card hands are there? $\binom{52}{5}$
- How many 5-card hands are flushes?
 - Choose a suit in one of 4 ways...
 - Given suit choose any 5 cards out of 13...
 - So $4 * \binom{13}{5}$

Probability of a Flush

- How many 5-card hands are there? $\binom{52}{5}$
- How many 5-card hands are flushes?
 - Choose a suit in one of 4 ways...
 - Given suit choose any 5 cards out of 13...
 - So $4 * \binom{13}{5}$
- So, probability of being dealt a flush is

$$\frac{4 * \binom{13}{5}}{\binom{52}{5}}$$

Probability of a Flush

- Probability of being dealt a flush is

$$\frac{4 * \binom{13}{5}}{\binom{52}{5}}$$

Probability of a Flush

- Probability of being dealt a flush is

$$\frac{4 * \binom{13}{5}}{\binom{52}{5}}$$

- How likely is this?

Probability of a Flush

- Probability of being dealt a flush is

$$\frac{4 * \binom{13}{5}}{\binom{52}{5}}$$

- How likely is this?
 - Not at all likely: $\approx 0.002 = 0.2\%$ ☹️

Likelihood of a Straight

- Straights are 5 cards of *consecutive rank*
 - Ace can be either end (high or low)
 - No wrap-arounds (e.g Q K A 2 3, suits don't matter)

Likelihood of a Straight

- Straights are 5 cards of *consecutive rank*
 - Ace can be either end (high or low)
 - No wrap-arounds (e.g Q K A 2 3, suits don't matter)
- What is the probability that we are dealt a straight?

Likelihood of a Straight

- Straights are 5 cards of *consecutive rank*
 - Ace can be either end (high or low)
 - No wrap-arounds (e.g Q K A 2 3, suits don't matter)
- What is the probability that we are dealt a straight?
- As before, #possible 5-card hands = $\binom{52}{5}$

Likelihood of a Straight

- Straights are 5 cards of **consecutive rank**
 - Ace can be either end (high or low)
 - No wrap-arounds (e.g Q K A 2 3, suits don't matter)
- What is the probability that we are dealt a straight?
- As before, #possible 5-card hands = $\binom{52}{5}$
- To find out the #straights:
 - Pick lower rank in 10 ways (A-10)
 - Pick a suit in 4 ways
 - Pick the 4 subsequent cards **from any suit** in 4^4 ways

Likelihood of a Straight

- Straights are 5 cards of **consecutive rank**
 - Ace can be either end (high or low)
 - No wrap-arounds (e.g Q K A 2 3, suits don't matter)
- What is the probability that we are dealt a straight?

• As before, #possible 5-card hands = $\binom{52}{5}$

• To find out the #straights:

- Pick lower rank in 10 ways (A-10)
- Pick a suit in 4 ways
- Pick the 4 subsequent cards **from any suit** in 4^4 ways

That's $10 * 4^5$ ways.
So, probability of a
straight = $\frac{10 * 4^5}{\binom{52}{5}}$

Caveat on Flushes

- [Wikipedia](#) says we're wrong about flushes!

Caveat on Flushes


- [Wikipedia](#) says we're wrong about flushes!
- Formally, our flushes included (for example) **3h 4h 5h 6h 7h**
 - Hands like these are called **straight flushes** and Wikipedia does not include them.

Caveat on Flushes

- [Wikipedia](#) says we're wrong about flushes!
- Formally, our flushes included (for example) **3h 4h 5h 6h 7h**
 - Hands like these are called **straight flushes** and Wikipedia does not include them.
 - *How many straight flushes are there?*

Caveat on Flushes

- [Wikipedia](#) says we're wrong about flushes!
- Formally, our flushes included (for example) **3h 4h 5h 6h 7h**
 - Hands like these are called **straight flushes** and Wikipedia does not include them.
 - **How many straight flushes are there?**
 - **40.** Here's why:
 - Pick rank: A through 10 (higher ranks don't allow straights) in **10 ways**
 - Pick suit in **4 ways**



Probability of Non-Straight Flush...

$$\frac{4 * \binom{13}{5} - 40}{\binom{52}{5}} = 0.001965$$

- This is how [Wikipedia](#) defines the probability of a flush. 😊

Probability of a Straight Flush...

$$\frac{40}{\binom{52}{5}} = 0.0000138517$$

Probability of a Straight Flush...

$$\frac{40}{\binom{52}{5}} = 0.0000138517$$

The expected # hands you need to play to get a straight flush is then

$$\lceil \frac{1}{0.0000138517} \rceil = 72,194$$

Same Caveat for Straights

- From the #straights we computed we will have to subtract the 40 possible straight flushes to get...

$$\frac{10 * 4^5 - 40}{\binom{52}{5}} = 0.003925$$

Same Caveat

- From the #straights we computed we will have to subtract the 40 possible straight flushes to get...

$$\frac{10*4^5 - 40}{\binom{52}{5}} = 0.003925 > 0.001965 = \text{probability of flush}$$

- *Flushes, being more rare, beat straights in poker.*

Probability of a Pair

- Try to calculate the probability of a pair!

Probability of a Pair

- Try to calculate the probability of a **pair!**
- Perhaps you thought of the problem like this:
 - The denominator will be $\binom{52}{5}$ (easy), so let's focus on the **numerator:**
 1. First choose rank in 13 ways.
 2. Then, choose two of four suits in $\binom{4}{2} = 6$ ways.
 3. Then, choose 3 cards out of 50 in $\binom{50}{3}$ ways.


Probability of a Pair

- Try to calculate the probability of a **pair!**
- Perhaps you thought of the problem like this:
 - The denominator will be $\binom{52}{5}$ (easy), so let's focus on the **numerator:**

1. First choose rank in 13 ways.

2. Then, choose two of four suits in $\binom{4}{2} = 6$ ways.

3. Then, choose 3 cards out of 50 in $\binom{50}{3}$ ways.



Numerator: $13 \times 6 \times \binom{50}{3}$

Probability of a Pair

- Try to calculate the probability of a **pair!**
- Perhaps you thought of the problem like this:
 - The denominator will be $\binom{52}{5}$ (easy), so let's focus on the **numerator:**

1. First choose rank in 13 ways.

2. Then, choose two of four suits in $\binom{4}{2} = 6$ ways.

3. Then, choose 3 cards out of 50 in $\binom{50}{3}$ ways.

Numerator: $13 \times 6 \times \binom{50}{3}$

• So, probability =
$$\frac{13 \times 6 \times \binom{50}{3}}{\binom{52}{5}}$$

Probability of a Pair


- Try to calculate the probability of a **pair!**
- Perhaps you thought of the problem like this:
 - The denominator will be $\binom{52}{5}$ (easy), so let's focus on the **numerator:**

1. First choose rank in 13 ways.

2. Then, choose two of four suits in $\binom{4}{2} = 6$ ways.

3. Then, choose 3 cards out of 50 in $\binom{50}{3}$ ways.

• So, probability =
$$\frac{13 \times 6 \times \binom{50}{3}}{\binom{52}{5}}$$



Numerator: $13 \times 6 \times \binom{50}{3}$

Is this accurate?

Yes

No

Probability of a Pair

- Try to calculate the probability of a **pair!**
- Perhaps you thought of the problem like this:
 - The denominator will be $\binom{52}{5}$ (easy), so let's focus on the **numerator:**

1. First choose rank in 13 ways.

2. Then, choose two of four suits in $\binom{4}{2} = 6$ ways.

3. Then, choose 3 cards out of 50 in $\binom{50}{3}$ ways.

• So, probability =
$$\frac{13 \times 6 \times \binom{50}{3}}{\binom{52}{5}}$$

Numerator: $13 \times 6 \times \binom{50}{3}$

Is this accurate? **Severe overcount!**

Yes

No

Don't Count Better Hands!

- In the computation before, we included:
 - 2-of-a-kind
 - 3-of-a-kind
 - 4-of-a-kind
 - Full House

Don't Count Better Hands!

- In the computation before, we included:
 - 2-of-a-kind
 - 3-of-a-kind
 - 4-of-a-kind
 - Full House
- To properly compute, we would have to subtract **all** better hands possible with at least one pair.

Joint Probability

Joint Probability (“AND” of Two Events)

- The probability that two events A and B occur **simultaneously** is known as the **joint** probability of A and B and is denoted in a number of ways:

Joint Probability (“AND” of Two Events)

- The probability that two events A and B occur **simultaneously** is known as the **joint** probability of A and B and is denoted in a number of ways:
 - $P(A \cap B)$ (Most useful from a set-theoretic perspective; **we'll be using this**)

Joint Probability (“AND” of Two Events)

- The probability that two events A and B occur **simultaneously** is known as the **joint** probability of A and B and is denoted in a number of ways:
 - $P(A \cap B)$ (Most useful from a set-theoretic perspective; **we'll be using this**)
 - $P(A, B)$ (One sees this a lot in Physics books)

Joint Probability (“AND” of Two Events)

- The probability that two events A and B occur **simultaneously** is known as the **joint** probability of A and B and is denoted in a number of ways:
 - $P(A \cap B)$ (Most useful from a set-theoretic perspective; **we'll be using this**)
 - $P(A, B)$ (One sees this a lot in Physics books)
 - $P(AB)$ (Perhaps most convenient, therefore most common)

Calculating Joints

- Probability that the first coin toss is heads and the second coin toss is tails

Calculating Joints

- Probability that the first coin toss is heads and the second coin toss is tails $\frac{1}{2} \times \frac{1}{2}$

Calculating Joints

- Probability that the first coin toss is heads and the second coin toss is tails $\frac{1}{2} \times \frac{1}{2}$
- Probability that the first die is **at most** a 2 and the second one is 5 **or** 6

Calculating Joints

- Probability that the first coin toss is heads and the second coin toss is tails $\frac{1}{2} \times \frac{1}{2}$
- Probability that the first die is **at most** a 2 and the second one is 5 **or** 6
 - # outcomes of die roll is 6
 - # outcomes where first die is at most 2 is 2
 - Hence, probability of first die roll being at most 2 is $\frac{1}{3}$

Calculating Joints

- Probability that the first coin toss is heads and the second coin toss is tails $\frac{1}{2} \times \frac{1}{2}$
- Probability that the first die is **at most** a 2 and the second one is 5 **or** 6
 - # outcomes of die roll is 6
 - # outcomes where first die is at most 2 is 2
 - Hence, probability of first die roll being at most 2 is $\frac{1}{3}$
 - Similarly, probability of second die roll being 5 or 6 is $\frac{1}{3}$.
 - Hence, probability that **both** events happen (joint probability) is $\frac{1}{9}$.

Calculating Joints

- Jason's going to flip a coin and then pick a card from a 52-card deck.
 - Probability that the coin is heads and the card has rank 8?



Calculating Joints

- Jason's going to flip a coin and then pick a card from a 52-card deck
 - Probability that the coin is heads and the card has rank 8?



- This is because $P(\textit{coin} = H) = \frac{1}{2}$ and $P(\textit{card_rank} = 8) = \frac{4}{52} = \frac{1}{13}$
 - So their joint probability is $\frac{1}{2} \times \frac{1}{13} = \frac{1}{26}$

The Law of Joint Probability

$$P(A \cap B) = P(A) \cdot P(B)$$

$$P(A_1 \cap A_2 \cap \cdots \cap A_n) = \prod_{i=1}^n P(A_i)$$

The Law of Joint Probability

$$P(A \cap B) = P(A) \cdot P(B)$$

$$P(A_1 \cap A_2 \cap \cdots \cap A_n) = \prod_{i=1}^n P(A_i)$$

- Unfortunately, this “law” is not always applicable!

The Law of Joint Probability

$$P(A \cap B) = P(A) \cdot P(B)$$

$$P(A_1 \cap A_2 \cap \cdots \cap A_n) = \prod_{i=1}^n P(A_i)$$

- Unfortunately, this “law” is not always applicable!
- It is applicable only when all the different events A_i are *independent* (sometimes called *marginally independent*) of each other.
- Let’s look at an example.

What If The Events Influence Each Other?

- Probability that a die is even and that it is 2.

What If The Events Influence Each Other?

- Probability that a die is even and that it is 2.
 - Probability that the die is even = $\frac{1}{2}$

What If The Events Influence Each Other?

- Probability that a die is even and that it is 2.
 - Probability that the die is even = $\frac{1}{2}$
 - Probability that the die is two = $\frac{1}{6}$

What If The Events Influence Each Other?

- Probability that a die is even and that it is 2.
 - Probability that the die is even = $\frac{1}{2}$
 - Probability that the die is two = $\frac{1}{6}$
 - Probability the die is even and the die is two = $\frac{1}{12}$???

What If The Events Influence Each Other?

- Probability that a die **is even and that it is 2**.
 - Probability that the die is even = $\frac{1}{2}$
 - Probability that the die is two = $\frac{1}{6}$
 - Probability the die is even **and** the die is two = $\frac{1}{12}$???
- **NO!**
 - What is the probability that the die is even and the die is 2?




What If The Events Influence Each Other?

- Probability that a die **is even and that it is 2**.
 - Probability that the die is even = $\frac{1}{2}$
 - Probability that the die is two = $\frac{1}{6}$
 - Probability the die is even **and** the die is two = $\frac{1}{12}$???
- **NO!**
 - What is the probability that the die is even and the die is 2?



Set-Theoretic Interpretation


- Notice that the event A: “Die roll is even” is a **superset** of the event B: “Die roll comes 2”



- Die roll even
- Die roll comes 2

Set-Theoretic Interpretation

- Notice that the event A: “Die roll is even” is a **superset** of the event B: “Die roll comes 2”



- Die roll even
- Die roll comes 2

- Since $A \cap B = A$, $P(A \cap B) = P(A) = \frac{1}{6}$

Calculating Joints

- [The University of Southern North Dakota](#) offers a Discrete Mathematics Course where the possible grades are A through G. (No + or -)

Calculating Joints

- [The University of Southern North Dakota](#) offers a Discrete Mathematics Course where the possible grades are A through G. (No + or -)
- What is the probability that Jason gets **both** an A and a G in that course?

Calculating Joints

- [The University of Southern North Dakota](#) offers a Discrete Mathematics Course where the possible grades are A through G. (No + or -)
- What is the probability that Jason gets **both** an A and a G in that course?
 - Clearly, it **can't** be

$$(probability\ Jason\ gets\ an\ A) \times (probability\ Jason\ gets\ a\ B) = \frac{1}{7} \times \frac{1}{7} = \frac{1}{49}$$

Calculating Joints

- [The University of Southern North Dakota](#) offers a Discrete Mathematics Course where the possible grades are A through G. (No + or -)
- What is the probability that Jason gets **both** an A and a G in that course?
 - Clearly, it **can't** be

~~$(\text{probability Jason gets an A}) \times (\text{probability Jason gets a B}) = \frac{1}{7} \times \frac{1}{7} = \frac{1}{49}$~~

Calculating Joints

- [The University of Southern North Dakota](#) offers a Discrete Mathematics Course where the possible grades are A through G. (No + or -)
- What is the probability that Jason gets **both** an A and a G in that course?
 - Clearly, it **can't** be

~~$(\text{probability Jason gets an A}) \times (\text{probability Jason gets a B}) = \frac{1}{7} \times \frac{1}{7} = \frac{1}{49}$~~

- It is **0**. Those two events cannot happen **jointly!**

Calculating Joints

- [The University of Southern North Dakota](#) offers a Discrete Mathematics Course where the possible grades are A through G. (No + or -)
- What is the probability that Jason gets **both** an A and a G in that course?
 - Clearly, it **can't** be

~~$(\text{probability Jason gets an A}) \times (\text{probability Jason gets a B}) = \frac{1}{7} \times \frac{1}{7} = \frac{1}{49}$~~

- It is **0**. Those two events cannot happen **jointly!**
- Events such as these are called **disjoint** or **mutually disjoint**.


Set-Theoretic Interpretation

- A = “Jason gets an A in USND’s 250”
- G=“Jason gets a G in USND’s 250”



Set-Theoretic Interpretation

- A = “Jason gets an A in USND’s 250”
- G = “Jason gets a G in USND’s 250”



- Note that $A \cap G = \emptyset$, so there are no common outcomes.
 - So $P(A \cap G) = 0$

Calculating Joints

- I have my original die again.
 - Probability that it comes up 1, 2 or 3 = $\frac{1}{2}$
 - Probability that it comes up 3, 4 or 5 = $\frac{1}{2}$
 - What is the probability that it comes up 1, 2 or 3 **and** 3, 4 or 5?

Calculating Joints

- I have my original die again.
 - Probability that it comes up 1, 2 or 3 = $\frac{1}{2}$
 - Probability that it comes up 3, 4 or 5 = $\frac{1}{2}$
 - What is the probability that it comes up 1, 2 or 3 **and** 3, 4 or 5?

$$\frac{1}{6}$$

$$\frac{1}{5}$$

$$\frac{1}{4}$$

$$\frac{1}{3}$$

Calculating Joints


- I have my original die again.
 - Probability that it comes up 1, 2 or 3 = $\frac{1}{2}$
 - Probability that it comes up 3, 4 or 5 = $\frac{1}{2}$
 - What is the probability that it comes up 1, 2 or 3 **and** 3, 4 or 5?



- Note that the only common outcome between the two events is **3**, which can come up only **once** out of **six** possibilities.


Set-Theoretic Interpretation

- Let A = dice comes up 1, 2, or 3
- Let B = dice comes up 3, 4, or 5
- Let C = dice comes up 1, 2, 3, 4, 5 OR 6



Set-Theoretic Interpretation

- Let A = dice comes up 1, 2, or 3
- Let B = dice comes up 3, 4, or 5
- Let C = dice comes up 1, 2, 3, 4, 5 OR 6



- Then, probability that the dice comes up 3 = $\frac{1}{6}$

Dependent and Independent Events

Independent Events *(informally)*

- Two events are independent if **one does not influence the other.**

Independent Events *(informally)*

- Two events are independent if **one does not influence the other.**
- Examples:
 - The event E_1 = “first coin toss” and E_2 = “second coin toss”
 - With the same die, the events E_1 = “roll 1”, E_2 = “roll 2”, E_3 = “roll 3”
 - Jason flips a coin and then picks a card.

Independent Events (*informally*)

- Two events are independent if **one does not influence the other.**
- Examples:
 - The event E1 = “first coin toss” and E2 = “second coin toss”
 - With the same die, the events E1 = “roll 1”, E2 = “roll 2”, E3 = “roll 3”
 - Jason flips a coin and then picks a card.
- Counter-examples:
 - E1 = “Die is even”, E2=“Die is 6”
 - E1= “Grade in 250” and “Passing 250”

Law of Joint Probability *(informally)*

- Two events are independent if **one does not influence the other.**
 - This definition is a bit **too informal**, so mathematicians tend to avoid it.

Law of Joint Probability (*informally*)

- Two events are independent if **one does not influence the other.**
 - This definition is a bit **too informal**, so mathematicians tend to avoid it.
- Formally, we define that A and B are **independent** if

$$P(A \cap B) = P(A) \cdot P(B)$$

Disjoint or Independent?

1. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It rains in Athens, Greece today"

Disjoint


Independent

Both

Neither


Disjoint or Independent?

1. $E_1 = \text{"It rains in College Park, MD today"}$
 $E_2 = \text{"It rains in Athens, Greece today"}$



Disjoint or Independent?

1. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It rains in Athens, Greece today"



Weather is weird!

2. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It is sunny in College Park, MD today"



Disjoint or Independent?

1. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It rains in Athens, Greece today"

Disjoint

Independent

Both

Neither

Weather is weird!

2. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It is sunny in College Park, MD today"

Disjoint

Independent

Both

Neither

Disjoint or Independent?

1. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It rains in Athens, Greece today"

Disjoint

Independent

Both

Neither

Weather is weird!

2. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It is sunny in College Park, MD today"

Disjoint

Independent

Both

Neither

3. $E_1 =$ Die #1 comes at most 4
 $E_2 =$ Die #2 comes at least 5

Disjoint

Independent

Both

Neither


Disjoint or Independent?

1. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It rains in Athens, Greece today"




Weather is weird!

2. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It is sunny in College Park, MD today"



3. $E_1 =$ Die #1 comes at most 4
 $E_2 =$ Die #2 comes at least 5




Disjoint or Independent?

1. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It rains in Athens, Greece today"




Weather is weird!

2. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It is sunny in College Park, MD today"



3. $E_1 =$ Die #1 comes at most 4
 $E_2 =$ Die #2 comes at least 5



4. $E_1 =$ Student gets a C
 $E_2 =$ Student passes the class




Disjoint or Independent?

1. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It rains in Athens, Greece today"



Weather is weird!

2. $E_1 =$ "It rains in College Park, MD today"
 $E_2 =$ "It is sunny in College Park, MD today"



3. $E_1 =$ Die #1 comes at most 4
 $E_2 =$ Die #2 comes at least 5



4. $E_1 =$ Student gets a C
 $E_2 =$ Student passes the class



Recap: “Disjoint” vs “Independent”

- Friends don't let friends get confused between “disjoint” and “independent”!

Disjoint	Independent
Has a set-theoretic interpretation!	Has a causality interpretation!
Means that $P(A \cap B) = 0$	Means that $P(A \cap B) = P(A) \cdot P(B)$
Means that $P(A \cup B) = P(A) + P(B)$	Means that $P(A \cup B) = P(A) + P(B) - P(A) \cdot P(B)$

Disjoint Probability (“OR” of Two Events)

- Jason rolls two dice.
 - **What is the probability that he rolls a 7 or a 9?**

Disjoint Probability (“OR” of Two Events)

- Jason rolls two dice.
 - **What is the probability that he rolls a 7 or a 9?**
 - #Ways to roll a 7 is 6.
 - #Ways to roll a 9 is 4: (6, 3), (5, 4), (4, 5), (3, 6)
 - #Ways to roll a 7 OR a 9 is then 10.
 - Therefore, the probability is $\frac{10}{36} = \frac{5}{18}$
 - Key: Rolling a 7 and a 9 are **disjoint events**.

Disjoint Probability (“OR”)

- 52-card deck
- Probability of drawing a face card (J, Q, K) or a heart

Disjoint Probability (“OR”)

- 52-card deck
- Probability of drawing a face card (J, Q, K) or a heart
 - **Are these disjoint?**

Disjoint Probability (“OR”)

- 52-card deck
- Probability of drawing a face card (J, Q, K) or a heart
 - **Are these disjoint?**
 - **NO**, for example, **Queen of hearts**
- How big is $Face_Card \cup Hearts$?

Disjoint Probability (“OR”)

- 52-card deck
- Probability of drawing a face card (J, Q, K) or a heart
 - **Are these disjoint?**
 - **NO**, for example, **Queen of hearts**
- How big is $Face_Card \cup Hearts$ (abbrv. F, H below)?
 - Use law of **inclusion / exclusion!**

$$|F \cup H| = |F| + |H| - |F \cap H| = 12 + 13 - 3 = 22$$

Disjoint Probability (“OR”)

- 52-card deck
- Probability of drawing a face card (J, Q, K) or a heart
 - **Are these disjoint?**
 - **NO**, for example, **Queen of hearts**
- How big is $Face_Card \cup Hearts$ (abbrv. F, H below)?
 - Use law of **inclusion / exclusion!**

$$|F \cup H| = |F| + |H| - |F \cap H| = 12 + 13 - 3 = 22$$

- So probability = $\frac{22}{52} = \frac{11}{26}$.

Alternative Viewpoint

- $P(F) = \frac{12}{52}$
- $P(H) = \frac{13}{52}$
- $P(F \cap H) = \frac{3}{52}$
- $P(F \cup H) = P(F) + P(H) - P(F \cap H)$

Alternative Viewpoint

- $P(F) = \frac{12}{52}$
- $P(H) = \frac{13}{52}$
- $P(F \cap H) = \frac{3}{52}$
- $P(F \cup H) = P(F) + P(H) - P(F \cap H)$
- We can also do:

$$\frac{\binom{13}{4} \binom{4}{1} \binom{4}{2} * 4^3}{\binom{52}{5}}$$

Probability of Unions

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

Probability of Unions

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

- If A and B are **independent**, we have

$$P(A \cup B) = P(A) + P(B) - P(A) \cdot P(B)$$

Probability of Unions

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

- If A and B are **independent**, we have

$$P(A \cup B) = P(A) + P(B) - P(A) \cdot P(B)$$

- If A and B are **disjoint**, we have

$$P(A \cup B) = P(A) + P(B)$$

Probability of Unions of 3 Sets

$$\begin{aligned} P(A \cup B \cup C) &= P(A) + P(B) + P(C) \\ &\quad - P(A \cap B) - P(B \cap C) - P(A \cap C) \\ &\quad \quad + P(A \cap B \cap C) \end{aligned}$$

Probability of Unions of 3 Sets

$$\begin{aligned} P(A \cup B \cup C) &= P(A) + P(B) + P(C) \\ &\quad - P(A \cap B) - P(B \cap C) - P(A \cap C) \\ &\quad + P(A \cap B \cap C) \end{aligned}$$

- If A, B and C are **pairwise independent**, we have :

$$\begin{aligned} P(A \cup B \cup C) &= P(A) + P(B) + P(C) - P(A) \cdot P(B) - P(B) \cdot P(C) - \\ &\quad P(A) \cdot P(C) + P(A \cdot B \cdot C) \end{aligned}$$

Probability of Unions of 3 Sets

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) \\ - P(A \cap B) - P(B \cap C) - P(A \cap C) \\ + P(A \cap B \cap C)$$

- If A, B and C are **pairwise independent**, we have :

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A) \cdot P(B) - P(B) \cdot P(C) - \\ P(A) \cdot P(C) + P(A \cdot B \cdot C)$$

- If A, B and C are **pairwise disjoint** (so $A \cap B = A \cap C = B \cap C = \emptyset$, so clearly $A \cap B \cap C = \emptyset$), we have

$$P(A \cup B \cup C) = P(A) + P(B) + P(C)$$

Conditional Probability and Bayes' Law

Conditional Probability


- If **A occurs**, then is B
 - a) More likely?
 - b) Equally likely?
 - c) Less likely?

Conditional Probability


- If **A occurs**, then is B
 - a) More likely?
 - b) Equally likely?
 - c) Less likely?
- Any of these could happen, it depends on the relationship between A and B.

Conditional Probability


- If **A occurs**, then is B
 - a) More likely?
 - b) Equally likely?
 - c) Less likely?
- Any of these could happen, it depends on the relationship between A and B.



OR



OR



OR ...

Examples

- We roll two dice
 - Event $A = \text{“Sum of the dice } S \equiv 0 \pmod{4}\text{”}$
 - Note that $P(A) = \frac{9}{36} = \frac{1}{4}$, since we have **nine** rolls of the dice that sum to a multiple of 4:
 $(1, 3), (2, 2), (3, 1), (2, 6), (3, 5), (4, 4), (5, 3), (6, 2), (6, 6)$

Examples

- We roll two dice
 - Event A = “Sum of the dice $S \equiv 0 \pmod{4}$ ”
 - Note that $P(A) = \frac{9}{36} = \frac{1}{4}$, since we have **nine** rolls of the dice that sum to a multiple of 4:
 $(1, 3), (2, 2), (3, 1), (2, 6), (3, 5), (4, 4), (5, 3), (6, 2), (6, 6)$
 - Event B = “The first die comes up 3”
 - Note that $P(B) = \frac{6}{36} = \frac{1}{6}$

Examples

- We roll two dice
 - Event A = “Sum of the dice $S \equiv 0 \pmod{4}$ ”
 - Note that $P(A) = \frac{9}{36} = \frac{1}{4}$, since we have **nine** rolls of the dice that sum to a multiple of 4:
 $(1, 3), (2, 2), (3, 1), (2, 6), (3, 5), (4, 4), (5, 3), (6, 2), (6, 6)$
 - Event B = “The first die comes up 3”
 - Note that $P(B) = \frac{6}{36} = \frac{1}{6}$
- What is the probability of **A given B**?

Examples

- What is the probability of **A given B**?

Examples

- What is the probability of **A given B**?
 - Outcomes of A are (1, 3), (2, 2), (3, 1), (2, 6), (3, 5), (4, 4), (5, 3), (6, 2), (6, 6)
 - Outcomes of B are (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6)
 - Outcomes of rolling two dice: (1, 1), (1, 2), ..., (6, 5), (6, 6)

Examples

- What is the probability of **A given B**?
 - Outcomes of A are (1, 3), (2, 2), **(3, 1)**, (2, 6), **(3, 5)**, (4, 4), (5, 3), (6, 2), (6, 6)
 - Outcomes of B are **(3, 1)**, (3, 2), (3, 3), (3, 4), **(3, 5)**, (3, 6)
 - Outcomes of rolling two dice: (1, 1), (1, 2), ..., (6, 5), (6, 6)

Examples

- What is the probability of **A given B**?
 - Outcomes of A are (1, 3), (2, 2), **(3, 1)**, (2, 6), **(3, 5)**, (4, 4), (5, 3), (6, 2), (6, 6)
 - Outcomes of B are **(3, 1)**, (3, 2), (3, 3), (3, 4), **(3, 5)**, (3, 6)
 - Outcomes of rolling two dice: (1, 1), (1, 2), ..., (6, 5), (6, 6)
- As discussed, $P(A) = \frac{9}{36} = \frac{1}{4}$
- However, once B occurs, instead of 36 outcomes, we now have...

Examples

- What is the probability of **A given B**?
 - Outcomes of A are (1, 3), (2, 2), **(3, 1)**, (2, 6), **(3, 5)**, (4, 4), (5, 3), (6, 2), (6, 6)
 - Outcomes of B are **(3, 1)**, (3, 2), (3, 3), (3, 4), **(3, 5)**, (3, 6)
 - Outcomes of rolling two dice: (1, 1), (1, 2), ..., (6, 5), (6, 6)
- As discussed, $P(A) = \frac{9}{36} = \frac{1}{4}$
- However, once B occurs, instead of 36 outcomes, we now have... **6 outcomes.**

Examples

- What is the probability of **A given B**?
 - Outcomes of A are (1, 3), (2, 2), **(3, 1)**, (2, 6), **(3, 5)**, (4, 4), (5, 3), (6, 2), (6, 6)
 - Outcomes of B are **(3, 1)**, (3, 2), (3, 3), (3, 4), **(3, 5)**, (3, 6)
 - Outcomes of rolling two dice: (1, 1), (1, 2), ..., (6, 5), (6, 6)
- As discussed, $P(A) = \frac{9}{36} = \frac{1}{4}$
- However, once B occurs, instead of **36** outcomes, we now have... **6 outcomes**.
 - Only **2** of them are outcomes that correspond to A.

Examples

- What is the probability of **A given B**?
 - Outcomes of A are (1, 3), (2, 2), **(3, 1)**, (2, 6), **(3, 5)**, (4, 4), (5, 3), (6, 2), (6, 6)
 - Outcomes of B are **(3, 1)**, (3, 2), (3, 3), (3, 4), **(3, 5)**, (3, 6)
 - Outcomes of rolling two dice: (1, 1), (1, 2), ..., (6, 5), (6, 6)
- As discussed, $P(A) = \frac{9}{36} = \frac{1}{4}$
- However, once B occurs, instead of 36 outcomes, we now have... **6 outcomes**.
 - Only **2** of them are outcomes that correspond to A.
 - Therefore, the probability of **A given B** is $\frac{2}{6} = \frac{1}{3}$

Examples

- We once again two roll dice
 - Event A = “Sum of the dice is ≥ 8 ”
 - Event B = “ First die is 4”

Examples

- We once again two roll dice
 - Event A = “Sum of the dice is ≥ 8 ”
 - Event B = “ First die is 4”
- If B happens, what is your intuition about the probability of A?

Examples

- We once again two roll dice
 - Event A = “Sum of the dice is ≥ 8 ”
 - Event B = “ First die is 4”
- If B happens, what is your intuition about the probability of A?



Examples

- We once again two roll dice
 - Event A = “Sum of the dice is ≥ 8 ”
 - Event B = “ First die is 4”
- If B happens, what is your intuition about the probability of A?



Let's see if
your
intuition
was correct!

Examples

- We once again two roll dice
 - Event A = “Sum of the dice is ≥ 8 ” $P(A) = ?$ (work on it)
 - Event B = “ First die is 4”

Examples

- We once again two roll dice
 - Event A = “Sum of the dice is ≥ 8 ” $P(A) = \frac{15}{36} = \frac{5}{12}$
 - Event B = “First die is a 4”

Examples

- We once again two roll dice
 - Event A = “Sum of the dice is ≥ 8 ” $P(A) = \frac{15}{36} = \frac{5}{12}$
 - Event B = “First die is a 4” $P(B) = \frac{1}{6}$

Examples

- We once again two roll dice
 - Event A = “Sum of the dice is ≥ 8 ” $P(A) = \frac{15}{36} = \frac{5}{12}$
 - Event B = “First die is a 4” $P(B) = \frac{1}{6}$
- Prob of A given B = Prob second dice is 4, 5, or 6 = $\frac{3}{6} = \frac{1}{2} > \frac{5}{12}$

By just $\frac{1}{12}$...



Conditional Probability

- Let A, B be two events. The conditional probability of A *given* B , denoted $P(A | B)$ is defined as follows:

$$P(A | B) = \frac{P(A \cap B)}{P(B)}$$

Re-Thinking Independent Events

- **Alternative definition of independent events:** Two events A and B will be called marginally independent, or just independent for short, if and only if

$$P(A|B) = P(A)$$

Re-Thinking Independent Events

- **Alternative definition of independent events:** Two events A and B will be called marginally independent, or just independent for short, if and only if

$$P(A|B) = P(A)$$

- Applying the definition of $P(A|B)$ we have:
 - $\frac{P(A \cap B)}{P(B)} = P(A) \Rightarrow P(A \cap B) = P(A) \cdot P(B)$, which is a relationship we had reached **earlier** when discussing the joint probability.

Complex Probabilities

- Suppose that I have two dice: a six-sided one and a ten-sided one.

Complex Probabilities

- Suppose that I have two dice: a six-sided one and a ten-sided one.
- I pick either one of them with probability $\frac{1}{2}$ and roll it.
 - What's the probability that the die comes up 6? (work on this yourselves NOW)

Complex Probabilities

- Suppose that I have two dice: a six-sided one and a ten-sided one.
- I pick either one of them with probability $\frac{1}{2}$
 - What's the probability that the die comes up 6? (work on this yourselves **NOW**)

$$\begin{aligned}P(\text{Roll} = 6) &= P(\text{Roll} = 6, \text{Die} = 6) + P(\text{Roll} = 6, \text{Die} = 10) = \\&= P(\text{Roll} = 6 | \text{Die} = 6) \times P(\text{Die} = 6) + P(\text{Roll} = 6 | \text{Die} = 10) \times P(\text{Die} = 10) \\&= \\&= \frac{1}{6} \times \frac{1}{2} + \frac{1}{10} \times \frac{1}{2} = \frac{1}{12} + \frac{1}{20} = \frac{2}{15} \approx 0.1333 \dots\end{aligned}$$

Complex Probabilities

- Suppose that I have two dice: a six-sided one and a ten-sided one.

Complex Probabilities

- Suppose that I have two dice: a six-sided one and a ten-sided one.
- Now **we change the problem** so that we pick the ten-sided die with prob $\frac{5}{9}$ and the six-sided die with prob $\frac{4}{9}$.

Complex Probabilities

- Suppose that I have two dice: a six-sided one and a ten-sided one.
- Now **we change the problem** so that we pick the ten-sided die with prob $\frac{5}{9}$ and the six-sided die with prob $\frac{4}{9}$.
- **Intuitively**, will the probability that I come up with a 6...

Complex Probabilities

- Suppose that I have two dice: a six-sided one and a ten-sided one.
- Now **we change the problem** so that we pick the ten-sided die with prob $\frac{5}{9}$ and the six-sided die with prob $\frac{4}{9}$.
- **Intuitively**, will the probability that I come up with a 6...

Go up

Go down

Stay the
same

Complex Probabilities

- Suppose that I have two dice: a six-sided one and a ten-sided one.
- Now **we change the problem** so that we pick the ten-sided die with prob $\frac{5}{9}$ and the six-sided die with prob $\frac{4}{9}$.
- **Intuitively**, will the probability that I come up with a 6...

Go up

Go down

Stay the
same

Let's see if your
intuition was
correct!

Complex Probabilities

- Suppose that I have two dice: a six-sided one and a ten-sided one.
- Now **we change the problem** so that we pick the ten-sided die with prob $\frac{5}{9}$ and the six-sided die with prob $\frac{4}{9}$.
- What's the probability **that I come up with a 6?**

$$\begin{aligned} P(\text{Roll} = 6) &= P(\text{Roll} = 6, \text{Die} = 6) + P(\text{Roll} = 6, \text{Die} = 10) = \\ &= P(\text{Roll} = 6 | \text{Die} = 6) \times P(\text{Die} = 6) + P(\text{Roll} = 6, \text{Die} = 10) \times P(\text{Die} = 10) = \\ &= \frac{1}{6} \times \frac{4}{9} + \frac{1}{10} \times \frac{5}{9} = \frac{2}{27} + \frac{1}{18} = \frac{7}{54} \approx 0.130 < \mathbf{0.133} \end{aligned}$$

Bayes' Law

- Suppose A and B are events in a sample space Ω . Then, the following is an identity:

$$P(A|B) = P(B|A) \frac{P(A)}{P(B)}$$

known as **Bayes' Law**

Questions



- If $P(A|B) = P(A)$, is it the case that $P(B|A) = P(B)$?

Yes

No

Questions

- If $P(A|B) = P(A)$, is it the case that $P(B|A) = P(B)$?





- Substituting $P(A|B)$ with $P(A)$ in the formulation of Bayes' Law, we have:

$$\cancel{P(A)} = P(B | A) \cdot \frac{\cancel{P(A)}}{P(B)} \Rightarrow 1 = \frac{P(B|A)}{P(B)} \Rightarrow P(B|A) = P(B)$$

Questions

- If $P(A|B) = P(A)$, is it the case that $P(B|A) = P(B)$?



(A ind B) iff (B ind A)

- Substituting $P(A|B)$ with $P(A)$ in the formulation of Bayes' Law, we have:

$$\cancel{P(A)} = P(B | A) \cdot \frac{\cancel{P(A)}}{P(B)} \Rightarrow 1 = \frac{P(B|A)}{P(B)} \Rightarrow P(B|A) = P(B)$$

Questions

- If $P(B) = 0$, then is $P(A|B)$ also 0?

Yes

No

Questions

- If $P(B) = 0$, then is $P(A|B)$ also 0?

Yes

No

- It is **undefined**, since $P(A | B) = P(B | A) \cdot \frac{P(A)}{P(B)}$

STOP

RECORDING