

Optional Project Solutions

May 6, 2026

Problem 2a: Prop Logic

Problem 2a: Prop Logic

Give a formula on 5 variables that has exactly 3 satisfying assignments.

Problem 2a: Prop Logic

Give a formula on 5 variables that has exactly 3 satisfying assignments.

Imagine a truth table on 5 vars which the first three rows are T but the rest are F. The formula you get is

Problem 2a: Prop Logic

Give a formula on 5 variables that has exactly 3 satisfying assignments.

Imagine a truth table on 5 vars which the first three rows are T but the rest are F. The formula you get is

$$(\neg v \wedge \neg w \wedge \neg x \wedge \neg y \wedge \neg z)$$

$$\vee(\neg v \wedge \neg w \wedge \neg x \wedge \neg y \wedge z)$$

$$\vee(\neg v \wedge \neg w \wedge \neg x \wedge y \wedge \neg z)$$

Problem 2a: Prop Logic

Problem 2a: Prop Logic

Give a formula on 3 variables that has exactly 5 satisfying assignments.

Problem 2a: Prop Logic

Give a formula on 3 variables that has exactly 5 satisfying assignments.

Imagine a truth table on 3 vars which the first five rows are T but the rest are F. The formula you get is

Problem 2a: Prop Logic

Give a formula on 3 variables that has exactly 5 satisfying assignments.

Imagine a truth table on 3 vars which the first five rows are T but the rest are F. The formula you get is

$$(\neg x \wedge \neg y \wedge \neg z)$$

$$\vee(\neg x \wedge \neg y \wedge z)$$

$$\vee(\neg x \wedge y \wedge \neg z)$$

$$\vee(\neg x \wedge y \wedge z)$$

$$\vee(x \wedge \neg y \wedge \neg z)$$

Problem 2c: Prop Logic

Problem 2c: Prop Logic

Give an a, b such that there is NO formula on a variables that has exactly b satisfying assignments.

Problem 2c: Prop Logic

Give an a, b such that there is NO formula on a variables that has exactly b satisfying assignments.

Take $a = 3$ and $b = 2^a + 1 = 9$.

Problem 2d: Prop Logic

Problem 2d: Prop Logic

Either

Problem 2d: Prop Logic

Either

give an a, b such that there is NO formula on a variables that has exactly b satisfying assignments AND there is NO formula on b variables that has exactly a satisfying assignments.

Problem 2d: Prop Logic

Either

give an a, b such that there is NO formula on a variables that has exactly b satisfying assignments AND there is NO formula on b variables that has exactly a satisfying assignments.

OR show there is no such a, b .

Problem 2d: Prop Logic

Either

give an a, b such that there is NO formula on a variables that has exactly b satisfying assignments AND there is NO formula on b variables that has exactly a satisfying assignments.

OR show there is no such a, b .

There is no such a, b :

Problem 2d: Prop Logic

Either

give an a, b such that there is NO formula on a variables that has exactly b satisfying assignments AND there is NO formula on b variables that has exactly a satisfying assignments.

OR show there is no such a, b .

There is no such a, b :

You would need both $b \geq 2^a + 1$ and $a \geq 2^b + 1$.

Problem 2d: Prop Logic

Either

give an a, b such that there is NO formula on a variables that has exactly b satisfying assignments AND there is NO formula on b variables that has exactly a satisfying assignments.

OR show there is no such a, b .

There is no such a, b :

You would need both $b \geq 2^a + 1$ and $a \geq 2^b + 1$.

If both are true then

Problem 2d: Prop Logic

Either

give an a, b such that there is NO formula on a variables that has exactly b satisfying assignments AND there is NO formula on b variables that has exactly a satisfying assignments.

OR show there is no such a, b .

There is no such a, b :

You would need both $b \geq 2^a + 1$ and $a \geq 2^b + 1$.

If both are true then

$$b \geq 2^a + 1 \geq 2^{2^b+1} + 1$$

which is impossible.

Problem 3: Domains Question

Give a domain $\mathbb{D} \subseteq \mathbb{R}$ such that:

Problem 3: Domains Question

Give a domain $\mathbb{D} \subseteq \mathbb{R}$ such that:
 $(\exists L)(\forall x)[L \leq x]$. (L is for Lowest).

Problem 3: Domains Question

Give a domain $\mathbb{D} \subseteq \mathbb{R}$ such that:

$(\exists L)(\forall x)[L \leq x]$. (L is for Lowest).

$(\exists H)(\forall x)[U \geq x]$. (H is for Highest).

Problem 3: Domains Question

Give a domain $\mathbb{D} \subseteq \mathbb{R}$ such that:

$(\exists L)(\forall x)[L \leq x]$. (L is for Lowest).

$(\exists H)(\forall x)[U \geq x]$. (H is for Highest).

$(\forall x \neq H)(\exists y)x < y \wedge (\forall z)[\neg(x < z < y)]$.

(For every element $x \neq H$ there is a *successor*.)

Problem 3: Domains Question

Give a domain $\mathbb{D} \subseteq \mathbb{R}$ such that:

$(\exists L)(\forall x)[L \leq x]$. (L is for Lowest).

$(\exists H)(\forall x)[U \geq x]$. (H is for Highest).

$(\forall x \neq H)(\exists y)x < y \wedge (\forall z)[\neg(x < z < y)]$.

(For every element $x \neq H$ there is a *successor*.)

$(\forall x \neq L)(\exists y)y < x \wedge (\forall z)[\neg(y < z < x)]$.

(For every element $x \neq L$ there is a *predecessor*.)

Problem 3: Domains Answer

Problem 3: Domains Answer

Let D be the set

$$\left\{-1, -\frac{1}{2}, -\frac{1}{3}, \dots\right\} \cup \left\{\dots, \frac{1}{3}, \frac{1}{2}, 1\right\}.$$

Problem 3: Domains Answer

Let D be the set

$$\left\{-1, -\frac{1}{2}, -\frac{1}{3}, \dots\right\} \cup \left\{\dots, \frac{1}{3}, \frac{1}{2}, 1\right\}.$$

-1 is the lowest element

Problem 3: Domains Answer

Let D be the set

$$\left\{-1, -\frac{1}{2}, -\frac{1}{3}, \dots\right\} \cup \left\{\dots, \frac{1}{3}, \frac{1}{2}, 1\right\}.$$

-1 is the lowest element

1 is the highest element

Problem 3: Domains Answer

Let D be the set

$$\left\{-1, -\frac{1}{2}, -\frac{1}{3}, \dots\right\} \cup \left\{\dots, \frac{1}{3}, \frac{1}{2}, 1\right\}.$$

-1 is the lowest element

1 is the highest element

Every element except for -1 and 1 has a successor and a predecessor.

Problem 4: Constructive Induction

Let a_i be defined as follows:

Problem 4: Constructive Induction

Let a_i be defined as follows:

$$a_0 = 40 \quad a_1 = 17 \quad a_2 = 40$$

Problem 4: Constructive Induction

Let a_i be defined as follows:

$$a_0 = 40 \quad a_1 = 17 \quad a_2 = 40$$

$$(\forall n \geq 3)[a_n = 3a_{n-1} + 2a_{n-2} + a_{n-3} + 7].$$

Problem 4: Constructive Induction

Let a_i be defined as follows:

$$a_0 = 40 \quad a_1 = 17 \quad a_2 = 40$$

$$(\forall n \geq 3)[a_n = 3a_{n-1} + 2a_{n-2} + a_{n-3} + 7].$$

Use const ind to find b, m such that

Problem 4: Constructive Induction

Let a_i be defined as follows:

$$a_0 = 40 \quad a_1 = 17 \quad a_2 = 40$$

$$(\forall n \geq 3)[a_n = 3a_{n-1} + 2a_{n-2} + a_{n-3} + 7].$$

Use const ind to find b, m such that
for all n , $a_n \equiv b \pmod{m}$.

Base Case

IB: Need

Base Case

IB: Need

$$40 \equiv b \pmod{m}.$$

Base Case

IB: Need

$$40 \equiv b \pmod{m}.$$

$$17 \equiv b \pmod{m}.$$

Base Case

IB: Need

$$40 \equiv b \pmod{m}.$$

$$17 \equiv b \pmod{m}.$$

$$40 \equiv b \pmod{m}.$$

IH and IS

IH and IS

IH: For all $n' < n$, $a_{n'} \equiv b \pmod{m}$.

IH and IS

IH: For all $n' < n$, $a_{n'} \equiv b \pmod{m}$.

IS: All \equiv are mod m .

IH and IS

IH: For all $n' < n$, $a_{n'} \equiv b \pmod{m}$.

IS: All \equiv are mod m .

$$a_n = 3a_{n-1} + 2a_{n-2} + a_{n-3} + 7 \equiv 3b + 2b + b + 7 = 6b + 7$$

IH and IS

IH: For all $n' < n$, $a_{n'} \equiv b \pmod{m}$.

IS: All \equiv are mod m .

$$a_n = 3a_{n-1} + 2a_{n-2} + a_{n-3} + 7 \equiv 3b + 2b + b + 7 = 6b + 7$$

So we need

$$6b + 7 \equiv b$$

IH and IS

IH: For all $n' < n$, $a_{n'} \equiv b \pmod{m}$.

IS: All \equiv are mod m .

$$a_n = 3a_{n-1} + 2a_{n-2} + a_{n-3} + 7 \equiv 3b + 2b + b + 7 = 6b + 7$$

So we need

$6b + 7 \equiv b$ which is equivalent to

IH and IS

IH: For all $n' < n$, $a_{n'} \equiv b \pmod{m}$.

IS: All \equiv are mod m .

$$a_n = 3a_{n-1} + 2a_{n-2} + a_{n-3} + 7 \equiv 3b + 2b + b + 7 = 6b + 7$$

So we need

$6b + 7 \equiv b$ which is equivalent to

$$5b + 7 \equiv 0.$$

IH and IS

IH: For all $n' < n$, $a_{n'} \equiv b \pmod{m}$.

IS: All \equiv are mod m .

$$a_n = 3a_{n-1} + 2a_{n-2} + a_{n-3} + 7 \equiv 3b + 2b + b + 7 = 6b + 7$$

So we need

$6b + 7 \equiv b$ which is equivalent to

$$5b + 7 \equiv 0.$$

Next Slide.

What we Need

What we Need

Need a, b to satisfy

What we Need

Need a, b to satisfy

$$40 \equiv b \pmod{m}.$$

$$17 \equiv b \pmod{m}.$$

$$40 \equiv b \pmod{m}.$$

$$5b + 7 \equiv 0.$$

What we Need

Need a, b to satisfy

$$40 \equiv b \pmod{m}.$$

$$17 \equiv b \pmod{m}.$$

$$40 \equiv b \pmod{m}.$$

$$5b + 7 \equiv 0.$$

Note that we need 17, 40 are all equivalent to each other mod m .

What we Need

Need a, b to satisfy

$$40 \equiv b \pmod{m}.$$

$$17 \equiv b \pmod{m}.$$

$$40 \equiv b \pmod{m}.$$

$$5b + 7 \equiv 0.$$

Note that we need 17, 40 are all equivalent to each other mod m .

$$40 \equiv 17 \text{ so } 23 \equiv 0.$$

What we Need

Need a, b to satisfy

$$40 \equiv b \pmod{m}.$$

$$17 \equiv b \pmod{m}.$$

$$40 \equiv b \pmod{m}.$$

$$5b + 7 \equiv 0.$$

Note that we need 17, 40 are all equivalent to each other mod m .

$$40 \equiv 17 \text{ so } 23 \equiv 0.$$

Lets try $m = 23$.

What we Need

Need a, b to satisfy

$$40 \equiv b \pmod{m}.$$

$$17 \equiv b \pmod{m}.$$

$$40 \equiv b \pmod{m}.$$

$$5b + 7 \equiv 0.$$

Note that we need 17, 40 are all equivalent to each other mod m .

$$40 \equiv 17 \text{ so } 23 \equiv 0.$$

Lets try $m = 23$.

Next slide

Try $m = 23$

Try $m = 23$

$$40 \equiv b \pmod{23}.$$

$$17 \equiv b \pmod{23}.$$

$$40 \equiv b \pmod{23}.$$

$$5b + 7 \equiv 0 \pmod{23}.$$

Try $m = 23$

$$40 \equiv b \pmod{23}.$$

$$17 \equiv b \pmod{23}.$$

$$40 \equiv b \pmod{23}.$$

$$5b + 7 \equiv 0 \pmod{23}.$$

The first three equations force $b = 17$.

Try $m = 23$

$$40 \equiv b \pmod{23}.$$

$$17 \equiv b \pmod{23}.$$

$$40 \equiv b \pmod{23}.$$

$$5b + 7 \equiv 0 \pmod{23}.$$

The first three equations force $b = 17$.

We still need

Try $m = 23$

$$40 \equiv b \pmod{23}.$$

$$17 \equiv b \pmod{23}.$$

$$40 \equiv b \pmod{23}.$$

$$5b + 7 \equiv 0 \pmod{23}.$$

The first three equations force $b = 17$.

We still need

$$5b + 7 \equiv 0 \pmod{23}.$$

Try $m = 23$

$$40 \equiv b \pmod{23}.$$

$$17 \equiv b \pmod{23}.$$

$$40 \equiv b \pmod{23}.$$

$$5b + 7 \equiv 0 \pmod{23}.$$

The first three equations force $b = 17$.

We still need

$$5b + 7 \equiv 0 \pmod{23}.$$

Since $b = 17$:

Try $m = 23$

$$40 \equiv b \pmod{23}.$$

$$17 \equiv b \pmod{23}.$$

$$40 \equiv b \pmod{23}.$$

$$5b + 7 \equiv 0 \pmod{23}.$$

The first three equations force $b = 17$.

We still need

$$5b + 7 \equiv 0 \pmod{23}.$$

Since $b = 17$:

$$(5 \times 17) + 7 \equiv 0 \pmod{23}.$$

Try $m = 23$

$$40 \equiv b \pmod{23}.$$

$$17 \equiv b \pmod{23}.$$

$$40 \equiv b \pmod{23}.$$

$$5b + 7 \equiv 0 \pmod{23}.$$

The first three equations force $b = 17$.

We still need

$$5b + 7 \equiv 0 \pmod{23}.$$

Since $b = 17$:

$$(5 \times 17) + 7 \equiv 0 \pmod{23}.$$

So YES, $b = 17$ and $m = 23$.

$n^{1/7}$ Problem and Begin Solution

$n^{1/7}$ Problem and Begin Solution

$$n \in \mathbb{N}, n \geq 2.$$

$n^{1/7}$ Problem and Begin Solution

$n \in \mathbb{N}$, $n \geq 2$.

Let $n = p_1^{n_1} \cdots p_k^{n_k}$, where p_1, \dots, p_k are primes.

$n^{1/7}$ Problem and Begin Solution

$n \in \mathbb{N}$, $n \geq 2$.

Let $n = p_1^{n_1} \cdots p_k^{n_k}$, where p_1, \dots, p_k are primes.

Use unique factorization to show:

If $n^{1/7}$ is rational then $(\forall 1 \leq i \leq k)[n_i \equiv 0 \pmod{7}]$.

$n^{1/7}$ Problem and Begin Solution

$n \in \mathbb{N}$, $n \geq 2$.

Let $n = p_1^{n_1} \cdots p_k^{n_k}$, where p_1, \dots, p_k are primes.

Use unique factorization to show:

If $n^{1/7}$ is rational then $(\forall 1 \leq i \leq k)[n_i \equiv 0 \pmod{7}]$.

Assume there exists $a, b \in \mathbb{N}$ such that $n^{1/7} = \frac{a}{b}$.

$n^{1/7}$ Problem and Begin Solution

$n \in \mathbb{N}$, $n \geq 2$.

Let $n = p_1^{n_1} \cdots p_k^{n_k}$, where p_1, \dots, p_k are primes.

Use unique factorization to show:

If $n^{1/7}$ is rational then $(\forall 1 \leq i \leq k)[n_i \equiv 0 \pmod{7}]$.

Assume there exists $a, b \in \mathbb{N}$ such that $n^{1/7} = \frac{a}{b}$.

$$(p_1^{n_1} \cdots p_k^{n_k})^{1/7} = \frac{a}{b}$$

$n^{1/7}$ Problem and Begin Solution

$n \in \mathbb{N}$, $n \geq 2$.

Let $n = p_1^{n_1} \cdots p_k^{n_k}$, where p_1, \dots, p_k are primes.

Use unique factorization to show:

If $n^{1/7}$ is rational then $(\forall 1 \leq i \leq k)[n_i \equiv 0 \pmod{7}]$.

Assume there exists $a, b \in \mathbb{N}$ such that $n^{1/7} = \frac{a}{b}$.

$$(p_1^{n_1} \cdots p_k^{n_k})^{1/7} = \frac{a}{b}$$

$$p_1^{n_1} \cdots p_k^{n_k} = \frac{a^7}{b^7}$$

$n^{1/7}$ Problem and Begin Solution

$n \in \mathbb{N}$, $n \geq 2$.

Let $n = p_1^{n_1} \cdots p_k^{n_k}$, where p_1, \dots, p_k are primes.

Use unique factorization to show:

If $n^{1/7}$ is rational then $(\forall 1 \leq i \leq k)[n_i \equiv 0 \pmod{7}]$.

Assume there exists $a, b \in \mathbb{N}$ such that $n^{1/7} = \frac{a}{b}$.

$$(p_1^{n_1} \cdots p_k^{n_k})^{1/7} = \frac{a}{b}$$

$$p_1^{n_1} \cdots p_k^{n_k} = \frac{a^7}{b^7}$$

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$n^{1/7}$ Problem and Begin Solution

$n \in \mathbb{N}$, $n \geq 2$.

Let $n = p_1^{n_1} \cdots p_k^{n_k}$, where p_1, \dots, p_k are primes.

Use unique factorization to show:

If $n^{1/7}$ is rational then $(\forall 1 \leq i \leq k)[n_i \equiv 0 \pmod{7}]$.

Assume there exists $a, b \in \mathbb{N}$ such that $n^{1/7} = \frac{a}{b}$.

$$(p_1^{n_1} \cdots p_k^{n_k})^{1/7} = \frac{a}{b}$$

$$p_1^{n_1} \cdots p_k^{n_k} = \frac{a^7}{b^7}$$

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

Next slide for exciting conclusion.

$n^{1/7}$ Solution

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$$a = p_1^{a_1} \cdots p_k^{a_k}$$

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$$a = p_1^{a_1} \cdots p_k^{a_k}$$

$$\text{hence } a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$$a = p_1^{a_1} \cdots p_k^{a_k}$$

$$b = p_1^{b_1} \cdots p_k^{b_k}$$

hence $a^7 = p_1^{7a_1} \cdots p_k^{7a_k}$.

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$$a = p_1^{a_1} \cdots p_k^{a_k}$$

$$b = p_1^{b_1} \cdots p_k^{b_k}$$

hence $a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$

hence

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$$a = p_1^{a_1} \cdots p_k^{a_k}$$

$$b = p_1^{b_1} \cdots p_k^{b_k}$$

$$\text{hence } a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$$\text{hence } b^7 = p_1^{7b_1} \cdots p_k^{7b_k}.$$

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$$a = p_1^{a_1} \cdots p_k^{a_k} \quad \text{hence } a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$$b = p_1^{b_1} \cdots p_k^{b_k} \quad \text{hence } b^7 = p_1^{7b_1} \cdots p_k^{7b_k}.$$

$$(p_1^{7b_1} \cdots p_k^{7b_k})(p_1^{n_1} \cdots p_k^{n_k}) = a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$$a = p_1^{a_1} \cdots p_k^{a_k} \quad \text{hence } a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$$b = p_1^{b_1} \cdots p_k^{b_k} \quad \text{hence } b^7 = p_1^{7b_1} \cdots p_k^{7b_k}.$$

$$(p_1^{7b_1} \cdots p_k^{7b_k})(p_1^{n_1} \cdots p_k^{n_k}) = a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$$p_1^{7b_1+n_1} \cdots p_k^{7b_k+n_k} = p_1^{7a_1} \cdots p_k^{7a_k}$$

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$$a = p_1^{a_1} \cdots p_k^{a_k} \quad \text{hence } a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$$b = p_1^{b_1} \cdots p_k^{b_k} \quad \text{hence } b^7 = p_1^{7b_1} \cdots p_k^{7b_k}.$$

$$(p_1^{7b_1} \cdots p_k^{7b_k})(p_1^{n_1} \cdots p_k^{n_k}) = a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$$p_1^{7b_1+n_1} \cdots p_k^{7b_k+n_k} = p_1^{7a_1} \cdots p_k^{7a_k}$$

Let $1 \leq i \leq k$. By Unique Factorization, p_i has to appear the same number of times on both sides.

$n^{1/7}$ Solution

$$b^7 p_1^{n_1} \cdots p_k^{n_k} = a^7.$$

$$a = p_1^{a_1} \cdots p_k^{a_k} \quad \text{hence } a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$$b = p_1^{b_1} \cdots p_k^{b_k} \quad \text{hence } b^7 = p_1^{7b_1} \cdots p_k^{7b_k}.$$

$$(p_1^{7b_1} \cdots p_k^{7b_k})(p_1^{n_1} \cdots p_k^{n_k}) = a^7 = p_1^{7a_1} \cdots p_k^{7a_k}.$$

$$p_1^{7b_1+n_1} \cdots p_k^{7b_k+n_k} = p_1^{7a_1} \cdots p_k^{7a_k}$$

Let $1 \leq i \leq k$. By Unique Factorization, p_i has to appear the same number of times on both sides.

Hence

$$7b_i + n_i = 7a_i$$

$$n_i = y(a_i - b_i) \equiv 0 \pmod{7}.$$

Darling's Lunch: Combinatorics

Darling's Lunch: Combinatorics

Bill makes his Darling lunch.

Darling's Lunch: Combinatorics

Bill makes his Darling lunch.

S sandwiches, F fruits, D desserts.

Darling's Lunch: Combinatorics

Bill makes his Darling lunch.

S sandwiches, F fruits, D desserts.

Lunch: s diff sandwich's, f diff fruits, and d deserts

Darling's Lunch: Combinatorics

Bill makes his Darling lunch.

S sandwiches, F fruits, D desserts.

Lunch: s diff sandwich's, f diff fruits, and d deserts

How many ways can Darling have lunch?

Darling's Lunch: Combinatorics

Bill makes his Darling lunch.

S sandwiches, F fruits, D desserts.

Lunch: s diff sandwich's, f diff fruits, and d deserts

How many ways can Darling have lunch? $\binom{S}{s} \binom{F}{f} D^d$.

Darling's Lunch: Combinatorics

Bill makes his Darling lunch.

S sandwiches, F fruits, D desserts.

Lunch: s diff sandwich's, f diff fruits, and d deserts

How many ways can Darling have lunch? $\binom{S}{s} \binom{F}{f} D^d$.

PBJ, BLUEBERRY, BLUEBERRY PIE: Darling does not want ALL THREE.

Darling's Lunch: Combinatorics

Bill makes his Darling lunch.

S sandwiches, F fruits, D desserts.

Lunch: s diff sandwich's, f diff fruits, and d deserts

How many ways can Darling have lunch? $\binom{S}{s} \binom{F}{f} D^d$.

PBJ, BLUEBERRY, BLUEBERRY PIE: Darling does not want ALL THREE.

How many ways can Darling have lunch?

Darling's Lunch: Combinatorics

Bill makes his Darling lunch.

S sandwiches, F fruits, D desserts.

Lunch: s diff sandwich's, f diff fruits, and d deserts

How many ways can Darling have lunch? $\binom{S}{s} \binom{F}{f} D^d$.

PBJ, BLUEBERRY, BLUEBERRY PIE: Darling does not want ALL THREE.

How many ways can Darling have lunch?

How many lunches **do** have all 3?: $\binom{S-1}{s-2} \binom{F-1}{f-1} D^{d-1}$.

Darling's Lunch: Combinatorics

Bill makes his Darling lunch.

S sandwiches, F fruits, D desserts.

Lunch: s diff sandwich's, f diff fruits, and d deserts

How many ways can Darling have lunch? $\binom{S}{s} \binom{F}{f} D^d$.

PBJ, BLUEBERRY, BLUEBERRY PIE: Darling does not want ALL THREE.

How many ways can Darling have lunch?

How many lunches **do** have all 3?: $\binom{S-1}{s-2} \binom{F-1}{f-1} D^{d-1}$.

So answer is $\binom{S}{s} \binom{F}{f} D^d - \binom{S-1}{s-1} \binom{F-1}{f-1} D^{d-1}$.

Number of Solutions

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that
 $x_1 + x_2 + x_3 + x_4 + x_5 = 1000$

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Four Bars.

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Four Bars.

1st space has at least 1.

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Four Bars.

1st space has at least 1.

2nd space has at least 2.

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Four Bars.

1st space has at least 1.

2nd space has at least 2.

3rd space has at least 3.

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Four Bars.

1st space has at least 1.

2nd space has at least 2.

3rd space has at least 3.

4th space has at least 4

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Four Bars.

1st space has at least 1.

2nd space has at least 2.

3rd space has at least 3.

4th space has at least 4

5th space has at least 5

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Four Bars.

1st space has at least 1.

2nd space has at least 2.

3rd space has at least 3.

4th space has at least 4

5th space has at least 5

So $1 + 2 + 3 + 4 + 5 = 15$ stars already placed.

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Four Bars.

1st space has at least 1.

2nd space has at least 2.

3rd space has at least 3.

4th space has at least 4

5th space has at least 5

So $1 + 2 + 3 + 4 + 5 = 15$ stars already placed.

Our problem: How many ways to arrange 4 bars and

$1000 - 15 = 985$ stars.

Number of Solutions

How many $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{N}^5$ such that

$$x_1 + x_2 + x_3 + x_4 + x_5 = 1000$$

and $x_1 \geq 1$, $x_2 \geq 2$, $x_3 \geq 3$, $x_4 \geq 4$, and $x_5 \geq 5$.

Stars and Bars!

Four Bars.

1st space has at least 1.

2nd space has at least 2.

3rd space has at least 3.

4th space has at least 4

5th space has at least 5

So $1 + 2 + 3 + 4 + 5 = 15$ stars already placed.

Our problem: How many ways to arrange 4 bars and

$1000 - 15 = 985$ stars.

$\frac{989}{4!985!}$. This is also $\binom{985}{4}$. Discuss.

Strange Poker-two 3-of-a-kind

Strange Poker-two 3-of-a-kind

Rank in $\{1, \dots, r\}$, suit in $\{1, \dots, s\}$, 6-card hands.

Strange Poker-two 3-of-a-kind

Rank in $\{1, \dots, r\}$, suit in $\{1, \dots, s\}$, 6-card hands.

1) Probability of getting two 3-of-a-kinds:

Strange Poker-two 3-of-a-kind

Rank in $\{1, \dots, r\}$, suit in $\{1, \dots, s\}$, 6-card hands.

1) Probability of getting two 3-of-a-kinds:

Two 3-of-a-kind: Pick 2 ranks $\binom{r}{2}$,

Strange Poker-two 3-of-a-kind

Rank in $\{1, \dots, r\}$, suit in $\{1, \dots, s\}$, 6-card hands.

1) Probability of getting two 3-of-a-kinds:

Two 3-of-a-kind: Pick 2 ranks $\binom{r}{2}$,

For the lower rank pick three suits: $\binom{s}{3}$.

Strange Poker-two 3-of-a-kind

Rank in $\{1, \dots, r\}$, suit in $\{1, \dots, s\}$, 6-card hands.

1) Probability of getting two 3-of-a-kinds:

Two 3-of-a-kind: Pick 2 ranks $\binom{r}{2}$,

For the lower rank pick three suits: $\binom{s}{3}$.

For the higher rank pick three suits: $\binom{s}{3}$.

Strange Poker-two 3-of-a-kind

Rank in $\{1, \dots, r\}$, suit in $\{1, \dots, s\}$, 6-card hands.

1) Probability of getting two 3-of-a-kinds:

Two 3-of-a-kind: Pick 2 ranks $\binom{r}{2}$,

For the lower rank pick three suits: $\binom{s}{3}$.

For the higher rank pick three suits: $\binom{s}{3}$.

Prob of getting three of a kind:

Strange Poker-two 3-of-a-kind

Rank in $\{1, \dots, r\}$, suit in $\{1, \dots, s\}$, 6-card hands.

1) Probability of getting two 3-of-a-kinds:

Two 3-of-a-kind: Pick 2 ranks $\binom{r}{2}$,

For the lower rank pick three suits: $\binom{s}{3}$.

For the higher rank pick three suits: $\binom{s}{3}$.

Prob of getting three of a kind:

$$\frac{\binom{r}{2} \binom{s}{3}^2}{\binom{rs}{6}}.$$

Strange Poker 4-of-a-kind

Strange Poker 4-of-a-kind

2) What is the probability of getting four-of-a-kinds?

Strange Poker 4-of-a-kind

2) What is the probability of getting four-of-a-kinds?

Four of a kind:

Strange Poker 4-of-a-kind

2) What is the probability of getting four-of-a-kinds?

Four of a kind: Pick a ranks r .

Strange Poker 4-of-a-kind

2) What is the probability of getting four-of-a-kinds?

Four of a kind: Pick a ranks r . Pick 4 suits $\binom{5}{4}$.

Strange Poker 4-of-a-kind

2) What is the probability of getting four-of-a-kinds?

Four of a kind: Pick a rank r . Pick 4 suits $\binom{5}{4}$.

Pick 2 more cards but they can't be of rank r .

Strange Poker 4-of-a-kind

2) What is the probability of getting four-of-a-kinds?

Four of a kind: Pick a rank r . Pick 4 suits $\binom{5}{4}$.

Pick 2 more cards but they can't be of rank r . $\binom{(r-1)5}{2}$.

Strange Poker 4-of-a-kind

2) What is the probability of getting four-of-a-kinds?

Four of a kind: Pick a ranks r . Pick 4 suits $\binom{s}{4}$.

Pick 2 more cards but they can't be of rank r . $\binom{(r-1)s}{2}$.

Prob of getting 4-of-a-kind:

Strange Poker 4-of-a-kind

2) What is the probability of getting four-of-a-kinds?

Four of a kind: Pick a ranks r . Pick 4 suits $\binom{s}{4}$.

Pick 2 more cards but they can't be of rank r . $\binom{(r-1)s}{2}$.

Prob of getting 4-of-a-kind:

$$\frac{r \binom{s}{4} \binom{(r-1)s}{2}}{\binom{rs}{6}}.$$

Strange Poker 4-of-a-kind

2) What is the probability of getting four-of-a-kinds?

Four of a kind: Pick a rank r . Pick 4 suits $\binom{s}{4}$.

Pick 2 more cards but they can't be of rank r . $\binom{(r-1)s}{2}$.

Prob of getting 4-of-a-kind:

$$\frac{r \binom{s}{4} \binom{(r-1)s}{2}}{\binom{rs}{6}}.$$

If you didn't know to not use the same rank, I will still mark it right.