

**Homework 11, Morally Due 12:30PM, Tue Apr 28 2026**

1. (0 points) What is your name?

**GO TO NEXT PAGE**

2. (35 points)

- (a) (10 points) Show that  $\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}$  with a combinatorial proof. That is, show that the Right Hand Side solves the problem *how many ways can I pick  $k$  Ramsey Theorists from a set of  $n$  Ramsey Theorists.*  
(You will use this in the next part.)
- (b) (15 points) Show that, for all  $1, b \geq 2$ ,  
 $R(a, b) \leq \binom{a+b-2}{a-1}$ .
- (c) (10 points) Use the result  $R(a, b) \leq \binom{a+b-2}{a-1}$  to get an asymptotic upper bound on  $R(k)$  that is LESS THAN  $2^{2k-1}$ .

**GO TO NEXT PAGE**

3. (30 points)

**Def** Let  $G = (V, E)$  be a graph.  $D \subseteq V$  is a *dominating set (DS)* if

$$(\forall v \in V)[v \in D \vee (\exists y \in D)[(x, y) \in E]].$$

Every graph has a DS of size  $n$ :  $D = V$ . We do better!

You will prove: *There exists a function  $\alpha$  such that,*

*a) For all  $d \in \mathbf{N}$ ,  $0 < \alpha(d) < 1$ .*

*b) The function  $\alpha(d)$  is DECREASING.*

*c) For every graph with min degree  $\geq d$  there is a dominating set of size  $\leq \alpha(d)n$ .*

On the next page we will state the theorem with the function  $\alpha$  and sketch the proof. YOU will fill in the details and find a function  $\alpha$  that works. We guide this with a series of embedded questions.

**Thm** There exists a function  $\alpha$  such that the following hold:

- $\alpha$  maps  $\mathbb{N}$  to  $[0, 1]$ .
- $\alpha$  is strictly DECREASING. Hence, for all  $d \geq 1$ ,  $\alpha(d) < 1$ .
- If  $G = (V, E)$  is a graph on  $n$  vertices with min degree  $\geq d$  then  $G$  has a dominating set of size  $\leq \alpha(d)n$ .

**Pf** Let  $p$  be a probability to be determined by YOU later.

Pick  $X \subseteq V$  as follows: For every  $v \in V$  choose  $v$  with probability  $p$ .

- (a) (0 points but you need it for later) What is  $E(|X|)$ ? It will be a function of  $n, p$ .
- (b) (0 points but you need it for later) Let  $Y \subseteq V - X$  be the vertices that DO NOT have an edge to an element of  $X$ . Formally

$$Y = \{y \in V - X : (\forall x \in X)[(x, y) \notin E]\}.$$

Give an upper bound on  $E(|Y|)$ . It will be a function of  $n, d, p$ . Note that  $X \cup Y$  is a dominating set. We later pick  $p$  so that  $|X \cup Y|$  is small.

- (c) (0 points but you need it for later) What is  $E(|X \cup Y|)$ ? (Hint: This is very easy by the linearity of expectation.)
- (d) (30 points) Pick  $p$  to make  $E(|X \cup Y|)$  smaller than  $n$  (Hint: Find an upper bound on  $E(|X \cup Y|)$  and minimize that bound. Use that  $(1 - p)^d \leq e^{-dp}$ .)

State and prove a theorem of the form:

**Thm** If  $G = (V, E)$  is a graph on  $n$  vertices with min degree  $\geq d$  then  $G$  has a dominating set of size  $\leq \alpha(d)n$ .

**GO TO NEXT PAGE**

4. (35 points)

- (a) (0 points but you need to do it for the rest of the problem) Write a program that will, given  $n$ , generate a random 2-coloring of  $\binom{[n]}{2}$  by, for each edge, color it RED with prob  $\frac{1}{2}$  and BLUE with prob  $\frac{1}{2}$ .
- (b) (0 points but you need to do it for the rest of the problem) Write a program that will, given  $n$  and a 2-coloring of  $\binom{[n]}{2}$ , count how many mono  $K_3$ 's there are.
- (c) (0 points but you need to do it for the rest of the problem) This problem just puts the two programs together. Write a program that will, given  $n$ , generate a random 2-coloring of  $\binom{[n]}{2}$  by, for each edge, color it RED with prob  $\frac{1}{2}$  and BLUE with prob  $\frac{1}{2}$  and then OUTPUTS the number of mono  $K_3$ 's.
- (d) (0 points but you need to do it for the rest of the problem) This problem mostly uses the last program. Write a program that will, given  $n$ , run the last program 100 times and gather up the number-of-triangles. Then just output the MIN and the MAX.
- (e) (35 points) Write a program that outputs a table with
- $n$  going from 4 to 20
  - Min number of mono  $K_3$ 's from last program.
  - Max number of mono  $K_3$ 's from last program.
  - How much bigger is the Max from what we are guaranteed by the theorems in class. We call this *over*.

Here is the format of the table (I made up the numbers and only give the first 3 rows.

$n$	min	max	over
4	1	2	2
5	1	2	1
6	2	8	3