

# BILL, RECORD LECTURE!!!!

BILL RECORD LECTURE!!!

# Well Quasi Orders: Kruskal Tree Theorem and the Graph Minor Theorem

Exposition by William Gasarch

February 18, 2026

# The Kruskal Tree Theorem

# Subgraph Ordering for Graphs & Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **subgraph** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

# Subgraph Ordering for Graphs & Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **subgraph** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

# Subgraph Ordering for Graphs & Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **subgraph** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

- Remove vertices (and the incident edge)

- Remove edges (the vertices stay)

# Subgraph Ordering for Graphs & Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **subgraph** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

Remove edges (the vertices stay)

We denote this  $H \preceq_{\text{sub}} G$ . It is an ordering.

# Subgraph Ordering for Graphs & Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **subgraph** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

Remove edges (the vertices stay)

We denote this  $H \preceq_{\text{sub}} G$ . It is an ordering.

**Def** A **colored graph** is a graph where you also have assigned to each vertex a color. There is no restriction on the coloring.

# Subgraph Ordering for Graphs & Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **subgraph** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

Remove edges (the vertices stay)

We denote this  $H \preceq_{\text{sub}} G$ . It is an ordering.

**Def** A **colored graph** is a graph where you also have assigned to each vertex a color. There is no restriction on the coloring.

**Def** Let  $H$  and  $G$  be colored graphs.  $H$  is a **subgraph** of  $G$  is defined exactly as for non-colored graphs.

# Subgraph Ordering for Graphs & Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **subgraph** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

Remove edges (the vertices stay)

We denote this  $H \preceq_{\text{sub}} G$ . It is an ordering.

**Def** A **colored graph** is a graph where you also have assigned to each vertex a color. There is no restriction on the coloring.

**Def** Let  $H$  and  $G$  be colored graphs.  $H$  is a **subgraph** of  $G$  is defined exactly as for non-colored graphs.

We denote this  $H \preceq_{\text{c-sub}} G$ . It is an ordering.

# Subgraph Ordering for Graphs & Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **subgraph** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

Remove edges (the vertices stay)

We denote this  $H \preceq_{\text{sub}} G$ . It is an ordering.

**Def** A **colored graph** is a graph where you also have assigned to each vertex a color. There is no restriction on the coloring.

**Def** Let  $H$  and  $G$  be colored graphs.  $H$  is a **subgraph** of  $G$  is defined exactly as for non-colored graphs.

We denote this  $H \preceq_{\text{c-sub}} G$ . It is an ordering.

Do some examples on the board.

# The Minor Ordering for Graphs and Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **minor** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

# The Minor Ordering for Graphs and Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **minor** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

# The Minor Ordering for Graphs and Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **minor** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

- Remove vertices (and the incident edge)

- Remove edges (the vertices stay)

# The Minor Ordering for Graphs and Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **minor** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

- Remove vertices (and the incident edge)

- Remove edges (the vertices stay)

- Remove an edge and merging its endpoints.

# The Minor Ordering for Graphs and Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **minor** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

- Remove vertices (and the incident edge)

- Remove edges (the vertices stay)

- Remove an edge and merging its endpoints.

We denote this  $H \preceq_m G$ . It is an ordering.

# The Minor Ordering for Graphs and Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **minor** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

Remove edges (the vertices stay)

Remove an edge and merging its endpoints.

We denote this  $H \preceq_m G$ . It is an ordering.

**Def** Let  $H$  and  $G$  be colored graphs.  $H$  is a **minor** of  $G$  is defined exactly as for non-colored graphs.

# The Minor Ordering for Graphs and Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **minor** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

Remove edges (the vertices stay)

Remove an edge and merging its endpoints.

We denote this  $H \preceq_m G$ . It is an ordering.

**Def** Let  $H$  and  $G$  be colored graphs.  $H$  is a **minor** of  $G$  is defined exactly as for non-colored graphs.

We denote this  $H \preceq_{c-m} G$ . It is an ordering.

# The Minor Ordering for Graphs and Colored Graphs

**Def** Let  $H$  and  $G$  be graphs.  $H$  is a **minor** of  $G$  if you can obtain  $H$  by taking  $G$  and do the following operations as many times as you want and in any order that you want:

Remove vertices (and the incident edge)

Remove edges (the vertices stay)

Remove an edge and merging its endpoints.

We denote this  $H \preceq_m G$ . It is an ordering.

**Def** Let  $H$  and  $G$  be colored graphs.  $H$  is a **minor** of  $G$  is defined exactly as for non-colored graphs.

We denote this  $H \preceq_{c-m} G$ . It is an ordering.

Do some examples on the board.

# The Kruskal Tree Theorem

# The Kruskal Tree Theorem

In the 1930's Andrew Vazsonyi conjectured that the set of trees under the minor ordering was a wqo.

# The Kruskal Tree Theorem

In the 1930's Andrew Vazsonyi conjectured that the set of trees under the minor ordering was a wqo.

In 1960 Joe Kruskal proved this for his PhD Thesis. Formally:

# The Kruskal Tree Theorem

In the 1930's Andrew Vazsonyi conjectured that the set of trees under the minor ordering was a wqo.

In 1960 Joe Kruskal proved this for his PhD Thesis. Formally:

**Thm** Let  $X$  be the set of trees. Then  $(X, \preceq_m)$  is a wqo.

# The Kruskal Tree Theorem

In the 1930's Andrew Vazsonyi conjectured that the set of trees under the minor ordering was a wqo.

In 1960 Joe Kruskal proved this for his PhD Thesis. Formally:

**Thm** Let  $X$  be the set of trees. Then  $(X, \preceq_m)$  is a wqo.

In 1963 Nash-Williams obtained a simpler proof.

# The Kruskal Tree Theorem

In the 1930's Andrew Vazsonyi conjectured that the set of trees under the minor ordering was a wqo.

In 1960 Joe Kruskal proved this for his PhD Thesis. Formally:

**Thm** Let  $X$  be the set of trees. Then  $(X, \preceq_m)$  is a wqo.

In 1963 Nash-Williams obtained a simpler proof.

We present the proof of Nash-Williams.

# The Kruskal Tree Theorem

In the 1930's Andrew Vazsonyi conjectured that the set of trees under the minor ordering was a wqo.

In 1960 Joe Kruskal proved this for his PhD Thesis. Formally:

**Thm** Let  $X$  be the set of trees. Then  $(X, \preceq_m)$  is a wqo.

In 1963 Nash-Williams obtained a simpler proof.

We present the proof of Nash-Williams.

Joe Kruskal was happy that there was a simpler proof.

# The Kruskal Tree Theorem

In the 1930's Andrew Vazsonyi conjectured that the set of trees under the minor ordering was a wqo.

In 1960 Joe Kruskal proved this for his PhD Thesis. Formally:

**Thm** Let  $X$  be the set of trees. Then  $(X, \preceq_m)$  is a wqo.

In 1963 Nash-Williams obtained a simpler proof.

We present the proof of Nash-Williams.

Joe Kruskal was happy that there was a simpler proof.

Clyde Kruskal learned the proof of the KTT from Bill Gasarch

# The Kruskal Tree Theorem

In the 1930's Andrew Vazsonyi conjectured that the set of trees under the minor ordering was a wqo.

In 1960 Joe Kruskal proved this for his PhD Thesis. Formally:

**Thm** Let  $X$  be the set of trees. Then  $(X, \preceq_m)$  is a wqo.

In 1963 Nash-Williams obtained a simpler proof.

We present the proof of Nash-Williams.

Joe Kruskal was happy that there was a simpler proof.

Clyde Kruskal learned the proof of the KTT from Bill Gasarch

Martin Kruskal learned the proof of the KTT from Bill Gasarch

# The Kruskal Tree Theorem

In the 1930's Andrew Vazsonyi conjectured that the set of trees under the minor ordering was a wqo.

In 1960 Joe Kruskal proved this for his PhD Thesis. Formally:

**Thm** Let  $X$  be the set of trees. Then  $(X, \preceq_m)$  is a wqo.

In 1963 Nash-Williams obtained a simpler proof.

We present the proof of Nash-Williams.

Joe Kruskal was happy that there was a simpler proof.

Clyde Kruskal learned the proof of the KTT from Bill Gasarch

Martin Kruskal learned the proof of the KTT from Bill Gasarch

The Kruskal Family should talk with each other more.

# Sketch of Proof of the KTT

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots,$

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.  
Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.  
 $T_i$  is the smallest  $i$ th element of a bad sequence that begins  
 $T_1, \dots, T_{i-1}$ .

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.

$T_i$  is the smallest  $i$ th element of a bad sequence that begins  $T_1, \dots, T_{i-1}$ .

$U_i$  is the **set** of trees you get if you remove the root  $r_i$  of  $T_i$ .

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.

$T_i$  is the smallest  $i$ th element of a bad sequence that begins  $T_1, \dots, T_{i-1}$ .

$U_i$  is the **set** of trees you get if you remove the root  $r_i$  of  $T_i$ .

$XX$  is union of all trees in all the  $U_i$ . ROOTS is the set of roots.

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.

$T_i$  is the smallest  $i$ th element of a bad sequence that begins  $T_1, \dots, T_{i-1}$ .

$U_i$  is the **set** of trees you get if you remove the root  $r_i$  of  $T_i$ .

$XX$  is union of all trees in all the  $U_i$ . ROOTS is the set of roots.

Prove  $XX$  is a wqo (this is the hard part).

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.

$T_i$  is the smallest  $i$ th element of a bad sequence that begins  $T_1, \dots, T_{i-1}$ .

$U_i$  is the **set** of trees you get if you remove the root  $r_i$  of  $T_i$ .

$XX$  is union of all trees in all the  $U_i$ . ROOTS is the set of roots.

Prove  $XX$  is a wqo (this is the hard part).

By prior Theorems,  $2^{\text{fin}XX} \times \text{ROOTS}$  is a wqo.

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.

$T_i$  is the smallest  $i$ th element of a bad sequence that begins  $T_1, \dots, T_{i-1}$ .

$U_i$  is the **set** of trees you get if you remove the root  $r_i$  of  $T_i$ .

$XX$  is union of all trees in all the  $U_i$ . ROOTS is the set of roots.

Prove  $XX$  is a wqo (this is the hard part).

By prior Theorems,  $2^{\text{fin}XX} \times \text{ROOTS}$  is a wqo.

Look at sequence  $(U_1, r_1), (U_2, r_2), (U_3, r_3), \dots$

## Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.

$T_i$  is the smallest  $i$ th element of a bad sequence that begins  $T_1, \dots, T_{i-1}$ .

$U_i$  is the **set** of trees you get if you remove the root  $r_i$  of  $T_i$ .

$XX$  is union of all trees in all the  $U_i$ . ROOTS is the set of roots.

Prove  $XX$  is a wqo (this is the hard part).

By prior Theorems,  $2^{\text{fin}XX} \times \text{ROOTS}$  is a wqo.

Look at sequence  $(U_1, r_1), (U_2, r_2), (U_3, r_3), \dots$

There exists  $i \leq j$  such that  $(U_i, r_i) \preceq (U_j, r_j)$ .

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.

$T_i$  is the smallest  $i$ th element of a bad sequence that begins  $T_1, \dots, T_{i-1}$ .

$U_i$  is the **set** of trees you get if you remove the root  $r_i$  of  $T_i$ .

$XX$  is union of all trees in all the  $U_i$ . ROOTS is the set of roots.

Prove  $XX$  is a wqo (this is the hard part).

By prior Theorems,  $2^{\text{fin}XX} \times \text{ROOTS}$  is a wqo.

Look at sequence  $(U_1, r_1), (U_2, r_2), (U_3, r_3), \dots$

There exists  $i \leq j$  such that  $(U_i, r_i) \preceq (U_j, r_j)$ .

Can use this to get  $T_i \preceq T_j$ .

# Sketch of Proof of the KTT

Assume, BWOC, that  $(X, \preceq_m)$  is not a wqo.  $\exists \geq 1$  bad seq.

Create min bad seq  $T_1, T_2, \dots$ . Size is number of vertices.

$T_i$  is the smallest  $i$ th element of a bad sequence that begins  $T_1, \dots, T_{i-1}$ .

$U_i$  is the **set** of trees you get if you remove the root  $r_i$  of  $T_i$ .

$XX$  is union of all trees in all the  $U_i$ . ROOTS is the set of roots.

Prove  $XX$  is a wqo (this is the hard part).

By prior Theorems,  $2^{\text{fin}XX} \times \text{ROOTS}$  is a wqo.

Look at sequence  $(U_1, r_1), (U_2, r_2), (U_3, r_3), \dots$

There exists  $i \leq j$  such that  $(U_i, r_i) \preceq (U_j, r_j)$ .

Can use this to get  $T_i \preceq T_j$ . We are Done!

# The Color Version of Kruskal Tree Theorem

**Thm** Let  $n \in \mathbb{N}$ . Let  $X$  be the set of  $n$ -colored trees. Then  $(X, \preceq_{c-m})$  is a wqo.

# The Color Version of Kruskal Tree Theorem

**Thm** Let  $n \in \mathbb{N}$ . Let  $X$  be the set of  $n$ -colored trees. Then  $(X, \preceq_{c-m})$  is a wqo.

Proof is similar to the proof of **Kruskal Tree Theorem**

# The Graph Minor Theorem

# The Graph Minor Theorem

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

He soon after conjectured that the set of **graphs** under minor was a wqo (his result would follow from that).

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

He soon after conjectured that the set of **graphs** under minor was a wqo (his result would follow from that).

This was proven by Robertson and Seymour.

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

He soon after conjectured that the set of **graphs** under minor was a wqo (his result would follow from that).

This was proven by Robertson and Seymour.

It is called **The Graph Minor Theorem** and abbreviated **GMT**.

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

He soon after conjectured that the set of **graphs** under minor was a wqo (his result would follow from that).

This was proven by Robertson and Seymour.

It is called **The Graph Minor Theorem** and abbreviated **GMT**.

The GMT was proved in a series of 20 papers which are over 500 pages written between 1983 and 2004.

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

He soon after conjectured that the set of **graphs** under minor was a wqo (his result would follow from that).

This was proven by Robertson and Seymour.

It is called **The Graph Minor Theorem** and abbreviated **GMT**.

The GMT was proved in a series of 20 papers which are over 500 pages written between 1983 and 2004.

The proof has not been simplified. I will not be presenting it.

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

He soon after conjectured that the set of **graphs** under minor was a wqo (his result would follow from that).

This was proven by Robertson and Seymour.

It is called **The Graph Minor Theorem** and abbreviated **GMT**.

The GMT was proved in a series of 20 papers which are over 500 pages written between 1983 and 2004.

The proof has not been simplified. I will not be presenting it.

**GMT** Let  $X$  be the set of graphs. Then  $(X, \preceq_m)$  is a wqo.

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

He soon after conjectured that the set of **graphs** under minor was a wqo (his result would follow from that).

This was proven by Robertson and Seymour.

It is called **The Graph Minor Theorem** and abbreviated **GMT**.

The GMT was proved in a series of 20 papers which are over 500 pages written between 1983 and 2004.

The proof has not been simplified. I will not be presenting it.

**GMT** Let  $X$  be the set of graphs. Then  $(X, \preceq_m)$  is a wqo.

I will be happy when there is a simpler proof.

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

He soon after conjectured that the set of **graphs** under minor was a wqo (his result would follow from that).

This was proven by Robertson and Seymour.

It is called **The Graph Minor Theorem** and abbreviated **GMT**.

The GMT was proved in a series of 20 papers which are over 500 pages written between 1983 and 2004.

The proof has not been simplified. I will not be presenting it.

**GMT** Let  $X$  be the set of graphs. Then  $(X, \preceq_m)$  is a wqo.

I will be happy when there is a simpler proof.

I don't think that will happen in my lifetime but...

# The Graph Minor Theorem

In the 1937 Klaus Wagner showed that a graph is planar IFF it does not have  $K_{3,3}$  or  $K_5$  as a minor.

He soon after conjectured that the set of **graphs** under minor was a wqo (his result would follow from that).

This was proven by Robertson and Seymour.

It is called **The Graph Minor Theorem** and abbreviated **GMT**.

The GMT was proved in a series of 20 papers which are over 500 pages written between 1983 and 2004.

The proof has not been simplified. I will not be presenting it.

**GMT** Let  $X$  be the set of graphs. Then  $(X, \preceq_m)$  is a wqo.

I will be happy when there is a simpler proof.

I don't think that will happen in my lifetime but...

It may happen in Ilya Hajiaghayi's lifetime since he might do it.

# An Open Question

# An Open Question

Why did **Joe Kruskal** get his name on his theorem but

# An Open Question

Why did **Joe Kruskal** get his name on his theorem but

**Robertson and Seymour** did not get their name on their theorem.