

# BILL, RECORD LECTURE!!!!

BILL RECORD LECTURE!!!

# When Does a 2-Coloring Yield a Mono Unit Square?

**Exposition by William Gasarch**

May 8, 2026

# Credit Where Credit is Due!

The main theorem of these slides is due to Paul Erdős, Ronald Graham, Peter Montgomery, Bruce L. Rothchild, Joel Spencer, Ernst G. Straus.

# Credit Where Credit is Due!

The main theorem of these slides is due to Paul Erdős, Ronald Graham, Peter Montgomery, Bruce L. Rothchild, Joel Spencer, Ernst G. Straus.

**[Journal of Combinatorial Theory \(A\), Vol. 14, 341-363, 1973](https://www.cs.umd.edu/~gasarch/TOPICS/eramsey/eramseyOne.pdf)**

<https://www.cs.umd.edu/~gasarch/TOPICS/eramsey/eramseyOne.pdf>

# What is a Mono Unit Square?

**Def** a **Mono Unit Square** is a unit square with all four corners the same color.

# What is a Mono Unit Square?

**Def** a **Mono Unit Square** is a unit square with all four corners the same color.

**Def** A coloring is **proper** if there is no unit square.

# What is a Mono Unit Square?

**Def** a **Mono Unit Square** is a unit square with all four corners the same color.

**Def** A coloring is **proper** if there is no unit square.

**Question** Is there a proper 2-coloring of  $\mathbb{R}^2$ ?

# What is a Mono Unit Square?

**Def** a **Mono Unit Square** is a unit square with all four corners the same color.

**Def** A coloring is **proper** if there is no unit square.

**Question** Is there a proper 2-coloring of  $\mathbb{R}^2$ ?

**Answer** Yes. We leave this for an exercise.

# What About Higher Dimensions?

# What About Higher Dimensions?

Here is all that is known:

# What About Higher Dimensions?

Here is all that is known:

- ▶ There is a proper 2-col of  $\mathbb{R}^2$ .

# What About Higher Dimensions?

Here is all that is known:

- ▶ There is a proper 2-col of  $\mathbb{R}^2$ .
- ▶ There is no proper 2-col of  $\mathbb{R}^4$ .

# What About Higher Dimensions?

Here is all that is known:

- ▶ There is a proper 2-col of  $\mathbb{R}^2$ .
- ▶ There is no proper 2-col of  $\mathbb{R}^4$ .

The problem is open in  $\mathbb{R}^3$ .

# What About Higher Dimensions?

Here is all that is known:

- ▶ There is a proper 2-col of  $\mathbb{R}^2$ .
- ▶ There is no proper 2-col of  $\mathbb{R}^4$ .

The problem is open in  $\mathbb{R}^3$ .

**Darling** That's too bad. We live in  $\mathbb{R}^3$ .

# We Will Prove...

# We Will Prove...

In this slide packet we will show that

# We Will Prove...

In this slide packet we will show that

For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a Mono Unit Square.

# We Will Prove...

In this slide packet we will show that

For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a Mono Unit Square.

In a later slide packet we will show

# We Will Prove...

In this slide packet we will show that

For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a Mono Unit Square.

In a later slide packet we will show

For all  $\text{COL}: \mathbb{R}^6 \rightarrow [2]$  there exists a Mono Unit Square.

# We Will Prove...

In this slide packet we will show that

For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a Mono Unit Square.

In a later slide packet we will show

For all  $\text{COL}: \mathbb{R}^6 \rightarrow [2]$  there exists a Mono Unit Square.

For all  $\text{COL}: \mathbb{R}^5 \rightarrow [2]$  there exists a Mono Unit Square.

# We Will Prove...

In this slide packet we will show that

For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a Mono Unit Square.

In a later slide packet we will show

For all  $\text{COL}: \mathbb{R}^6 \rightarrow [2]$  there exists a Mono Unit Square.

For all  $\text{COL}: \mathbb{R}^5 \rightarrow [2]$  there exists a Mono Unit Square.

The  $\mathbb{R}^5$  result is really an observation about the  $\mathbb{R}^6$  proof.

# We Will Prove...

In this slide packet we will show that

For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a Mono Unit Square.

In a later slide packet we will show

For all  $\text{COL}: \mathbb{R}^6 \rightarrow [2]$  there exists a Mono Unit Square.

For all  $\text{COL}: \mathbb{R}^5 \rightarrow [2]$  there exists a Mono Unit Square.

The  $\mathbb{R}^5$  result is really an observation about the  $\mathbb{R}^6$  proof.

We will also have comments about the proof of:

## We Will Prove...

In this slide packet we will show that

For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a Mono Unit Square.

In a later slide packet we will show

For all  $\text{COL}: \mathbb{R}^6 \rightarrow [2]$  there exists a Mono Unit Square.

For all  $\text{COL}: \mathbb{R}^5 \rightarrow [2]$  there exists a Mono Unit Square.

The  $\mathbb{R}^5$  result is really an observation about the  $\mathbb{R}^6$  proof.

We will also have comments about the proof of:

For all  $\text{COL}: \mathbb{R}^4 \rightarrow [2]$  there exists a Mono Unit Square.

# Vectors We Need

## Notation

# Vectors We Need

## Notation

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, 0\right) \quad e_2 = \left(0, \frac{1}{\sqrt{2}}, 0\right) \quad e_3 = \left(0, 0, \frac{1}{\sqrt{2}}\right).$$

# Vectors We Need

## Notation

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, 0\right)$$

$$e_2 = \left(0, \frac{1}{\sqrt{2}}, 0\right)$$

$$e_3 = \left(0, 0, \frac{1}{\sqrt{2}}\right).$$

$$E = \{e_1, e_2, e_3\}.$$

# Vectors We Need

## Notation

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, 0\right) \quad e_2 = \left(0, \frac{1}{\sqrt{2}}, 0\right) \quad e_3 = \left(0, 0, \frac{1}{\sqrt{2}}\right).$$

$$E = \{e_1, e_2, e_3\}.$$

$$f_1 = \left(\frac{1}{\sqrt{2}}, 0, 0, \dots, 0\right) \text{ 9 coordinates,}$$

$$f_2 = \left(0, \frac{1}{\sqrt{2}}, 0, \dots, 0\right) \quad \dots \quad f_9 = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

# Vectors We Need

## Notation

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, 0\right) \quad e_2 = \left(0, \frac{1}{\sqrt{2}}, 0\right) \quad e_3 = \left(0, 0, \frac{1}{\sqrt{2}}\right).$$

$$E = \{e_1, e_2, e_3\}.$$

$$f_1 = \left(\frac{1}{\sqrt{2}}, 0, 0, \dots, 0\right) \text{ 9 coordinates,}$$

$$f_2 = \left(0, \frac{1}{\sqrt{2}}, 0, \dots, 0\right) \quad \dots \quad f_9 = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

$$F = \{f_1, \dots, f_9\}.$$

# Vectors We Need

## Notation

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, 0\right) \quad e_2 = \left(0, \frac{1}{\sqrt{2}}, 0\right) \quad e_3 = \left(0, 0, \frac{1}{\sqrt{2}}\right).$$

$$E = \{e_1, e_2, e_3\}.$$

$$f_1 = \left(\frac{1}{\sqrt{2}}, 0, 0, \dots, 0\right) \text{ 9 coordinates,}$$

$$f_2 = \left(0, \frac{1}{\sqrt{2}}, 0, \dots, 0\right) \quad \dots \quad f_9 = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

$$F = \{f_1, \dots, f_9\}.$$

**Notation** If  $1 \leq i \leq 3$  and  $1 \leq k \leq 9$  then  $e_i f_k \in \mathbb{R}^{12}$  is the concat of the coordinates.

# Vectors We Need

## Notation

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, 0\right) \quad e_2 = \left(0, \frac{1}{\sqrt{2}}, 0\right) \quad e_3 = \left(0, 0, \frac{1}{\sqrt{2}}\right).$$

$$E = \{e_1, e_2, e_3\}.$$

$$f_1 = \left(\frac{1}{\sqrt{2}}, 0, 0, \dots, 0\right) \text{ 9 coordinates,}$$

$$f_2 = \left(0, \frac{1}{\sqrt{2}}, 0, \dots, 0\right) \quad \dots \quad f_9 = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

$$F = \{f_1, \dots, f_9\}.$$

**Notation** If  $1 \leq i \leq 3$  and  $1 \leq k \leq 9$  then  $e_i f_k \in \mathbb{R}^{12}$  is the concat of the coordinates.

## Example

$$e_2 f_7 = \left(0, \frac{1}{\sqrt{2}}, 0, 0, 0, 0, 0, 0, 0, 0, \frac{1}{\sqrt{2}}, 0, 0\right).$$

# Lemma Left To Reader

**Lemma** Let  $1 \leq i < j \leq 3$  and  $1 \leq k < \ell \leq 9$ .

# Lemma Left To Reader

**Lemma** Let  $1 \leq i < j \leq 3$  and  $1 \leq k < \ell \leq 9$ .

Then the following points form a unit square as shown:

$$e_i f_\ell \quad e_j f_\ell$$

$$e_i f_k \quad e_j f_k$$

# The $\mathbb{R}^{12}$ Theorem

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

Let  $\text{COL}': F \rightarrow [8]$  be defined as

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

Let  $\text{COL}': F \rightarrow [8]$  be defined as

$$\text{COL}'(f_k) = \text{COL}(e_1 f_k)\text{COL}(e_2 f_k)\text{COL}(e_3 f_k).$$

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

Let  $\text{COL}': F \rightarrow [8]$  be defined as

$$\text{COL}'(f_k) = \text{COL}(e_1 f_k)\text{COL}(e_2 f_k)\text{COL}(e_3 f_k).$$

There must be two  $f_k$ 's that map to the same triple-of-colors.

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

Let  $\text{COL}': F \rightarrow [8]$  be defined as

$$\text{COL}'(f_k) = \text{COL}(e_1 f_k)\text{COL}(e_2 f_k)\text{COL}(e_3 f_k).$$

There must be two  $f_k$ 's that map to the same triple-of-colors.

We assume  $f_4$  and  $f_7$  map to the same color.

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

Let  $\text{COL}': F \rightarrow [8]$  be defined as

$$\text{COL}'(f_k) = \text{COL}(e_1 f_k) \text{COL}(e_2 f_k) \text{COL}(e_3 f_k).$$

There must be two  $f_k$ 's that map to the same triple-of-colors.

We assume  $f_4$  and  $f_7$  map to the same color.

The triple-of-colors must have two of the same. We assume the triple is RRB.

Hence

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

Let  $\text{COL}': F \rightarrow [8]$  be defined as

$$\text{COL}'(f_k) = \text{COL}(e_1 f_k)\text{COL}(e_2 f_k)\text{COL}(e_3 f_k).$$

There must be two  $f_k$ 's that map to the same triple-of-colors.

We assume  $f_4$  and  $f_7$  map to the same color.

The triple-of-colors must have two of the same. We assume the triple is RRB.

Hence

$$\text{COL}'(f_4) = \text{COL}(e_1 f_4)\text{COL}(e_2 f_4)\text{COL}(e_3 f_4) = RRB.$$

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

Let  $\text{COL}': F \rightarrow [8]$  be defined as

$$\text{COL}'(f_k) = \text{COL}(e_1 f_k)\text{COL}(e_2 f_k)\text{COL}(e_3 f_k).$$

There must be two  $f_k$ 's that map to the same triple-of-colors.

We assume  $f_4$  and  $f_7$  map to the same color.

The triple-of-colors must have two of the same. We assume the triple is  $RRB$ .

Hence

$$\text{COL}'(f_4) = \text{COL}(e_1 f_4)\text{COL}(e_2 f_4)\text{COL}(e_3 f_4) = RRB.$$

$$\text{COL}'(f_7) = \text{COL}(e_1 f_7)\text{COL}(e_2 f_7)\text{COL}(e_3 f_7) = RRB.$$

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

Let  $\text{COL}': F \rightarrow [8]$  be defined as

$$\text{COL}'(f_k) = \text{COL}(e_1 f_k)\text{COL}(e_2 f_k)\text{COL}(e_3 f_k).$$

There must be two  $f_k$ 's that map to the same triple-of-colors.

We assume  $f_4$  and  $f_7$  map to the same color.

The triple-of-colors must have two of the same. We assume the triple is RRB.

Hence

$$\text{COL}'(f_4) = \text{COL}(e_1 f_4)\text{COL}(e_2 f_4)\text{COL}(e_3 f_4) = RRB.$$

$$\text{COL}'(f_7) = \text{COL}(e_1 f_7)\text{COL}(e_2 f_7)\text{COL}(e_3 f_7) = RRB.$$

Hence the following are the same color:

$$e_2 f_4 \quad e_2 f_7$$

$$e_1 f_7 \quad e_1 f_4$$

# The $\mathbb{R}^{12}$ Theorem

**Thm** For all  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$  there exists a mono unit square.

Let  $\text{COL}: \mathbb{R}^{12} \rightarrow [2]$

Let  $\text{COL}': F \rightarrow [8]$  be defined as

$$\text{COL}'(f_k) = \text{COL}(e_1 f_k)\text{COL}(e_2 f_k)\text{COL}(e_3 f_k).$$

There must be two  $f_k$ 's that map to the same triple-of-colors.

We assume  $f_4$  and  $f_7$  map to the same color.

The triple-of-colors must have two of the same. We assume the triple is RRB.

Hence

$$\text{COL}'(f_4) = \text{COL}(e_1 f_4)\text{COL}(e_2 f_4)\text{COL}(e_3 f_4) = RRB.$$

$$\text{COL}'(f_7) = \text{COL}(e_1 f_7)\text{COL}(e_2 f_7)\text{COL}(e_3 f_7) = RRB.$$

Hence the following are the same color:

$$e_2 f_4 \quad e_2 f_7$$

$$e_1 f_7 \quad e_1 f_4$$

# The $\mathbb{R}^{12}$ Theorem: More Colors

# The $\mathbb{R}^{12}$ Theorem: More Colors

The following you should be able to prove.

# The $\mathbb{R}^{12}$ Theorem: More Colors

The following you should be able to prove.

# The $\mathbb{R}^{12}$ Theorem: More Colors

The following you should be able to prove.

**Thm** For all  $c$ , there exists  $n$  such that, for all  $\text{COL}: \mathbb{R}^n \rightarrow [c]$  there exists a mono unit square.

# The $\mathbb{R}^{12}$ Theorem: More Dimension

# The $\mathbb{R}^{12}$ Theorem: More Dimension

Def

# The $\mathbb{R}^{12}$ Theorem: More Dimension

## Def

A **1-dim unit cube** is congruent to the 2 points  $\{0, 1\}$ .

# The $\mathbb{R}^{12}$ Theorem: More Dimension

## Def

A **1-dim unit cube** is congruent to the 2 points  $\{0, 1\}$ .

A **2-dim unit cube** is congruent to the  $2^2$  points  $\{0, 1\} \times \{0, 1\}$ .

# The $\mathbb{R}^{12}$ Theorem: More Dimension

## Def

A **1-dim unit cube** is congruent to the 2 points  $\{0, 1\}$ .

A **2-dim unit cube** is congruent to the  $2^2$  points  $\{0, 1\} \times \{0, 1\}$ .

A **3-dim unit cube** is congruent to the  $2^3$  points  $\{0, 1\}^3$ .

# The $\mathbb{R}^{12}$ Theorem: More Dimension

## Def

A **1-dim unit cube** is congruent to the 2 points  $\{0, 1\}$ .

A **2-dim unit cube** is congruent to the  $2^2$  points  $\{0, 1\} \times \{0, 1\}$ .

A **3-dim unit cube** is congruent to the  $2^3$  points  $\{0, 1\}^3$ .

An  **$m$ -dim unit cube** is congruent to the  $2^m$  points  $\{0, 1\}^m$ .

# The $\mathbb{R}^{12}$ Theorem: More Dimension

## Def

A **1-dim unit cube** is congruent to the 2 points  $\{0, 1\}$ .

A **2-dim unit cube** is congruent to the  $2^2$  points  $\{0, 1\} \times \{0, 1\}$ .

A **3-dim unit cube** is congruent to the  $2^3$  points  $\{0, 1\}^3$ .

An  **$m$ -dim unit cube** is congruent to the  $2^m$  points  $\{0, 1\}^m$ .

**Def** Given a coloring  $\text{COL}: \mathbb{R}^n \rightarrow [c]$ , a **mono unit  $m$ -cube** is a set of  $2^m$  points, all the same color, that form a mono  $m$ -cube

# The $\mathbb{R}^{12}$ Theorem: More Dimension

## Def

A **1-dim unit cube** is congruent to the 2 points  $\{0, 1\}$ .

A **2-dim unit cube** is congruent to the  $2^2$  points  $\{0, 1\} \times \{0, 1\}$ .

A **3-dim unit cube** is congruent to the  $2^3$  points  $\{0, 1\}^3$ .

An  **$m$ -dim unit cube** is congruent to the  $2^m$  points  $\{0, 1\}^m$ .

**Def** Given a coloring  $\text{COL}: \mathbb{R}^n \rightarrow [c]$ , a **mono unit  $m$ -cube** is a set of  $2^m$  points, all the same color, that form a mono  $m$ -cube

You should be able to prove the following.

# The $\mathbb{R}^{12}$ Theorem: More Dimension

## Def

A **1-dim unit cube** is congruent to the 2 points  $\{0, 1\}$ .

A **2-dim unit cube** is congruent to the  $2^2$  points  $\{0, 1\} \times \{0, 1\}$ .

A **3-dim unit cube** is congruent to the  $2^3$  points  $\{0, 1\}^3$ .

An  **$m$ -dim unit cube** is congruent to the  $2^m$  points  $\{0, 1\}^m$ .

**Def** Given a coloring  $\text{COL}: \mathbb{R}^n \rightarrow [c]$ , a **mono unit  $m$ -cube** is a set of  $2^m$  points, all the same color, that form a mono  $m$ -cube

You should be able to prove the following.

**Thm** For all  $c, m$ , there exists  $n$  such that, for all  $\text{COL}: \mathbb{R}^n \rightarrow [c]$  there exists a mono unit  $m$ -cube.

# Ramsey Sets

# Ramsey Sets

**Def** Let  $X \subseteq \mathbb{R}^m$ .  $X$  is a **Ramsey Set** if,

# Ramsey Sets

**Def** Let  $X \subseteq \mathbb{R}^m$ .  $X$  is a **Ramsey Set** if, for all  $c$ , there exists  $n$  such that

# Ramsey Sets

**Def** Let  $X \subseteq \mathbb{R}^m$ .  $X$  is a **Ramsey Set** if,  
for all  $c$ , there exists  $n$  such that  
for all  $\text{COL}: \mathbb{R}^n \rightarrow [c]$  there exists a mono copy of  $X$ .

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

For all COL:  $\mathbb{R}^{c+1} \rightarrow [c]$  there is a mono  $X$ :

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

For all COL:  $\mathbb{R}^{c+1} \rightarrow [c]$  there is a mono  $X$ :

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, \dots, 0\right), \dots, e_{c+1} = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

For all COL:  $\mathbb{R}^{c+1} \rightarrow [c]$  there is a mono  $X$ :

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, \dots, 0\right), \dots, e_{c+1} = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

Two of them are the same color and are a copy of  $X$ .

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

For all  $\text{COL}: \mathbb{R}^{c+1} \rightarrow [c]$  there is a mono  $X$ :

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, \dots, 0\right), \dots, e_{c+1} = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

Two of them are the same color and are a copy of  $X$ .

**Note** We wanted an  $n$  such that  $\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there would be a mono  $X$ .

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

For all  $\text{COL}: \mathbb{R}^{c+1} \rightarrow [c]$  there is a mono  $X$ :

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, \dots, 0\right), \dots, e_{c+1} = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

Two of them are the same color and are a copy of  $X$ .

**Note** We wanted an  $n$  such that  $\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there would be a mono  $X$ .

e got a finite set of points in  $\mathbb{R}^n$  such that just coloring those points you get a mono  $X$ .

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

For all  $\text{COL}: \mathbb{R}^{c+1} \rightarrow [c]$  there is a mono  $X$ :

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, \dots, 0\right), \dots, e_{c+1} = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

Two of them are the same color and are a copy of  $X$ .

**Note** We wanted an  $n$  such that  $\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there would be a mono  $X$ .

e got a finite set of points in  $\mathbb{R}^n$  such that just coloring those points you get a mono  $X$ .

**Theorem** (AC) Let  $X \subseteq \mathbb{R}^m$ , finite. Assume  $\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there exists a mono  $X$ .

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

For all COL:  $\mathbb{R}^{c+1} \rightarrow [c]$  there is a mono  $X$ :

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, \dots, 0\right), \dots, e_{c+1} = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

Two of them are the same color and are a copy of  $X$ .

**Note** We wanted an  $n$  such that  $\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there would be a mono  $X$ .

e got a finite set of points in  $\mathbb{R}^n$  such that just coloring those points you get a mono  $X$ .

**Theorem** (AC) Let  $X \subseteq \mathbb{R}^m$ , finite. Assume  $\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there exists a mono  $X$ .

Then there is a finite set  $Z \subseteq \mathbb{R}^n$  such that

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

For all  $\text{COL}: \mathbb{R}^{c+1} \rightarrow [c]$  there is a mono  $X$ :

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, \dots, 0\right), \dots, e_{c+1} = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

Two of them are the same color and are a copy of  $X$ .

**Note** We wanted an  $n$  such that  $\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there would be a mono  $X$ .

e got a finite set of points in  $\mathbb{R}^n$  such that just coloring those points you get a mono  $X$ .

**Theorem** (AC) Let  $X \subseteq \mathbb{R}^m$ , finite. Assume

$\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there exists a mono  $X$ .

Then there is a finite set  $Z \subseteq \mathbb{R}^n$  such that

$\forall \text{COL}: Z \rightarrow [c]$  there exists a mono  $X$ .

# Ramsey Sets: Only need a finite subseq of $\mathbb{R}^n$

## Examples

$X = \{0\}, \{1\} \subseteq \mathbb{R}^1$  is Ramsey.

For all COL:  $\mathbb{R}^{c+1} \rightarrow [c]$  there is a mono  $X$ :

$$e_1 = \left(\frac{1}{\sqrt{2}}, 0, \dots, 0\right), \dots, e_{c+1} = \left(0, \dots, 0, \frac{1}{\sqrt{2}}\right).$$

Two of them are the same color and are a copy of  $X$ .

**Note** We wanted an  $n$  such that  $\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there would be a mono  $X$ .

e got a finite set of points in  $\mathbb{R}^n$  such that just coloring those points you get a mono  $X$ .

**Theorem** (AC) Let  $X \subseteq \mathbb{R}^m$ , finite. Assume  $\forall \text{COL}: \mathbb{R}^n \rightarrow [c]$  there exists a mono  $X$ .

Then there is a finite set  $Z \subseteq \mathbb{R}^n$  such that

$\forall \text{COL}: Z \rightarrow [c]$  there exists a mono  $X$ .

We won't prove this.

# We Showed That

# We Showed That

We showed that the unit square is a Ramsey set.

# We Showed That

We showed that the unit square is a Ramsey set.

We showed that, for all  $m$ , the unit  $m$ -cube is Ramsey.

# Obtaining More Ramsey Sets

# Obtaining More Ramsey Sets

1) If  $x \in \mathbb{R}^{m_1}$  and  $y \in \mathbb{R}^{m_2}$  then  $x * y \in \mathbb{R}^{m_1+m_2}$  is formed by concatenating the coordinates.

# Obtaining More Ramsey Sets

1) If  $x \in \mathbb{R}^{m_1}$  and  $y \in \mathbb{R}^{m_2}$  then  $x * y \in \mathbb{R}^{m_1+m_2}$  is formed by concatenating the coordinates.

2) Let  $X \subseteq \mathbb{R}^{m_1}$  and  $Y \subseteq \mathbb{R}^{m_2}$ . Then  
 $X * Y = \{x * y : x \in X \wedge y \in Y\}$ .

# Obtaining More Ramsey Sets

1) If  $x \in \mathbb{R}^{m_1}$  and  $y \in \mathbb{R}^{m_2}$  then  $x * y \in \mathbb{R}^{m_1+m_2}$  is formed by concatenating the coordinates.

2) Let  $X \subseteq \mathbb{R}^{m_1}$  and  $Y \subseteq \mathbb{R}^{m_2}$ . Then  
 $X * Y = \{x * y : x \in X \wedge y \in Y\}$ .

The unit square is the product of  $\{0, 1\}$  and  $\{0, 1\}$ .

# Obtaining More Ramsey Sets

# Obtaining More Ramsey Sets

We showed the mono Unit Square is Ramsey.

# Obtaining More Ramsey Sets

We showed the mono Unit Square is Ramsey.

The unit square is the product of  $\{0, 1\}$  and  $\{0, 1\}$ .

# Obtaining More Ramsey Sets

We showed the mono Unit Square is Ramsey.

The unit square is the product of  $\{0, 1\}$  and  $\{0, 1\}$ .

Our proof was a scaled down version of the proof of this theorem:

# Obtaining More Ramsey Sets

We showed the mono Unit Square is Ramsey.

The unit square is the product of  $\{0, 1\}$  and  $\{0, 1\}$ .

Our proof was a scaled down version of the proof of this theorem:

**Thm** If  $X$  and  $Y$  are Ramsey then  $X * Y$  is Ramsey.