

AI Project-First Assignment

Counterfeit Coins Problem: Setting

Counterfeit Coins Problem: Setting

(We may consider this later.)

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

What is known?

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

What is known?

Very few exact values are known.

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

What is known?

Very few exact values are known.

Some asymptotic values are known.

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

What is known?

Very few exact values are known.

Some asymptotic values are known.

There does not seem to be any math structure to use.

Counterfeit Coins Problem: Project

Counterfeit Coins Problem: Project

Lets use AI on it!

Counterfeit Coins Problem: Project

Lets use AI on it!

How?

Counterfeit Coins Problem: Project

Lets use AI on it!

How?

Set it up as a game. We look at $c = 2$.

Counterfeit Coins Problem: Project

Lets use AI on it!

How?

Set it up as a game. We look at $c = 2$.

Weighter picks two sets of x coins to balance.

Counterfeit Coins Problem: Project

Lets use AI on it!

How?

Set it up as a game. We look at $c = 2$.

Weighter picks two sets of x coins to balance.

Spoiler gives an answer that is consistent with having 2 counterfeits.

Counterfeit Coins Problem: Project

Lets use AI on it!

How?

Set it up as a game. We look at $c = 2$.

Weighter picks two sets of x coins to balance.

Spoiler gives an answer that is consistent with having 2 counterfeits.

Repeat this until Weighter knows what the counterfeit coins are.

Counterfeit Coins Problem: Project

Lets use AI on it!

How?

Set it up as a game. We look at $c = 2$.

Weighter picks two sets of x coins to balance.

Spoiler gives an answer that is consistent with having 2 counterfeits.

Repeat this until Weighter knows what the counterfeit coins are.

Weighter's goal is to minimize the number of weightings.

Counterfeit Coins Problem: Project

Lets use AI on it!

How?

Set it up as a game. We look at $c = 2$.

Weighter picks two sets of x coins to balance.

Spoiler gives an answer that is consistent with having 2 counterfeits.

Repeat this until Weighter knows what the counterfeit coins are.

Weighter's goal is to minimize the number of weightings.

Spoiler's goal is to maximize the number of weightings.

Counterfeit Coins Problem: Project

Lets use AI on it!

How?

Set it up as a game. We look at $c = 2$.

Weighter picks two sets of x coins to balance.

Spoiler gives an answer that is consistent with having 2 counterfeits.

Repeat this until Weighter knows what the counterfeit coins are.

Weighter's goal is to minimize the number of weightings.

Spoiler's goal is to maximize the number of weightings.

Train two AI's (perhaps neural nets) to do well at these games and see if the number-of-weightings converges to a number that might be the truth.

Counterfeit Coins Problem: Project

Lets use AI on it!

How?

Set it up as a game. We look at $c = 2$.

Weighter picks two sets of x coins to balance.

Spoiler gives an answer that is consistent with having 2 counterfeits.

Repeat this until Weighter knows what the counterfeit coins are.

Weighter's goal is to minimize the number of weightings.

Spoiler's goal is to maximize the number of weightings.

Train two AI's (perhaps neural nets) to do well at these games and see if the number-of-weightings converges to a number that might be the truth.

Bonus You get to learn more about the classical versions of this problem.

The Two Numbers Game: Setting

The Two Numbers Game: Setting

Two Number Game begin with two numbers. Say $(12, 37)$.

The Two Numbers Game: Setting

Two Number Game begin with two numbers. Say $(12, 37)$.

Each player either

The Two Numbers Game: Setting

Two Number Game begin with two numbers. Say $(12, 37)$.

Each player either

subtracts one of the digits (1, 2, 3, or 5) from one of the numbers $(12, 35)$, or

The Two Numbers Game: Setting

Two Number Game begin with two numbers. Say (12, 37).

Each player either

- subtracts one of the digits (1, 2, 3, or 5) from one of the numbers (12, 35), or

- replace an even number x (12 here) with $x/2$ (6 here).

The Two Numbers Game: Setting

Two Number Game begin with two numbers. Say (12, 37).

Each player either

- subtracts one of the digits (1, 2, 3, or 5) from one of the numbers (12, 35), or

- replace an even number x (12 here) with $x/2$ (6 here).

First player to get one of the numbers ≤ 5 wins.

Example with (12, 35)

The Two Numbers Game: Setting

Two Number Game begin with two numbers. Say (12, 37).

Each player either

- subtracts one of the digits (1,2,3, or 5) from one of the numbers (12,35), or

- replace an even number x (12 here) with $x/2$ (6 here).

First player to get one of the numbers ≤ 5 wins.

Example with (12,35)

If ONE replaces 12 by 6 then (6,35). TWO then wins (6-3,35).

The Two Numbers Game: Setting

Two Number Game begin with two numbers. Say (12, 37).

Each player either

subtracts one of the digits (1, 2, 3, or 5) from one of the numbers (12, 35), or

replace an even number x (12 here) with $x/2$ (6 here).

First player to get one of the numbers ≤ 5 wins.

Example with (12, 35)

If ONE replaces 12 by 6 then (6, 35). TWO then wins (6-3, 35).

If ONE replace 12 with $12 - 2 = 10$, $12 - 3 = 9$, $12 - 5 = 7$ TWO wins.

The Two Numbers Game: Setting

Two Number Game begin with two numbers. Say (12, 37).

Each player either

- subtracts one of the digits (1, 2, 3, or 5) from one of the numbers (12, 35), or

- replace an even number x (12 here) with $x/2$ (6 here).

First player to get one of the numbers ≤ 5 wins.

Example with (12, 35)

If ONE replaces 12 by 6 then (6, 35). TWO then wins (6-3, 35).

If ONE replace 12 with $12 - 2 = 10$, $12 - 3 = 9$, $12 - 5 = 7$ TWO wins.

If ONE replaces 35 with $35 - 12 = 23$ then I DONT KNOW.

The Two Numbers Game: Setting

Two Number Game begin with two numbers. Say $(12, 37)$.

Each player either

subtracts one of the digits (1,2,3, or 5) from one of the numbers $(12,35)$, or

replace an even number x (12 here) with $x/2$ (6 here).

First player to get one of the numbers ≤ 5 wins.

Example with $(12,35)$

If ONE replaces 12 by 6 then $(6,35)$. TWO then wins $(6-3,35)$.

If ONE replace 12 with $12 - 2 = 10$, $12 - 3 = 9$, $12 - 5 = 7$ TWO wins.

If ONE replaces 35 with $35 - 12 = 23$ then I DONT KNOW.

For which (a, b) does ONE win? This has not been researched much.

The Two Numbers Game: First Assignment

The Two Numbers Game: First Assignment

Write a program that helps two humans play each other.

The Two Numbers Game: First Assignment

Write a program that helps two humans play each other.
PLAY each other and try to figure out some strategies.

The Two Numbers Game: First Assignment

Write a program that helps two humans play each other.

PLAY each other and try to figure out some strategies.

Write a program that plays the game. Put in as little or as much strategy as you want, but it has to make legal moves.

The Two Numbers Game: First Assignment

Write a program that helps two humans play each other.

PLAY each other and try to figure out some strategies.

Write a program that plays the game. Put in as little or as much strategy as you want, but it has to make legal moves.

PLAY against it and see how it does.

The Two Numbers Game: First Assignment

Write a program that helps two humans play each other.

PLAY each other and try to figure out some strategies.

Write a program that plays the game. Put in as little or as much strategy as you want, but it has to make legal moves.

PLAY against it and see how it does.

You will demo this. In the next bill-meeting, I will play against it. That demo should go smoothly **that is one of the reasons you have the non-bill meetings!**

The Two Numbers Game: First Assignment

Write a program that helps two humans play each other.

PLAY each other and try to figure out some strategies.

Write a program that plays the game. Put in as little or as much strategy as you want, but it has to make legal moves.

PLAY against it and see how it does.

You will demo this. In the next bill-meeting, I will play against it.

That demo should go smoothly **that is one of the reasons you have the non-bill meetings!**

Write a program that has the computer play itself.

Circle Crossing Game: First Project

Circle Crossing Game: First Project

Given a circle with n points on it. (You can assume they are equally spaced.)

Circle Crossing Game: First Project

Given a circle with n points on it. (You can assume they are equally spaced.)

ONE and TWO take turns picking two of the points and drawing a line between them.

Circle Crossing Game: First Project

Given a circle with n points on it. (You can assume they are equally spaced.)

ONE and TWO take turns picking two of the points and drawing a line between them.

ONE goes first.

Circle Crossing Game: First Project

Given a circle with n points on it. (You can assume they are equally spaced.)

ONE and TWO take turns picking two of the points and drawing a line between them.

ONE goes first.

First player to draw a line that crosses another line loses.

Circle Crossing Game: First Project

Given a circle with n points on it. (You can assume they are equally spaced.)

ONE and TWO take turns picking two of the points and drawing a line between them.

ONE goes first.

First player to draw a line that crosses another line loses.

Same programs as with the Number game.

Circle Crossing Game: First Project

Given a circle with n points on it. (You can assume they are equally spaced.)

ONE and TWO take turns picking two of the points and drawing a line between them.

ONE goes first.

First player to draw a line that crosses another line loses.

Same programs as with the Number game.

A human playing the game with another human or with the machine should be able to see the circle and the line. There should be a nice interface.

Counterfeit Coins Problem: Setting

Counterfeit Coins Problem: Setting

(We may consider this later.)

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

What is known?

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

What is known?

Very few exact values are known.

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

What is known?

Very few exact values are known.

Some asymptotic values are known.

Counterfeit Coins Problem: Setting

(We may consider this later.)

The following problem is well known:

There are 12 coins. One is counterfeit and is either heavier or lighter. You have a balance scale so you can take x coins on one side and x coins on the other and either it balances or one is heavier. How many weightings do you need to find which coin is counterfeit?

There are many variants on this problem:

Instead of 1 counterfeit coin, c counterfeit coins.

You may be given some information about the c counterfeits.

What is known?

Very few exact values are known.

Some asymptotic values are known.

There does not seem to be any math structure to use.

Counterfeit Coins Problem: Project

Counterfeit Coins Problem: Project

Lets use AI on it!

Counterfeit Coins Problem: Project

Lets use AI on it!

How?